

1 **Cryptic species and hybridisation in corals: challenges and opportunities for**
2 **conservation and restoration**

3
4

5	Cynthia Riginos ^{1,2,3}	0000-0002-5485-4197
6	Iva Popovic ¹	0000-0001-6582-4236
7	Zoe Meziere ¹	0000-0001-6874-8523
8	Vhon Oliver Garcia ¹	0000-0002-8933-452X
9	Ilha Byrne ¹	0000-0003-3909-2902
10	Samantha M Howitt ¹	0009-0002-6272-5261
11	Hisatake Ishida ¹	0000-0001-8401-8395
12	Kevin R Bairos-Novak ¹	0000-0002-0152-1452
13	Adriana Humanes ⁴	0000-0001-7962-2635
14	Hugo Scharfenstein ^{3,5}	0000-0003-3408-2350
15	Thomas Richards ¹	0000-0001-5945-6545
16	Ethan J. Briggs ¹	0000-0002-3475-5669
17	Vanessa Clark ¹	0000-0001-6125-9399
18	Chuan Lei ¹	
19	Mariam Khan ¹	
20	Katharine E Prata ¹	0000-0001-6679-0066

21
22 Corresponding author: Riginos

23
24 ¹ School of the Environment, The University of Queensland, St Lucia, QLD, Australia

25 ² Department of Marine Sciences – Tjärnö Marine Laboratory, University of Gothenburg,
26 Strömstad, Sweden

27 ³ Australian Institute of Marine Science, Townsville, QLD, Australia

28 ⁴ School of Natural and Environmental Sciences, Newcastle University, Newcastle upon Tyne,
29 UK

30 ⁵ School of BioSciences, The University of Melbourne, Melbourne, VIC, Australia

31
32 **Keywords**

33 cryptic species, adaptation, hybridization, coral, introgression, gene flow, conservation,
34 restoration, speciation

35

Abstract

shallow water coral reefs only ?

36 Conservation and management of coral reef ecosystems will depend on accurate assessments
37 of reef-building coral species diversity. However, the true diversity of corals may be
38 obfuscated by the presence of cryptic species, which are likely much more pervasive than is
39 currently recognised. Additionally, cryptic species may sometimes hybridize, resulting in gene
40 introgression between species. Here, we investigate the prevalence of cryptic coral species
41 via a structured literature review and find that over 50% of population genomic studies show
42 evidence for divisions within taxonomically recognised species and that such closely-related
43 taxa are often linked by gene flow. We find that cryptic taxa frequently segregate by
44 environment, especially depth, and may differ by phenotypic characteristics including
45 resilience to heat stress. This hidden biodiversity creates challenges for coral conservation
46 and restoration planning that are not well appreciated, including hiding true population
47 declines, biasing estimates for species' phenotypic breadth, overestimating the resilience of
48 species to stressors, yielding uncertainty in evolutionary dynamics inferred from past studies,
49 and creating reproductive barriers that may limit mating between local and translocated
50 corals. Increasing awareness that coral cryptic species with incomplete species boundaries
51 are common and building this expectation into conservation and restoration plans is an
52 important pathway forward. Rich opportunities for interdisciplinary collaboration among
53 coral and speciation biologists could fill key knowledge gaps relevant to conservation. We
54 detail recommendations for best practice and strategies for identifying cryptic taxa and
55 hybrids and urge their consideration in all future studies on corals.

remove "gene"
or replace by
genetic, as
genes are not
the only
elements that
introgress

56


57 **1 Introduction: Hidden dimensions of coral biodiversity pose conservation** 58 **challenges**

59 Coral reefs are highly biodiverse and productive ecosystems (Reaka-Kudla, 1997; Fisher *et al.*,
60 2015) that substantially contribute to human well-being (Moberg & Folke, 1999; Adey, 2000).
61 Yet, reef-building corals are imperilled by rising temperatures and other anthropogenic
62 stressors worldwide (Hughes *et al.*, 2017; Knowlton *et al.*, 2021; Souter *et al.*, 2021). Thus,
63 there is great urgency to inventory coral biodiversity and to deepen knowledge regarding the
64 processes that create and maintain this diversity to guide conservation and restoration
65 actions.

66

67 For corals it has long been recognised that morphological variation is unlikely to align well
68 with genetic variation. Molecular based phylogenetic investigations are uncovering
69 unexpected relationships and unanticipated numbers of distinct taxa (reviewed by Kitahara
70 *et al.*, 2016; Cowman *et al.*, 2020), and similarly, multilocus population genetic surveys
71 routinely find distinct genetic taxa within morphologically defined taxonomic species. These
72 so-called *cryptic species* (recent reported genomic examples include: Gomez-Corrales &

73 Prada, 2020; Underwood *et al.*, 2020; Wepfer *et al.*, 2020; Afiq-Rosli *et al.*, 2021; Bongaerts
74 *et al.*, 2021a; Feldman *et al.*, 2021; Fifer *et al.*, 2021; Rippe *et al.*, 2021; Zayasu *et al.*, 2021;
75 Adam *et al.*, 2022; Prata *et al.*, 2022; Rivera *et al.*, 2022; Matias *et al.*, 2023; Voolstra *et al.*,
76 2023; Meziere *et al.*, 2024) reinforce observations previously made with microsatellites and
77 few nuclear markers (such as Bongaerts *et al.*, 2010; Souter, 2010; Ladner & Palumbi, 2012;
78 Schmidt-Roach *et al.*, 2012; Prada & Hellberg, 2013; Boulay *et al.*, 2014; Prada *et al.*, 2014;
79 Warner, van Oppen & Willis, 2015; Gélín *et al.*, 2017) and demonstrate how species
80 designations based on morphology alone can underestimate true community diversity (Fig 1).
81 Indeed, most conservation management and restoration plans implicitly assume that coral
82 species are recognisable and biologically valid entities (Baums, 2008; Anthony *et al.*, 2017;
83 National Academies of Sciences, 2018; Colton *et al.*, 2022). These assumptions may
84 unintentionally bias conservation and restoration efforts when such biodiversity is not
85 acknowledged. Yet, detecting and delineating distinct genetic groups presents a substantial
86 challenge (which we discuss in more detail in Section 2.1). Rather than adopting a specific
87 criterion for species status, we use the terms *genetic groups* or *taxa* to signify their genetic
88 distinctiveness. We avoid terms such as clade or lineage, as monophyly across all or many
89 gene trees is unlikely when divergence is recent. We use *cryptic* to signify that taxa appear
90 morphologically similar to the “untrained human eye”, at least under field conditions.



to be even more convincing, suggest to add photographs taxonomic characters such a structure, to show the read th just a matter of amateurs loo morphology

91
 92 **Figure 1** – Examples of closely related cryptic species. Details can be found in the original studies:
 93 *Stylophora pistillata* (Meziere *et al.*, 2024), *Porites* sp. (Starko *et al.*, 2023), and *Agaricia lamarcki*
 94 (Prata *et al.*, 2022).

95
 96 Further complicating taxonomic delineations, cryptic coral taxa may be linked by occasional
 97 gene flow, or hybridisation. Morphologically intermediate individuals are often encountered
 98 in the field (Veron, 1995; Richards *et al.*, 2008), and many species can be crossed at least
 99 under experimental conditions (Isomura *et al.*, 2016; Chan, Peplow & van Oppen, 2019;
 100 Kitanobo *et al.*, 2022). Genomic studies confirm that hybridisation is common in corals, where
 101 gene flow has been documented between distinct taxa (Cooke *et al.*, 2020; Fifer *et al.*, 2021;
 102 Prada & Hellberg, 2021; Rippe *et al.*, 2021; Prata *et al.*, 2022; Matias *et al.*, 2023; Starko *et al.*,
 103 2023; Zhang *et al.*, 2023; Meziere *et al.*, 2024), including historical gene flow between taxa
 104 separated by over 15 million years (reviewed by van Oppen & Gates, 2006; Willis *et al.*, 2006;
 105 Mao & Satoh, 2019; González, Rivera-Vicéns & Schizas, 2021; Hobbs *et al.*, 2021; Pinsky, Clark
 106 & Bos, 2023). Cryptic genetic groups or taxa, potentially connected by limited gene flow,
 107 present distinct challenges for understanding coral biodiversity. Failing to recognize cryptic
 108 coral taxa and appropriately adjust interpretations can result in misleading conclusions about
 109 fundamentally important aspects of biodiversity measurement, including: i) underestimating
 110 species diversity, ii) overestimating within-species variances for various traits and tolerances,

111 iii) biasing inferences regarding within-species patterns of genetic diversity and population
112 structure, and iv) invalidating assumptions about reproductive compatibility.

113

114 First, biodiversity inventories typically determine the count and abundances of distinct
115 species. Common methods, such as field surveys, are usually based on morphological
116 identification of live organisms. If distinct taxa are not appropriately recognised and
117 delineated, then total species counts will be greatly underestimated, presenting an unstable
118 base from which to draw conclusions about coral population sizes, species extinction
119 vulnerabilities, and biodiversity valuations. For example, estimates of census population sizes
120 for present-day corals have been recently debated (Dietzel *et al.*, 2021; Muir *et al.*, 2022), as
121 smaller population sizes would imply greater vulnerability to extinction (Dietzel *et al.*, 2021;
122 Muir *et al.*, 2022).

123

124 Second, assessment of trait diversity is compromised if cryptic taxa are not identified, and
125 trait measurement occurs on an amalgamation of distinct taxa. The agglomerate will contain
126 more variation than the distinct groups, which will inflate estimates of both genetic and
127 phenotypic variance. This phenomenon can affect interpretation for a diverse array of traits.
128 For example, species range estimates would be upwardly biased if cryptic species are
129 geographically restricted within the broader range of a morphospecies. Environmental niche
130 breadth can also be overestimated: for instance, many coral species that were previously
131 considered **depth** generalists could resolve into taxa with more restricted depth distributions
132 when cryptic species are considered (e.g., Bongaerts *et al.*, 2021a). Similarly, a presumed
133 species may appear to have a generalist phenotype but actually comprise multiple taxa that
134 are more specialized, as appears to be the case for bleaching responses among *Orbicella*
135 *faveolata* taxa (Gomez-Corrales & Prada, 2020). In summary, single-species conclusions
136 drawn from agglomerated data from multiple cryptic taxa are likely to be overly optimistic in
137 terms of coral resilience.

138

139 Third, common measures of population genetics and gene flow may be inherently biased and
140 incorrect when cryptic taxa are examined as one. For example, using allele frequencies to
141 make inferences about microevolutionary dynamics without delineating separate gene pools
142 can yield incorrect results and interpretations. Lumping genetically distinct groups will inflate
143 apparent within-population diversity (based on measures of allelic diversity or expected
144 heterozygosity) and will therefore cause inbreeding to be overestimated – known as the
145 Wahlund effect (as discussed in Schmidt, Thia & Hoffmann, 2023). When comparing across
146 geographic locations, measures of population structure (notably F_{ST}) can be biased either
147 upwards or downwards, depending on the particular mix of cryptic taxa sampled. These
148 phenomena are neatly illustrated and discussed by Sheets *et al.* (2018) for *Acropora*
149 *hyacinthus* in the western Pacific (see also Pante *et al.*, 2014; Warner *et al.*, 2015).
150 Additionally, common metrics of population genetics such as genetic diversity,
151 differentiation, inbreeding and effective population size may be biased when hybridization

152 between differentiated taxa is not taken into account (Hoban *et al.*, 2022). For example,
153 hybridisation may inflate the measured diversity for populations that include individuals of
154 mixed ancestry, while gene flow or dispersal inferences may be biased in either direction.
155 Together, cryptic taxa and hybridisation are likely to greatly affect the accuracy of studies
156 aiming to assess and monitor genetic diversity.

157

158 Finally, the existence of cryptic taxa implies that there is some form of reproductive isolation
159 between taxa. Intrinsic barriers such as incompatible loci are likely to reduce gene flow among
160 taxa in nature, so populations of a morphospecies may not be as strongly linked by gene flow
161 as is often assumed. In restoration contexts, intrinsic barriers could lead to translocated
162 individuals being unable to mate with local individuals and could prevent crossing in captivity
163 and/or reduce hybrid fitness (outbreeding depression). Extrinsic barriers to reproduction
164 arise when survival or fertility differs between taxa based on their surrounding environments,
165 and as a result, they inhabit different niches. This means that cryptic taxa may not be
166 ecologically or functionally equivalent. Restoration actions that involve coral outplanting (i.e.,
167 fragments or sexually propagated colonies) risk mis-matching the source taxon's niche with
168 their new destination's environmental conditions. Conversely, hybridisation between taxa
169 could enhance fitness or phenotypic breadth of resultant offspring (Chan *et al.*, 2018).

170

171 Figure 2 outlines some possible consequences of cryptic coral taxa to current restoration
172 actions such as direct transplantation and coral gardening, as well as actions that are being
173 actively researched for potential use in coral restoration (National Academies of Sciences,
174 2018; Anthony *et al.*, 2020; Hein *et al.*, 2020; Bay *et al.*, 2023).







175

Interventions

- Direct transplantation & coral gardening:**
Transplantation of colonies or fragments from one reef to another. Coral gardening involves rearing corals in a nursery setting (land or sea-based) before transplantation to the reef.
- Gamete and larval capture and seeding:**
Collecting gametes and larvae from the wild and releasing larvae onto other reef areas.
- Intraspecific breeding within reefs:^a**
Sexual reproduction of targeted colonies and releasing progeny to increase population size (supportive breeding) and/or increase frequency of heat tolerant alleles (selective breeding).
- Assisted gene flow:^b**
Human assisted movement, with or without crossing corals in captivity, to promote gene exchange beyond the natural dispersal scale but within the current species range.
- Assisted migration:**
Human assisted movement, with or without crossing corals in captivity, over large spatial scales within or beyond the current species range.
- Hybridization among species:**
Crossing species to create novel genotypes and phenotypes.
- Conditioning:**
Stress exposure to make corals more tolerant of future stresses.
- Algal-symbiont manipulation:^c**
Replacing algal symbionts with more heat tolerant strains, that may be sourced from a reef or experimentally evolved
- Probiotics:^d**
Supplementing an established community of microbes with beneficial strains.
- Gene-editing:**
Altering select genes to yield more heat tolerant phenotypes

Equivalent terms for interventions:
^aManaged selection for environmental tolerance
^bIntraspecific breeding between reefs; outcrossing within species
^cExperimental evolution
^dMicrobiome manipulation

Potential consequences arising from cryptic species or hybridisation

- Taxon-environment mismatch**

Cryptic taxa differ in niche and other phenotypic attributes such that transplanted fragments, colonies or host genotypes that do not match the local environment perform poorly. This may include different responses to stress.
- Reproductive isolation**

Mating success between cryptic taxa is reduced relative to same-taxon matings. This can prevent transplants from mating with local colonies and can diminish crossing success for interventions that rely on sexual reproduction.
- Competitive dominance of transplanted taxon**

If the transplanted taxon has greater growth, survival, reproductive output, etc. then the transplanted taxon may outcompete the local taxon, potentially leading to the loss of locally adapted genetic diversity. However, increased coral cover may be viewed as a net benefit.
- Competitive dominance of hybrids**

Hybrids are expected to show more extreme phenotypes than their parental taxa: if hybrid colonies have greater growth, survival, reproductive output, etc. then outplanted hybrids may outcompete the local taxon, potentially leading to the loss of locally adapted genetic diversity. However, increased coral cover may be viewed as a net benefit.
- Host-microbe mismatching**

Beneficial microbes, including dinoflagellate symbionts, may confer benefits to some cryptic host taxa but not others.
- Gene-manipulation mismatching**

Edited genes may result in alleles that benefit some cryptic host taxa but not others.

176

177
178
179
180
181
182
183
184

Figure 2 - How cryptic coral taxa and hybridisation could affect coral reef restoration actions that aim to preserve biological diversity, counter population declines, and promote resilience to climate change through biological adaptation. Terminologies follow (van Oppen *et al.*, 2015; National Academies of Sciences, 2018; Hein *et al.*, 2020; Bay *et al.*, 2023) and are not necessarily mutually exclusive. For example, assisted migration could be undertaken on fragments (direct transplantation), larvae, or via the progeny of captive, sexually propagated corals. In all graphics, the coral colony shown as a larger size indicates greater fitness (survival, reproductive output, thermal tolerance, etc) leading to competitive dominance.

185
186
187
188
189
190
191
192
193
194

As outlined above, cryptic taxa and hybridisation have substantial implications for understanding and predicting the ecological and evolutionary dynamics of corals, and thus will be essential for making informed conservation and restoration decisions into the future. In this review, we critically assess the prevalence of hidden biodiversity among corals. By corals, we refer to benthic Anthozoans including Scleractinians (hard corals) as well as Octocorallians (soft corals). We focus primarily on population *genomic* studies (supported by whole genome or reduced representation sequencing of genome-wide variation) because thousands of loci are often required to detect recent evolutionary delineations between taxa. This paper builds upon important earlier reviews and syntheses of cryptic taxa (Richards, Berry & van Oppen, 2016) and hybridisation in corals (van Oppen & Gates, 2006; Willis *et al.*,

195 2006; Hobbs *et al.*, 2021; Pinsky *et al.*, 2023) that primarily focused on studies with few
196 markers (microsatellites, few sequenced loci, or allozymes).

197

198 Here we explore how evolutionary processes in corals can and should shape conservation and
199 restoration priorities and actions. Specifically, we: 1) develop criteria for recognising cryptic
200 taxa from population genetic data, 2) describe and evaluate surveyed genomic literature
201 against these guidelines to show that cryptic taxa frequently exist in sympatry, 3) determine
202 whether cryptic divergence in cnidarian hosts is accompanied by divergence in their
203 photosynthetic symbionts, 4) explore the environmental factors segregating cryptic taxa, 5)
204 review examples of hybridisation and/or gene flow between taxa, and 6) discuss how corals
205 could be exciting model systems for speciation and adaptation studies. The main text is
206 supported by text boxes that, A) provide a worked example of delineating coral taxa, B)
207 demonstrate how cryptic taxa are commonly overlooked in coral experiments, and C) outline
208 best practices for designing studies when cryptic taxa are likely to be encountered.

209 **2 Closely related coral taxa are common in sympatry and frequently connected by** 210 **gene flow**

211 To gauge the prevalence and impacts of cryptic coral taxa and hybridisation, we evaluate
212 findings from a structured literature search alongside a broader literature review. Although
213 our primary intent is to draw inferences from more informative genome-based studies,
214 population genomic studies for corals are limited both in their number and in scope. Thus, we
215 also include some discussion of noteworthy results based on microsatellite surveys and also
216 review microbial symbiont diversity because of their relation to cryptic host taxa (under
217 Sections 2.2 and 2.3).

218

219 **2.1 Detecting cryptic coral species using criteria based on reproductive isolation**

220 Many coral taxa potentially inhabit the “grey zone” of the speciation continuum (de Queiroz,
221 2005; Roux *et al.*, 2016). That is, genetically distinct groups may coexist within dispersal range,
222 yet these groups may not have all the contingent properties of distinct ‘species’ (sensu de
223 Queiroz, 2005), including complete reproductive isolation. However, for groups with
224 overlapping ranges to remain genetically distinct, reproductive barriers of some form must
225 be present. Reproductive barriers between groups may occur due to differences in
226 microhabitats or spawning times resulting in infrequent fertilisation, producing offspring with
227 phenotypes that do not match microhabitats, or other **intrinsic genetic incompatibilities**.
228 When reproductive barriers are not complete, the outcome of interbreeding between groups
229 is controlled by the relative strength of selection (which promotes divergence), and the scale
230 of gene exchange (which promotes homogenisation). Even low levels of gene exchange
231 between otherwise distinct groups will cause alleles to be shared. Allele sharing will also be
232 prevalent if divergence time between groups is recent whereby groups share ancestral

this paragraph would benefit from some
key references

233 variation. Thus, allele sharing is expected to be common between cryptic coral taxa that are
234 recently diverged (with or without ongoing gene flow).

235

236 Population genomic surveys have the power to detect subtly differentiated genetic groups
237 where allele frequency differences between groups are small. The statistical power of
238 common methods is determined by the number of loci examined as well as the extent of
239 genetic covariance among loci, where distinct populations exhibit non-random associations
240 of alleles at various loci across the genome (i.e., *linkage disequilibria*). While physical linkage
241 on a chromosome alone will cause covariance among loci, genome-wide covariance also
242 arises as a direct consequence of population structure (reflecting distinct gene pools subject
243 to independent outcomes of genetic drift and selection). For example, reproductively isolated
244 taxa or geographically separated populations will be differentiated across loci due to genetic
245 drift and selection. In contrast, gene flow among them will erode both their allele frequency
246 differences and linkage disequilibria. Statistical power for identifying small allele frequency
247 differences and linkage disequilibria in empirical surveys is increased by sampling many
248 individuals and many loci, where subtle differences between groups can be missed when few
249 individuals or few loci are sampled. For example, two co-occurring genetic groups of
250 *Montastraea cavernosa* were clearly delineated using thousands of loci and yet ambiguous
251 with 9 microsatellite loci (Sturm *et al.*, 2020).

252

253 We focus on two general approaches for **detecting subtle differentiation** between genetic
254 groups: **ordination-based analyses and model-based clustering**. Ordination-based analyses,
255 such as principal components analyses (PCA), principal coordinates analysis,
256 multidimensional scaling, etc., describe multidimensional relationships among entities and
257 are based on (and visually represent) the genetic covariance matrix (Patterson, Price & Reich,
258 2006). Model-based clustering analyses – typified by admixture detection analyses such as
259 STRUCTURE (Pritchard, Stephens & Donnelly, 2000) and ADMIXTURE (Alexander, Novembre
260 & Lange, 2009) – partition groups (K) based on associations among alleles and loci. Ordination
261 and model-based clustering approaches are unsupervised machine learning methods that are
262 valuable for exploring relationships between individuals (that is, without pre-assigning
263 individuals to “populations”, as is required by F-statistics and other population-level metrics).
264 These methods are some of the most common and routinely employed methods in genomic
265 surveys and provide complementary insights into spatial patterns of genetic diversity. We
266 omit results from **supervised methods, such as discriminant analysis**, that maximize variance
267 using user-assigned groupings (see **Thia, 2022 for extended discussion**).

268

269 We define taxa as distinct groups of individuals (genotypic clusters) within named species that
270 maintain their distinctiveness even when their ranges overlap, therefore they have the

271 opportunity to mate (in line with Mallet, 1995). This definition aligns to many species
272 concepts (de Queiroz, 2005), under the assumption that some form of reproductive isolation
273 must be occurring. Instances where distinct genetic groups are found together (i.e., sympatric
274 within the scale of dispersal distance) provide the strongest circumstantial evidence for some
275 degree of reproductive isolation between groups, as reduced gene flow due to dispersal
276 restrictions cannot be the primary cause of genetic divergence in these cases. These “taxa”
277 might be considered species, depending on the definition of species employed (de Queiroz,
278 2005); as our focus is not taxonomy but evolutionary processes and their implications, the
279 term “taxon” is more flexible. Thus, we focus on evidence for *sympatric cryptic coral taxa*.
280 When distinct genetic groups are geographically separated (i.e., allopatric), then divergence
281 may solely reflect dispersal barriers to gene exchange and therefore are not informative for
282 inferring reproductive isolation.

283

284 We propose three requirements for identifying and delineating taxa using ordination and
285 model-based clustering:

- 286 1) Distinct genetic groups occur in sympatry relative to their dispersal ability.
- 287 2) Ordination (e.g., PCA) clusters these distinct genetic groups based on
288 genotypes of individuals and/or model-based clustering indicates that
289 individuals belong to separate groups.
- 290 3) When genetic distance between sympatric individuals is greater than the
291 genetic distance between allopatric individuals. This is evidenced by
292 divergence between groups across the primary ordination axis and/or at low
293 hypothesized group numbers (K values) as compared to axes or groups that
294 describe geographic structure.

295

296 To determine if published genomic surveys of corals typically test for and find evidence for
297 cryptic taxa following the above criteria (see Textbox 1 for worked example), we searched the
298 Web of Science for published papers displaying graphical results from ordination (PCA,
299 primarily) and/or model-based clustering analyses (Structure and ADMIXTURE). We focus on
300 studies that genotyped individual cnidarian genomes for 1000s of loci or more and sampled
301 from two or more geographic locations (see Appendix for details of literature search). We
302 additionally collected information on depth ranges and symbiont composition where
303 reported. The literature search uncovered 41 papers describing results for 31 species, where
304 some papers included multiple species and some species were studied across multiple papers,
305 yielding a total of 51 unique records (available as supplemental data). Although we did not
306 restrict the search by sampling depth, none of the recovered records included species deeper
307 than mesophotic depths.

308


309 In our review of published studies, we apply the above criteria (1-3) conservatively by only
310 looking for sympatric differentiation i) along the first principal component axis (referred to as
311 'PC1') in ordination analyses, and ii) for model-based clustering examining outcomes when
312 individuals were assigned to only two groups (K=2). We consider individuals sampled ≤ 10 km
313 apart as being broadly sympatric since dispersal distances for many coral species are
314 unknown.

315

316 As shown in Fig. 3, over half of the coral population genomic studies (23 records out of 39
317 that could be evaluated against all three criteria) showed evidence for distinct cryptic taxa.
318 While we might expect genetic differentiation to be greater among brooding corals that have
319 less innate dispersal ability relative to broadcast spawning corals (Knowlton, 2001), the
320 relative proportions of sympatric versus non-sympatric groups did not differ between
321 brooding and broadcast spawning corals ($P > 0.2$ with Fisher's exact test using either
322 ordination or clustering). Criterion 3 is based on genetically distinct groups co-occurring. This
323 evidence is stronger when this groups co-occur across many geographic locations. Thus, we
324 investigated whether pairs of genetically distinct coral groups were repeated across multiple
325 sites using model-based clustering results based on author-selected K values. Across studies,
326 it was common for cryptic groups to be sympatric at multiple sampled sites (Fig. 3). This
327 observation strengthens the conclusion that closely-related, but distinct taxa, can co-occur
328 over extensive geographic areas and implicates some degree of reproductive isolation
329 maintaining the distinctiveness of each group. In summary, cryptic taxa are common features
330 in population genetic studies of corals.

331

332 Our literature review shows that cryptic taxa are prevalent, implying that many conclusions
333 related to biodiversity, species traits, and population genetic structure based at the
334 morphospecies level are likely to be inaccurate based on the reasons discussed previously
335 (see Section 1). Even when genomic data to evaluate cryptic taxa are presented, five studies
336 that we assessed failed to include any mention of cryptic species or taxa, despite their results
337 showing evidence for genetically distinct groups. Additionally, some of the studies reviewed
338 here computed summary statistics such as heterozygosity, or F-statistics using all individuals
339 from the sampling locations, despite evidence for cryptic taxa co-occurring within locations.
340 We surmise that inferences regarding the absence of potential cryptic taxa based on analyses
341 using few loci (including microsatellite studies) are likely lacking in terms of the power to
342 detect recently differentiated taxa, and therefore many published studies – including studies
343 published by authors of this review – may inadvertently base conclusions on heterogeneous
344 mixes of cryptic taxa.



345
346
347
348
349
350
351
352
353
354
355

Figure 3 - For many genomic studies of corals, the greatest axis of genetic differentiation defines groups that are sympatric. Results by species are summarized as either meeting or not meeting the three criteria for cryptic taxa under the strictest definition where sympatric differentiation is aligned to the first axis (ordination) or K = 2 (model-based clustering). Some studies had both ordination and model-based clustering, and some species were included in more than one study, thus multiple points can appear against each species. Studies that did not present ordination or model-based clustering are not shown. The number of locations where distinct groups were sympatric was based on K clustering, using the K value preferred by the authors, where a dash indicates that this value could not be inferred. From the 51 studies examined, 39 presented results that could be evaluated against criteria 1-3.

356

357 **2.2 No clear patterns for symbionts associated with cryptic coral host taxa**

358 An important aspect of coral biodiversity is the diverse microbial community living within the
359 cnidarian host, where mutualistic relationships with endosymbiotic dinoflagellates from the
360 family Symbiodiniaceae are known to affect whole organism physiology (LaJeunesse *et al.*,
361 2018). The specificity of coral-symbiont pairings is influenced by host-symbiont genetic
362 interactions, host reproduction, symbiont transfer mode, and environment (see Baker, 2003;
363 Davies *et al.*, 2023; Turnham *et al.*, 2023 for further readings). While symbiont evolution is
364 not the focus of this review, joint consideration of coral hosts and their microbes is relevant
365 in understanding the specificity of associations that can affect traits, local adaptation, and
366 restoration actions.

367

368 Among the population genomic studies that we surveyed, 43% genotyped dinoflagellate
369 symbionts alongside the coral hosts. Several studies (e.g., Howells *et al.*, 2016; van Oppen *et al.*,
370 2018; Gomez-Corrales & Prada, 2020; Bongaerts *et al.*, 2021a; Prata *et al.*, 2022; Buitrago-
371 Lopez *et al.*, 2023; Starko *et al.*, 2023) used internal transcribed spacer (*ITS*) sequences to
372 characterize the community of symbiont lineages living within a colony, aligning with
373 traditional marker-based methods (Davies *et al.*, 2023) for symbiont characterisation. Some

374 studies also used incidentally recorded symbiont sequences retrieved from whole-colony
375 sequencing (either reduced representation or shotgun whole genome sequencing) to make
376 inferences about symbionts, including reconstructing organelle diversity (Bongaerts *et al.*,
377 2017; Forsman *et al.*, 2017; Gonzalez-Zapata *et al.*, 2018; Cooke *et al.*, 2020; Bongaerts *et al.*,
378 2021a; Matias *et al.*, 2023; Zhang *et al.*, 2023) or summarizing symbiont genomes with *k*-mer
379 analyses (Zhang *et al.*, 2023).

380
381 A central issue for coral adaptation and diversification is the extent to which symbiotic
382 associations are flexible. Brooding corals are expected to have less flexible symbiont
383 associations because they often exhibit vertical symbiont transmission from the maternal
384 colony (Knowlton, 2001; Johnston, Cunning & Burgess, 2022; Turnham *et al.*, 2023). However,
385 in our review, only one brooding coral (*Stylophora pistillata*) that was genotyped for both
386 symbionts and hosts (Buitrago-Lopez *et al.*, 2023) preventing adequate comparisons among
387 reproductive modes. Symbiont types were commonly found to be shared across cryptic host
388 lineages for *S. pistillata* (Buitrago-Lopez *et al.*, 2023), for broadcast spawners (*O. faveolata*:
389 Gomez-Corrales & Prada, 2020; *Pachyseris speciosa*: Bongaerts *et al.*, 2021a; *Pocillopora*
390 *verrucosa*: Buitrago-Lopez *et al.*, 2023; *Porites lobata*: Rivera *et al.*, 2022; *Porites sp.*: Starko
391 *et al.*, 2023) for mixed mode *Pocillopora damicornis* (van Oppen *et al.*, 2018), and for taxa
392 with unknown modes of reproduction (*Agaricia fragilis*: Bongaerts *et al.*, 2017; *Agaricia*
393 *lamarcki*: Prata *et al.*, 2022). Some symbiont types appeared specific to cryptic taxa (*O.*
394 *faveolata*: Gomez-Corrales & Prada, 2020; *P. speciosa*: Bongaerts *et al.*, 2021a; *P. verrucosa*:
395 Buitrago-Lopez *et al.*, 2023; massive *Porites sp.*: Starko *et al.*, 2023; *P. damicornis*: van Oppen
396 *et al.*, 2018), whereas other sets of cryptic taxa were not associated with specific symbiont
397 types (*Agaricia fragilis*: Bongaerts *et al.*, 2017; *Agaricia lamarcki*: Prata *et al.*, 2022; *S.*
398 *pistillata*: Buitrago-Lopez *et al.*, 2023). Thus, some symbiont strains may be specialized, as
399 evidenced by symbiont lineages only associating with one host group; however, this pattern
400 is not always a characteristic accompanying cryptic cnidarian taxa.

401
402 Moving beyond single marker genotyping of symbionts (Davies *et al.*, 2023; Ishida *et al.*, 2023;
403 Zhang *et al.*, 2023) may provide better resolution of host-symbiont associations. For example,
404 Rivera *et al.* (2022) found that symbiont identities among *P. lobata* did not align to host taxa
405 using ITS genotyping but were concordant when using higher resolution SNP-based analyses.
406 Additional studies are needed, however, to determine whether alternative methods
407 harnessing whole-genome data to summarize symbiont diversity will reveal more nuanced
408 patterns than ITS based results.

409 **2.3 Depth and microhabitats can segregate cryptic taxa**

411 For coral hosts, the existence of distinct genetic groups in sympatry implies that
412 differentiation may be maintained by strong natural selection arising from local environments
413 (Richardson *et al.*, 2014). Such strong selection and genotype-by-environment associations
414 may merely reflect pre-existing reproductive isolation (Bierne *et al.*, 2011), although

415 divergent natural selection can also contribute to reproductive isolation (Schluter, 2001;
416 Rundle & Nosil, 2005). For marine species, including corals, depth often delineates closely-
417 related species (Knowlton, 1993) and could drive ecological speciation in corals (González *et*
418 *al.*, 2020). In coral reef communities, depth is a predominant structuring aspect, with distinct
419 species turnover between shallow (approximately < 30 m) and mesophotic (approximately
420 30-150 m) **depth zones**. **Importantly**, these transitions exist across communities because
421 many environmental factors covary with depth, such as light intensity and spectrum (Lesser,
422 Slattery & Leichter, 2009), temperature (Kahng *et al.*, 2019), nutrients (Leichter, Stokes &
423 Genovese, 2008), water flow (Muir *et al.*, 2015), as well as disturbance frequency and severity
424 (Bongaerts & Smith, 2019), creating highly contrasted habitats often only meters apart. The
425 strongest environmental differences are within the first few meters due to the exponential
426 decay of light in the aquatic environment – the most important environmental factor for light-
427 dependent scleractinians. Thus, it is highly likely that recently diverged taxa will also be
428 partitioned by depth, matching patterns of species turnover.

429
430 Among the population genomic studies surveyed, cryptic taxa differed in abundance by
431 depth-associated habitats. Eleven studies to date undertook extensive sampling over
432 replicated depth-associated habitat contrasts (i.e., more than one site >10 km apart with two
433 depth habitats sampled within each site). Replicated differentiation by depth was found for
434 *Agaricia fragilis* (i.e., shallow vs. mesophotic, Bongaerts *et al.*, 2017), *Agaricia lamarcki* (i.e.,
435 shallow vs. mesophotic, Prata *et al.*, 2022), *Isopora brueggemanni* (i.e., lagoon vs. slope;
436 Thomas *et al.*, 2019), *Pocillopora damnicornis* (i.e., flat vs. slope, van Oppen *et al.*, 2018) and
437 *Montastraea cavernosa* (i.e., shallow vs. mesophotic, Sturm *et al.*, 2022) but not for *Agaricia*
438 *grahamae* (i.e., upper vs. lower mesophotic, Prata *et al.*, 2022), *Stephanocoenia intersepta*
439 (i.e., shallow vs. mesophotic, Bongaerts *et al.*, 2017), *Acropora digitifera* (i.e., lagoon vs. slope,
440 Thomas *et al.*, 2019), ***Agaricia undata*** (i.e., shallow vs. upper mesophotic and upper vs. lower
441 mesophotic, Gonzalez-Zapata *et al.*, 2018). Thus, differentiation by depth frequently
442 discriminates cryptic coral taxa.

443
444 The most extensive investigations of differentiation by depth have been for *Seriatopora*
445 *hystrix* (brooder) on the Great Barrier Reef, Australia and *Eunicea flexuosa* (broadcast
446 spawner) in the Caribbean. Although most of the studies discussed below pre-date genomic-
447 scale genotyping and thus rely on small numbers of microsatellite loci for making genetic
448 inferences, they are noteworthy because they meticulously documented small scale depth
449 and habitat segregation of host genotypes and also used reciprocal transplantation
450 experiments to further support evidence for depth-associated adaptive differentiation.
451 Bongaerts *et al.* (2010) showed that genetically distinct groups of *S. hystrix* were associated
452 with reef zone and depth position (backreef at 2 m, shallow reef at 6 m, deep reef at ~27 m;
453 locations within 200 m distant horizontally). **For the gorgonian *E. flexuosa*, Prada & Hellberg**
454 **(2013) demonstrated that individuals from <5 m and >20 m depths were distinct and joined**
455 **by a narrow hybrid zone (< 200 m horizontally).** This depth separation was likely maintained

456 by selection against juveniles (Prada & Hellberg, 2014). Reciprocal transplantation supported
457 evidence for local adaptation in both cases (Bongaerts *et al.*, 2011; Prada & Hellberg, 2013).
458 Together, these two examples (*S. hystrix* and *E. flexuosa*) show that cryptic coral taxa could
459 comprise distinct depth-specific ecotypes maintained by selection and furthermore illustrate
460 the utility and feasibility of using experiments to strengthen evidence for local adaptation
461 across small spatial scales. There is an unrealised opportunity to combine future genomic
462 investigations with manipulative experiments to advance the understanding of speciation and
463 adaptation (discussed further in Section 3).

464


465 While depth has been one of the most investigated environmental axes of local genetic
466 differentiation in corals, cryptic taxa can be associated with other habitat differences (and
467 sometimes are evident with few markers (Warner *et al.*, 2015)). In one example, broad
468 physiological differences between *A. hyacinthus* taken from two reef pools less than 500m
469 apart but varying in natural thermal variability (coined “highly” and “moderately” variable
470 pools (reviewed by Thomas *et al.*, 2018)); were partially explained by the relative abundance
471 of four cryptic species (identified using ordination of SNP-based genotypes) that differed in
472 abundance between the two pools (Rose *et al.*, 2017). Similarly, substantial differences in
473 skeletal morphologies and gene expression were found between *P. lutea* colonies sampled
474 from a mangrove lagoon versus colonies from a reef approximately 500 m away (Scucchia *et al.*,
475 2023); however, these phenotypic differences also aligned with genome-wide
476 differentiation measured by transcriptome-derived SNPs, suggesting that mangrove and reef
477 populations could simply be different taxa (applying the criteria in Section 2.1).

478

479 It is important to note that the structure and composition of coral-associated microbial
480 communities also often vary along environmental gradients, including those associated with
481 depth and disturbance (Klaus *et al.*, 2007; Bongaerts *et al.*, 2013; Howells *et al.*, 2013; Quigley
482 *et al.*, 2022), and likely help mediate coral holobiont adaptation to environmental stress.
483 Endosymbiotic dinoflagellates (LaJeunesse *et al.*, 2018) enable certain coral species to reside
484 within specific light environments, where mesophotic and shallow depth coral hosts often
485 harbour distinct symbiont strains, whereas depth-generalists appear to be able to host many
486 symbiont strains (Bongaerts *et al.*, 2013). Consistent with these previous findings, several of
487 the studies examined here reported greater spatial or environmental partitioning among
488 symbionts as compared to hosts (e.g., *Astrangia poculata*: Aichelman & Barshis, 2020; *P.*
489 *verrucosa*: Buitrago-Lopez *et al.*, 2023; *S. pistillata*: Buitrago-Lopez *et al.*, 2023; *Platygyra*
490 *daedalea*: Howells *et al.*, 2016; *Acropora tenuis*: Matias *et al.*, 2023; *A. lamarcki*: Prata *et al.*,
491 2022; but not so for *A. digitifera*: Zhang *et al.*, 2023). Intriguingly, Starko *et al.* (2023)
492 demonstrated that a distinct symbiont community associated with one cryptic taxon of
493 massive *Porites sp.* changed following a heatwave, such that the post heatwave composition
494 better matched the symbiont communities living in the other two cryptic taxa. Thus,
495 symbionts may change to track local environments (Baker, 2003), although this flexibility may
496 differ among host taxa (Quigley *et al.*, 2022).

497

498 Whereas differentiation by depth and habitat appear to be fairly common in corals (and their
499 associated microbes), sampling strategies for many coral genomic studies are surprisingly
500 underpowered in their ability to detect genetic differentiation by depth or habitat. Among the
501 population genomic studies examined here, 25% failed to report sampling depth (or any other
502 relevant habitat, including our own work, e.g. Matias *et al.*, 2023). Presumably, most of the
503 genotyped corals across the studies reviewed here were sampled on SCUBA from < 30 m and
504 were sampled from a similar depth range across all sites. Among studies that did report depth,
505 many sampling regimes had depth confounded with geography where each location was
506 sampled at a single depth. A minority of studies (21%), however, implemented a structured
507 sampling design where the same depth was sampled at more than one location (Fig. 4). More
508 complete reporting on depth and other microenvironmental attributes alongside sampling
509 study designs that replicate environmental contrasts are needed to advance our
510 understanding of how cryptic taxa, reproductive barriers, heterogeneous environments, and
511 natural selection interact to shape coral biodiversity.



512

513

514

515

516

517

518

519

520

521

522

Figure 4 - Summary of depth sampling schemes for studies that reported depth and sampled at more than one depth. Numbers indicate the number of distinct locations that were sampled per depth. Locations were considered distinct if the nearest locations were depth contrasts (e.g., adjacent sites sampled at 5 and 15 m were considered as two locations); otherwise, same depth locations within 10 km were collapsed as a single location. Dotted lines connect locations from the same study and thick grey lines indicate the sampling range (as reported by authors). Citations are as follows: a) (Thomas *et al.*, 2019); b) (Bongaerts *et al.*, 2017); c) (Prata *et al.*, 2022); d) (Gonzalez-Zapata *et al.*, 2018); e) (Bongaerts *et al.*, 2021a); f) (Shilling *et al.*, 2023); g) (Rippe *et al.*, 2021); h) (Aichelman & Barshis, 2020); i) (Drury *et al.*, 2020); j) (Sturm *et al.*, 2020); k) (Sturm *et al.*, 2022); l) (Aurelle *et al.*, 2022); m) (van Oppen *et al.*, 2018); n) (Meziere *et al.*, 2024).

523

524 **2.4 Cryptic taxa can differ in their susceptibility to thermal stress**

525 Limited but growing evidence shows that phenotypes of closely related cryptic coral taxa can
526 differ with respect to environmental tolerances, which may affect where they can live (i.e.,
527 their niche, including depth) and how they might respond to climate change and other
528 anthropogenic stressors. For example, under experimental conditions, one genetic group of
529 *A. hyacinthus* showed greater resistance to bleaching conditions under experimental trials
530 compared to the three other groups, matching within reef microhabitat distributions of these
531 taxa in American Samoa (Rose *et al.*, 2021). Likewise, heat resilience and survival may differ
532 between cryptic taxa in the wild. By monitoring tagged colonies during an extreme heatwave,
533 Starko *et al.* (2023) found that mortality was significantly higher for one of three massive
534 *Porites* taxa on Kiritimati. In Palau, cores from *P. lobata* showed differences in past growth
535 rates and stress banding between four cryptic taxa – indicative of taxa-specific differences
536 over long time periods (1970-2014), with the two most heat tolerant taxa predominating at
537 the warmest location (Rivera *et al.*, 2022). Similarly, survival through a heatwave differed
538 among *Pocillopora* taxa distributed across Moorean reefs (Burgess *et al.*, 2021). These studies
539 illustrate that cryptic taxa are not necessarily interchangeable, especially with respect to heat
540 stress responses and survival over both short and long-time frames.

541

542 In Textbox 2, we investigate whether thermal biology studies are attuned to the coral cryptic
543 taxon “problem”. If cryptic coral taxa consistently differ in their phenotypes, including
544 response to thermal stress, then experimental outcomes need to be evaluated with respect
545 to taxonomic identity. Yet, we find that only 8% of such studies included genotyping that
546 could identify cryptic taxa, suggesting that many studies could be inadvertently evaluating
547 multiple taxa and thereby supporting incorrect or biased conclusions.

548

549 **2.5 Gene flow and introgression link taxa across divergence histories**

550 Interbreeding between genetically distinct groups of corals could elevate genetic diversity
551 and contribute to adaptation. Hybridisation in corals has long been suspected (van Oppen &
552 Gates, 2006; Willis *et al.*, 2006), but studies using few genetic markers often lack the
553 resolution to appropriately investigate hybridisation in the context of recently diverged taxa,
554 where genetic similarities can result from either shared ancestral diversity or gene flow. If
555 hybrids interbreed with parental species (backcrossing), interspecific recombination leads to
556 introgression of genetic material from one lineage into the other. Thus, by analysing
557 thousands of genomic SNPs, genomic studies can often resolve whether genetic similarities
558 are due to gene flow and sometimes identify genomic regions that have experienced high or
559 low levels of gene flow as well as identify where selection may have shaped introgression
560 patterns (Taylor & Larson, 2019).

561

562 Coral population genomic studies often find individuals with possible hybrid ancestries, but
563 only two studies included explicit tests for recent hybridisation (such as tests implemented in
564 NewHybrids (Anderson & Thompson, 2002)) that probabilistically assign individuals as

565 putative first-generation and early backcrosses. Hybrid individuals were found among
566 *Agaricia* taxa (Prata *et al.*, 2022) but not among *S. pistillata* taxa (Meziere *et al.*, 2024). Other
567 studies have identified likely hybrid individuals based on the proportion of assignment to
568 different groups from model-based clustering outputs (e.g. Cooke *et al.*, 2020; Kitchen *et al.*,
569 2020; Bongaerts *et al.*, 2021a; Fifer *et al.*, 2021; Rippe *et al.*, 2021; Rivera *et al.*, 2022; Matias
570 *et al.*, 2023). Using a majority background group assignment score of < 0.75 for distinct
571 sympatric genetic groups as an indication of possible recent hybridisation, we identified
572 potential hybrid individuals for 21 of the 34 species surveyed; however, these mixed ancestry
573 individuals are based on original author preferred K groups and therefore do not necessarily
574 reflect admixture between partially reproductively isolated groups (see Textbox 3 for further
575 discussion). Nonetheless, the prevalence of individuals with mixed ancestry suggests that
576 hybridisation could be ongoing for many taxa.

577
578 Historical gene flow between divergent species can be an important aspect of their
579 evolutionary history (Shinzato *et al.*, 2015; Voolstra *et al.*, 2023) even without early
580 generation hybrid individuals being frequent. Unique signatures of historical (“ancient”) gene
581 flow between taxa have been demonstrated by contrasting population genetic demographic
582 models using alternative speciation scenarios (e.g., no gene flow versus periodic or ongoing
583 gene flow) (Gutenkunst *et al.*, 2009; Beaumont, 2010; Sousa & Hey, 2013; Fraïsse *et al.*, 2021)
584 thereby resolving the contributions of shared ancestral polymorphisms and introgression to
585 shared genetic variation among taxa. Where demographic modeling has been applied to
586 corals, divergence time estimates among cryptic taxa ranged between a few thousand years
587 (e.g., ~10k between *A. digitifera*: Zhang *et al.*, 2023) and up to 9 million years (i.e., *Galaxea*
588 *fascicularis*: Wepfer *et al.*, 2020). However, most studies reported recent divergence times
589 (~0.2-1 million years: Cooke *et al.*, 2020; Bongaerts *et al.*, 2021a; Fifer *et al.*, 2021; Matias *et*
590 *al.*, 2023; Meziere *et al.*, 2024) that predate Holocene reef configurations. This suggests that
591 divergence began before species shifted into their present-day ranges. Additionally, several
592 studies found strong support for models of speciation with ongoing or episodic gene flow
593 between genetically differentiated and sympatric and/or depth associated cryptic coral taxa
594 with low-to-moderate levels of genetic exchange (e.g., Wepfer *et al.*, 2020; Fifer *et al.*, 2021;
595 Rippe *et al.*, 2021; Prata *et al.*, 2022; Tsuchiya *et al.*, 2022; Matias *et al.*, 2023; Starko *et al.*,
596 2023; Zhang *et al.*, 2023; Meziere *et al.*, 2024) (Prada & Hellberg, 2021). These findings
597 suggest that recent divergence with gene flow is common across a range of phylogenetic and
598 functional coral diversity and is geographically widespread.

599
600 An emerging observation among diverse metazoans is that gene flow between closely-related
601 species is variable across genomes (Ravinet *et al.*, 2017) due to local adaptation or
602 reproductive incompatibilities that reduce gene flow for selected regions (Bay & Rugg, 2017;
603 Martin & Jiggins, 2017). In cases where variable introgression rates were tested explicitly
604 using demographic modeling of coral divergence, models including heterogeneous gene flow
605 rates received the highest support (Fifer *et al.*, 2021; Rippe *et al.*, 2021; Starko *et al.*, 2023;

606 Meziere *et al.*, 2024), providing evidence for genomic barriers to gene flow in the sympatric
607 taxa studied and supporting the contention that some degree of reproductive isolation helps
608 maintain genetic cohesion of these interbreeding taxa. For example, in *S. pistillata*, more
609 divergent taxa had a higher proportion of their genomes experiencing reduced gene flow
610 compared to the less divergent taxa, implying that islands of differentiation become wider as
611 speciation proceeds (Meziere *et al.*, 2023). These findings are consistent with morphologically
612 similar taxa being at various stages of the divergence process (Roux *et al.*, 2016).

613

614 Low levels of gene flow can contribute to adaptation via introgression (Martin & Jiggins,
615 2017), where alleles derived from a different species can affect a specific beneficial trait (e.g.
616 resistance to hypoxia at high altitude in humans: Huerta-Sánchez *et al.*, 2015; and winter coat
617 colour in hares Giska *et al.*, 2019). In corals, a genomic region (approximately 220 kb) that
618 appear to contribute to increased bleaching-tolerance for one *A. hyacinthus* taxon relative to
619 other cryptic *A. hyacinthus* taxa was likely acquired through hybridisation with *A. millepora*
620 (Rose *et al.*, 2021). This finding is consistent with some evidence suggesting that historical
621 introgression events may have coincided with *Acropora* range expansions, with possible links
622 to ecological opportunities associated with their diversification (Mao, Economo & Satoh,
623 2018). While these studies implicate a role for hybridisation in adaptive evolution, and the
624 possibility that large chromosomal variants may be important for adaptation in the context
625 of gene flow, there have been no comprehensive investigations of adaptive introgression in
626 corals to date.

627

628 **2.6 Implications: investigations that do not test for cryptic taxa risk biased estimates of** 629 **ecologically important traits**

630 The preceding review of population genomic studies shows that cryptic coral taxa are
631 common (Section 2.1) and may be adapted to and associated with different
632 microenvironments – especially depth (2.3). These cryptic taxa, however, are often linked by
633 sharing symbiont strains (2.2) and via some gene exchange (2.5). Therefore, cryptic taxa may
634 be distinct in terms of ecology, physiology, and evolution, but how to describe and delineate
635 taxa is not obvious, as there may not be complete divisions or obvious diagnostic
636 characteristics. Even with access to genomic-scale genotyping, taxonomic resolution is
637 affected by sampling and open to interpretation. It is clear, however, that gross morphology
638 assessed by humans under field conditions is unreliable for recognising closely related taxa,
639 as over 50% of studies in our literature search show genetic evidence for cryptic taxa (Fig. 3).
640 Simply put, any coral investigation that does not include genotyping, risks treating a
641 heterogeneous mix of partially reproductively isolated taxa as a single species. To get a sense
642 of how extensive this issue could be for other fields in coral biology, in Textbox 2 we show
643 that only 8% of coral experiments on thermal tolerance include genotyping that could identify
644 cryptic taxa. Although it is outside the scope of this paper to examine the thousands of papers
645 in coral biology that examine phenotypes including niches and geographic distributions, we
646 would expect that most experimental and ecological studies are not combined with

647 genotyping in a way that can detect cryptic taxa, if they are present. Given that unaccounted
648 cryptic taxa can bias estimates of species' traits (alongside estimates of biodiversity and
649 evolutionary dynamics - Section 1), inferences from many studies need to be viewed with
650 scepticism.

651

652 **3 Corals as untapped systems for studying speciation and adaptation in a changing** 653 **world**

654 Although cryptic species and gene flow (hybridisation) between closely related coral taxa are
655 more common than previously thought, corals present largely overlooked systems for
656 studying speciation and adaptation. Such studies could bring new insights to processes
657 generating coral biodiversity and would also clarify biological attributes of corals that may
658 sway conservation management decisions and strategies.

659

660 The emerging consensus that closely-related coral taxa are frequently sympatric at coarse
661 spatial scales yet segregate by depth or other micro environmental characteristics aligns well
662 with models of ecological speciation (Rundle & Nosil, 2005). Furthermore, the presence of
663 distinct cryptic taxa in close geographic proximity suggests that selection may be very strong
664 (a high selection to migration ratio: Richardson *et al.*, 2014)). To what extent cryptic taxa differ
665 phenotypically or in terms of competitive ability is largely unknown, although differences in
666 bleaching susceptibility among some cryptic taxa suggest differing vulnerabilities to climate
667 change (Section 2.4). As sessile organisms, corals are well-suited to manipulative experiments
668 such as common garden or reciprocal translocation designs. Additionally, their clonal nature
669 means that genetically identical fragments from the same colony can be exposed to differing
670 treatments, offering rich opportunities to combine experiments with genomic analyses to
671 holistically investigate the interactions between taxon identity, phenotype, and environment
672 (Pinsky *et al.*, 2023; Richards *et al.*, 2023).

673

674 Divergence dates between cryptic taxa often pre-date Holocene reef configurations, implying
675 that old standing genetic diversity is spread across contemporary reefs that are characterized
676 by spatially complex yet replicated microhabitats and environmental gradients. Thus, corals
677 would be ideal for investigations that explore the genetic mechanisms of parallel divergence
678 (e.g. analogous to fishes that have colonized post-glacial lakes: Rougeux, Bernatchez &
679 Gagnaire, 2017; De-Kayne *et al.*, 2022). Such investigations would also provide insights on the
680 geographic distribution of standing genetic variation, which may be under increasing selective
681 pressure due to pervasive environmental changes. For example, knowing whether
682 geographically distant populations do or do not share alleles for advantageous traits can guide
683 decisions regarding the utility of assisted migration. This is because evolutionary rescue is
684 only worth considering if donor and recipient populations differ in functional standing genetic
685 diversity.

686

687 It is likely that endogenous barriers to reproduction suppress gene flow between taxa to some
688 extent. Analyses to date support evolutionary genomic models that allow different genomic
689 regions to be more or less permeable to gene flow, consistent with chromosomal inversions
690 or other structural variants contributing to reproductive isolation (see section 2.5). Future
691 studies that use chromosomal resolution genotyping will be critical to forming a deeper
692 understanding of how species boundaries are maintained (e.g., Leitwein *et al.*, 2020) and can
693 guide decisions on assisted migration or choosing broodstock for selective breeding (Fig. 2).

694

695 Individual colonies with genotypes consistent with recent hybrid ancestry have been noted
696 (Section 2.5). To date, these individuals have largely been treated as curiosities and not
697 subject to focused study. Studies of hybridization and hybrid zones also provide important
698 insights on speciation and adaptation (Hewitt, 1988) and would yield key background
699 information to evaluate suggestions that hybrid corals are viable in nature and could be used
700 in restoration. Potential restoration interventions based on hybridisation rest on the
701 supposition that hybrid corals differ in phenotypes from parental species through some
702 combination of hybrid vigor or transgressive segregation. Promisingly, first generation (F1)
703 crosses among *Acropora* species have showed that some crosses survived as well as or better
704 than parentals in some conditions (Chan *et al.*, 2018). Aside from the transect studies of Prada
705 & Hellberg (2014), we know of no systematic attempt to map the spatial and environmental
706 distributions of hybrids relative to parental taxa, and yet this knowledge can yield important
707 insights into natural selection and dispersal of cryptic taxa (Barton & Hewitt, 1985). Mapping
708 hybrids would also allow restoration biologists to easily locate naturally-occurring hybrid
709 corals with a wide diversity of genotypes that could be used as broodstock in restoration
710 methods that rely on sexually-produced offspring (Table 1).

711

712 Throughout this review, we have focused primarily on the cnidarian component of coral
713 genomes to document evidence for cryptic species and hybridization. However, in considering
714 how future studies could build on these observations to better understand speciation and
715 adaptation processes, it will also be important to integrate genetic analyses of microbes –
716 especially symbiotic dinoflagellates – with those of the cnidarian host. An exciting line of
717 investigation would be to try to understand the co-evolutionary dynamics of hosts and
718 symbionts in reference to environmental adaptation and speciation, where environmental
719 heterogeneity likely exerts direct selection on both corals and their symbionts (i.e. the coral
720 holobiont) and indirect selection via host-symbiont genetic interactions.

721

722

4. Conclusions

723 In this review, we demonstrate that cryptic coral taxa are extremely common and are often
724 connected by gene flow. Many previous studies have emphasized the importance of such

725 cryptic coral taxa (Gomez-Corrales & Prada, 2020; Bongaerts *et al.*, 2021a; Burgess *et al.*,
726 2021; Feldman *et al.*, 2021; Rippe *et al.*, 2021; Zayasu *et al.*, 2021; Prata *et al.*, 2022; Rivera *et*
727 *al.*, 2022; Matias *et al.*, 2023; Pinsky *et al.*, 2023; Starko *et al.*, 2023; Voolstra *et al.*, 2023;
728 Meziere *et al.*, 2024) and here we use a systematic examination of published population
729 genomic studies to document that cryptic taxa are indeed widespread across coral families
730 (Fig. 3).

731

732 The prevalence of cryptic coral taxa means that many accepted understandings and
733 conclusions regarding coral biology could be incorrect. In this review, we argue that hidden
734 taxonomic diversity can affect conservation thinking and, in the extreme, mislead possible
735 management decisions if ignored. In Section 1, we highlight how ignoring cryptic taxa can:
736 bias estimates of spatial biodiversity patterns; make species appear to have larger ranges,
737 trait spaces, and niches; and can skew inferences regarding intraspecific population structure
738 and gene flow in unpredictable ways. Thus, as a field, we are unable to confidently generalize
739 species distributions, ranges, and phenotypes including resilience to heat stress without
740 adequate study of potential cryptic taxa (Textbox 3). Identifying locations with high genetic
741 diversity that may harbour greater adaptive potential or inferring sites with high gene flow
742 and dispersal will crucially depend on analyses that are able to detect and account for distinct
743 taxa.

744

745 Although observations of cryptic coral taxa are frequent, our collective knowledge regarding
746 the evolutionary dynamics that enable closely-related taxa with incomplete species
747 boundaries to persist in sympatry remains limited. There is a vast potential to unite coral
748 studies with the insights and approaches into speciation and adaptation from other fields and
749 organisms. For example, partnerships between coral ecologists, physiologists, and population
750 geneticists may bridge insights into coral population microevolution to climate change, while
751 collaborations with experts from other fields may broker novel analyses and genomic-based
752 approaches to speciation and hybridisation in corals.

753


754 **The reality, however, is that the future for corals and coral reefs is perilous (Knowlton *et al.*,**
755 **2021). Management decisions cannot wait for perfect information.** Therefore, it is incumbent
756 for evolutionary biologists to identify attributes that do or will affect coral health and
757 evolutionary trajectories for the short-term future. An important next step will be to explore
758 how conservation and management actions can best proceed with a newfound expectation
759 that coral species boundaries are unlikely to be well defined – a conservation challenge that
760 ultimately afflicts many other taxa apart from corals (Roux *et al.*, 2016).

761

762 **Textbox 1: Applying taxonomic delineation with reproductive isolation criteria**

763 We propose three requirements for identifying and delineating taxa using genomic-informed
 764 ordination and model-based clustering approaches. **In the empirical example that follows**, we
 765 detail how coral cryptic taxa were identified using three criteria (section 2.1) and highlight
 766 difficulties with interpretation.

767
 768 Colonies of the brooding coral, *Agaricia agaricites*, were sampled at four sites ~10-15 km
 769 apart along southwest Curaçao, and collections were further subset to three depths (5, 10
 770 and 20 m) at each site. Genotyping using reduced representation sequencing of 335 colonies
 771 and 1,629 SNP-loci revealed distinct genetic groups co-occurring within four sampled sites.
 772 This study provides a clear example of cryptic taxa identified according to criteria 1-3.
 773 Furthermore, taxa occupied unique depth ranges (AA1 occurs predominantly at 20 m,
 774 whereas AA2 occurred at all depth sampled) suggesting divergence of taxa by habitat (Fig.
 775 T1A & C).
 776



777
 778
 779 **Figure T1 - *Agaricia agaricites* resolved into two distinct taxa and fulfil criteria 1, 2 and 3 for cryptic**
 780 **taxon delineation. A) PC1 resolves two sympatric groups at every sampling site. PC2 represents an**
 781 **example of geographic partitioning in AA2 (Site 1 vs Site 2 - 4) and therefore does not necessarily**
 782 **imply reproductive isolation. B) Shows that the percentage of variation explaining PC1 (14.65%) is ~5x**
 783 **more than PC2 (3.01%). C) Analyses using ADMIXTURE for K=2 assigned individuals to each group with**
 784 **high confidence ($q > 0.9$) and showed that AA1 and AA2 were sympatric at all sites. D) There is a**
 785 **significant drop in cross-validation error between K=1 and K=2, and greater log-likelihood, supporting**

786 the selection of $K = 2$. All three criteria are met in delineating AA1 and AA2 cryptic taxa within *A.*
787 *agaricites*.

788

789 This example also illustrates some complications for interpreting differentiation among
790 putative taxa. The second PC axis (also mirrored in ADMIXTURE outcome for $K=3$ and $K=4$)
791 shows partitioning that largely aligns with geographic separation and so would not be
792 considered as delineating distinct taxa under our criterion of sympatry. If the variation
793 captured by the second PC axis (and $K=3$ and $K=4$) reflects geographic differentiation,
794 geographically mismatched genotypes likely reflect recent immigration (shown by dashes
795 among $K=3$ and $K=4$ in Fig. T1C). Distinguishing migrants from distinct taxa may be especially
796 difficult when sampling numbers are low. However, if immigration is high (> 1 migrant per
797 generation) and there are no barriers to reproduction then the structure between
798 populations is expected to dissipate over a few generations (Waples & Gaggiotti, 2006).

799

800 This example highlights the utility of PCA and cluster-based modeling methods for identifying
801 cryptic coral taxa. However, additional subsetting and filtering steps are necessary to
802 thoroughly scrutinize data for consistent patterns and reveal accurate groupings. To better
803 understand the possible biases of both PCA and assignment methods, we refer readers to
804 (Pritchard *et al.*, 2000; McVean, 2009; Puechmaille, 2016).

805

806 **Textbox 2: Are coral experiments designed to detect cryptic taxa?**

807 Overlooking cryptic taxa in experiments can bias interpretations of experimental results. To
808 ascertain how substantial this issue might be for coral studies, we focus on experiments
809 related to thermal tolerance as a subset of coral studies more generally. Marine heatwaves
810 have caused extensive coral mortality and bleaching events globally (Leggat *et al.*, 2019), and
811 thus numerous coral studies have aimed to ascertain intra- and inter-specific differences in
812 phenotypic and physiological heat stress responses using experiments (e.g., common
813 gardens, reciprocal transplants, etc.) and natural heating events. Mounting evidence suggests
814 that cryptic species and hybrids display contrasting responses to heat stress (Section 2.4), and
815 so experimental results may be more accurate when taking into account potential cryptic taxa
816 – identified using genomic-scale genotyping .


817

818 The Coral Research and Development Accelerator Platform (CORDAP) database (Ortiz,
819 Humanes & Scharfenstein, 2023) represents a curated search for papers that study thermal
820 biology of corals. We screened the database to identify records which used genome-wide
821 data of the coral host (i.e., coral SNP data) and those that conducted either ordination or
822 model-based clustering (as in 2.1). We evaluated the database to determine:

- 823 1. The number of studies within the database undertaking marine heatwave
824 experiments that genotyped corals for multiple unlinked markers (i.e., created data
825 that could be used for ordination or model-based clustering).
- 826 2. The proportion of these studies that performed either an ordination or model-based
827 clustering based on individual genotypes.
- 828 3. Whether there is evidence for cryptic taxa based on applying the criteria outlined in
829 Section 2.1.

830

831 We found that very few experimental studies genotyped coral colonies - from 562 studies,
832 222 studies undertook marine heatwave experiments, of which 59 studies included any sort
833 of host genotyping and only 17 used high-resolution genome-wide markers such as SNPs. Still
834 fewer studies undertook either ordination or model-based clustering on their genomic data
835 (n= 9; Fig. T2). Of those nine studies, three showed indications of cryptic taxa in line with
836 proportions of cryptic taxa in population genomic surveys (Fig. 3). It is likely that many studies
837 will have inadvertently sampled multiple taxa and therefore we would anticipate that
838 reported variances among individuals within studies will be greater than true variances within
839 cryptic taxa (see Section 1). This could manifest as a bias in overestimated thermal tolerance
840 breadth and thus may also mask differences in measured tolerance in comparative tests
841 between morphospecies. The CORDAP database focuses on one group of studies, but we
842 would anticipate that similar issues arise across all coral experimental work that does not
843 leverage genomic-level genotyping of individual colonies.



844

845 **Figure T2** – Proportion of individual studies from the Coral Research and Development Accelerator
846 Platform (CORDAP) thermal tolerance experiment database that: (1) record genotype data capable of
847 identifying cryptic taxa via genome-wide data of hosts, and (2) test for cryptic taxa via ordination-
848 based analyses or model-based clustering analyses.

849

850

851 **Textbox 3: Best practice recommendations**

852 Future genomic surveys of corals should be designed with the expectation that cryptic species
853 could be encountered. This means undertaking structured and replicated sampling,
854 reassessing field collection protocols, and testing for cryptic taxa as part of bioinformatic and

855 population genetic analyses. Ensuring that all data and metadata are thoroughly documented
856 will create the best chances that future investigations can re-examine published data as novel
857 methods emerge and thus bypassing the need for additional fieldwork and genotyping in
858 some cases.

859

860 ***Spatial sampling at the colony level***

861 The best evidence for discriminating cryptic taxa from population structure is when distinct
862 taxa are observed in sympatry (or close geographic proximity) at multiple locations (Section
863 2.1). We acknowledge that there is an element of chance in co-sampling distinct taxa. Given
864 that depth appears to be the most common axis of differentiation, however, researchers who
865 are planning to sample across depth should ensure that the same depths are sampled at
866 multiple distinct locations to enable the detection of repeated co-occurrences of distinct taxa.

867

868 Alongside structured sampling, investigators would greatly enhance their data's value (and
869 scope for future inference) by transitioning from a population sampling mindset to focusing
870 on individual sampling and seeking to capture as much environmental context as possible at
871 the colony level. For instance, "cryptic" species may in fact be morphologically distinguishable
872 based on subtle characteristics (for example, *S. pistillata*, Meziere *et al.*, 2024) and therefore
873 taking comprehensive photographs that can be examined later (see, for example, Protocol for
874 Coral Collection & Curation by Project Phoenix:
875 <https://coralprojectphoenix.org/resources/#protocols>) may allow diagnostic characters to be
876 identified post hoc. For a subset of samples, it would be useful to retain larger colony
877 fragments that would be suitable as museum voucher specimens, if permits allow. Recording
878 each colony's geolocation and depth can greatly support analyses based on depth (i.e. as a
879 continuous rather than categorical predictor) and space, while simultaneously could provide
880 insights on the microhabitat attributes of cryptic taxa and hybrid individuals (as in (Prada &
881 Hellberg, 2014)). A particularly exciting technology that can greatly advance this colony-
882 focussed perspective is photogrammetry (Bongaerts *et al.*, 2021b). We recognise that moving
883 the focus from coral populations to colonies will require more time, effort, and expense, but
884 the insights will be far richer.

885

886 Datasets that link genomic genotyping with ecological context at the colony level will be
887 immensely valuable for gaining insights into ecological and evolutionary processes relevant
888 to conservation. To maximise this value, investigators should strive to make all facets of their
889 data open, which includes linking genotypes with all recorded metadata including metadata
890 that might not be relevant to the original study (but that might be of value to others).
891 Analytical pipelines also need to be fully reproducible by enabling consistency in
892 bioinformatics and analytical decisions across studies such that outcomes can be confidently
893 compared. No doubt, all this extra documentation is a substantial amount of work, and
894 therefore should be incorporated into initial project planning. Coral biologists can take
895 inspiration from plant population geneticists who have greatly advanced insights and impacts

ie "pseudocryptic"
species? perhaps
use that term ?

I suggest being a
bit more specific
on this topic, eg
talk about the
FAIR guidelines
[https://doi.org/
10.1038/
sdata.2016.18](https://doi.org/10.1038/sdata.2016.18)

896 by sharing highly curated data sets that have been re-used and supported a myriad of
897 additional studies after their initial publication, for example, the IntraBioDiv (Meirmans *et al.*,
898 2011) dataset of 27 co-distributed alpine plant genotypes has supported numerous
899 reanalyses and test cases. Additionally, the genomic (and phenotypic) datasets for lodgepole
900 pine and spruce from the AdapTree group ([https://adaptree.forestry.ubc.ca/about/scientific-
901 summary/](https://adaptree.forestry.ubc.ca/about/scientific-summary/)) have greatly advanced understandings of spatial adaptive diversity in trees.

902

903 ***Adjusting bioinformatic pipelines and analyses***

904 Bioinformatics and population genetic analyses also need to be sensitive to the possibility of
905 cryptic taxa (see also Section 2.1 and Textbox 1). Missing data thresholds and other data
906 quality filters are employed as standard practice on individuals and loci. However, sensitivity
907 of different missing data thresholds to test taxon assignment and hierarchy hypotheses are
908 not often mentioned. The more divergent groups are, the fewer sites they will share and thus
909 blanket missing data thresholds on heterogeneous samples may bias patterns. Applying
910 different missing data filters and subsetting datasets by selecting an even representation of
911 predetermined groups (from initial structure analyses) or isolating certain groups can help in
912 determining if the assignment and hierarchy of groups is stable and robust to the filters
913 selected (Pritchard *et al.*, 2000; McVean, 2009; Puechmaile, 2016). Intermediate or admixed
914 individuals may appear as hybrid individuals, but the causes of these patterns are many, *e.g.*,
915 unexplained variance due to geographic structure, under-sampled taxa, admixture with
916 unsampled taxa, or higher levels of missing data for some individuals. Thus, we suggest formal
917 hybrid tests are employed for clarification (*e.g.*, NewHybrids). Investigators should be
918 transparent regarding how biases or decisions were handled when reporting groupings. We
919 suggest following advice from (Meirmans, 2015) by always reporting multiple K values when
920 using assignment methods, as clustering analyses represent a heuristic approach that is open
921 to interpretation for all biologically-sensible K values, even if an optimal K-value is selected
922 by the user-defined summary statistic. Similarly, PCA results should present the percent of
923 variation explained and include multiple axes. Ultimately, we hope that the guidelines
924 presented here can be used as a framework to detect coral cryptic taxa in future population
925 genomic investigations.

926

927 ***In the absence of genotypes....***

928 While population geneticists are the target audience for our recommendations, any coral
929 biologist whose data interpretations could be affected by cryptic species would do well to
930 incorporate genotyping in their project planning. For instance, genotyping is largely
931 overlooked in experiments (Textbox 2). For experimental work, we propose that future
932 studies should: 1) where possible, include larger sample sizes ($n > 30$) to screen for cryptic
933 genetic population structure (this will ensure downstream comparisons in individual
934 phenotypic differences are not confounded by cryptic speciation); 2) follow guidelines from
935 2.1 to recognise cryptic species; 3) report initial data checking methods and results (*e.g.*,
936 screening population structure) in publications and reports (*e.g.*, in supplementary items) to

937 assist the interpretation of individual- and population-level differences; and 4) clarify
938 definitions and conventions for terms such as "cryptic species" and establish common
939 terminology.

940

941

Acknowledgements

942 We thank S. Starko for photos of *Porites* sp.

943

Data, scripts, code, and supplementary information availability

944 Data and supplementary information are available linked to this pdf at EcoEvoRxiv.

945

Conflict of interest disclosure

946 The authors declare that they have no financial conflicts of interest in relation to the content of the
947 article.

948

Funding

949 CR was supported by a guest professorship ("Marks professor") funded by Göteborgs Universitet during
950 part of the manuscript preparation.

951

References

- 952
- 953 ADAM, A. A. S., THOMAS, L., UNDERWOOD, J., GILMOUR, J. & RICHARDS, Z. T. (2022). Population
954 connectivity and genetic offset in the spawning coral *Acropora digitifera* in Western
955 Australia. *Molecular Ecology* **31**, 3533-3547. 10.1111/mec.16498
- 956 ADEY, W. H. (2000). Coral reef ecosystems and human health: biodiversity counts! *Ecosystem*
957 *health* **6**(4), 227-236. 10.1046/j.1526-0992.2000.006004227.x
- 958 AFIQ-ROSLI, L., WAINWRIGHT, B. J., GAJANUR, A. R., LEE, A. C., OOI, S. K., CHOU, L. M. & HUANG, D.
959 (2021). Barriers and corridors of gene flow in an urbanized tropical reef system. *Evol*
960 *Appl* **14**(10), 2502-2515. 10.1111/eva.13276
- 961 AICHELMAN, H. E. & BARSHIS, D. J. (2020). Adaptive divergence, neutral panmixia, and algal
962 symbiont population structure in the temperate coral *Astrangia poculata* along the
963 Mid-Atlantic United States. *PeerJ* **8**, e10201. 10.7717/peerj.10201
- 964 ALEXANDER, D. H., NOVEMBRE, J. & LANGE, K. (2009). Fast model-based estimation of ancestry in
965 unrelated individuals. *Genome research* **19**(9), 1655-1664. 10.1101/gr.094052.109
- 966 ANDERSON, E. C. & THOMPSON, E. A. (2002). A model-based method for identifying species
967 hybrids using multilocus genetic data. *Genetics* **160**, 1217-1229.
968 10.1093/genetics/160.3.1217
- 969 ANTHONY, K. R. N., BAY, L. K., COSTANZA, R., FIRN, J., GUNN, J., HARRISON, P., HEYWARD, A., LUNDGREN,
970 P., MEAD, D., MOORE, T., MUMBY, P. J., VAN OPPEN, M. J. H., ROBERTSON, J., RUNGE, M. C.,
971 SUGGETT, D. J., SCHAFFELKE, B., WACHENFELD, D. & WALSH, T. (2017). New interventions are

972 needed to save coral reefs. *Nature Ecology & Evolution* **1**(10), 1-3. 10.1038/s41559-
973 017-0313-5

974 ANTHONY, K. R. N., HELMSTEDT, K. J., BAY, L. K., FIDELMAN, P., HUSSEY, K. E., LUNDGREN, P., MEAD, D.,
975 MCLEOD, I. M., MUMBY, P. J., NEWLANDS, M., SCHAFFELKE, B., WILSON, K. A. & HARDISTY, P. E.
976 (2020). Interventions to help coral reefs under global change-A complex decision
977 challenge. *PloS ONE* **15**(8), e0236399. 10.1371/journal.pone.0236399

978 AURELLE, D., PRATLONG, M., OURY, N., HAGUENAUER, A., GELIN, P., MAGALON, H., ADJEROUD, M.,
979 ROMANS, P., VIDAL-DUPIOL, J., CLAEREBOUTD, M., NOUS, C., REYNES, L., TOULZA, E., BONHOMME,
980 F., MITTA, G. & PONTAROTTI, P. (2022). Species and population genomic differentiation in
981 *Pocillopora* corals (Cnidaria, Hexacorallia). *Genetica* **150**(5), 247-262. 10.1007/s10709-
982 022-00165-7

983 BAKER, A. C. (2003). Flexibility and specificity in coral-algal symbiosis: diversity, ecology, and
984 biogeography of *Symbiodinium*. *Annual Review of Ecology, Evolution, and Systematics*
985 **34**(1), 661-689. 10.1146/annurev.ecolsys.34.011802.132417

986 BARTON, N. H. & HEWITT, G. M. (1985). Analysis of hybrid zones. *Annual Review of Ecology and*
987 *Systematics* **16**(1), 113-148. 10.1146/annurev.es.16.110185.000553

988 BAUMS, I. B. (2008). A restoration genetics guide for coral reef conservation. *Molecular Ecology*
989 **17**(12), 2796-811. 10.1111/j.1365-294X.2008.03787.x

990 BAY, L. K., ORTIZ, J. C., HUMANES, A., RIGINOS, C., BAUMS, I. B., SCHARFENSTEIN, H., ARANDA, M., PEIXOTO,
991 R., NIEHAUS, A. C. & LE PORT, A. (2023). R&D technology roadmap for understanding
992 natural adaptation and assisted evolution of corals to climate change.

993 BAY, R. A. & RUEGG, K. (2017). Genomic islands of divergence or opportunities for introgression?
994 *Proceedings of the Royal Society B* **284**(1850). 10.1098/rspb.2016.2414

995 BEAUMONT, M. A. (2010). Approximate Bayesian Computation in evolution and ecology. *Annual*
996 *Review of Ecology, Evolution, and Systematics* **41**(1), 379-406.
997 papers3://publication/doi/10.1146/annurev-ecolsys-102209-144621

998 BIERNE, N., WELCH, J., LOIRE, E., BONHOMME, F. & DAVID, P. (2011). The coupling hypothesis: why
999 genome scans may fail to map local adaptation genes. *Molecular Ecology* **20**(10),
1000 2044-2072. 10.1111/j.1365-294X.2011.05080.x

1001 BONGAERTS, P., COOKE, I. R., YING, H., WELS, D., DEN HAAN, S., HERNANDEZ-AGREDA, A., BRUNNER, C. A.,
1002 DOVE, S., ENGLEBERT, N., EYAL, G., FORET, S., GRINBLAT, M., HAY, K. B., HARI, S., HAYWARD, D.
1003 C., LIN, Y., MIHALJEVIC, M., MOYA, A., MUIR, P., SINNIGER, F., SMALLHORN-WEST, P., TORDA, G.,
1004 RAGAN, M. A., VAN OPPEN, M. J. H. & HOEGH-GULDBERG, O. (2021a). Morphological stasis
1005 masks ecologically divergent coral species on tropical reefs. *Current Biology* **31**(11),
1006 2286-2298 e8. 10.1016/j.cub.2021.03.028

1007 BONGAERTS, P., DUBÉ, C. E., PRATA, K. E., GIJSBERS, J. C., ACHLATHIS, M. & HERNANDEZ-AGREDA, A.
1008 (2021b). Reefscape genomics: Leveraging advances in 3D imaging to assess fine-scale
1009 patterns of genomic variation on coral reefs. *Frontiers in Marine Science* **8**, 638979.
1010 10.3389/fmars.2021.638979

1011 BONGAERTS, P., FRADE, P. R., OGIER, J. J., HAY, K. B., VAN BLEIJSWIJK, J., ENGLEBERT, N., VERMEIJ, M. J. A.,
1012 BAK, R. P. M., VISSER, P. M. & HOEGH-GULDBERG, O. (2013). Sharing the slope: depth
1013 partitioning of agariciid corals and associated *Symbiodinium* across shallow and
1014 mesophotic habitats (2-60 m) on a Caribbean reef. *BMC Evolutionary Biology* **13**(1), 1-
1015 1. 10.1186/1471-2148-13-205

1016 BONGAERTS, P., RIGINOS, C., BRUNNER, R., ENGLEBERT, N., SMITH, S. R. & HOEGH-GULDBERG, O. (2017).
1017 Deep reefs are not universal refuges: Reseeding potential varies among coral species.
1018 *Science Advances* **3**(2), e1602373. 10.1126/sciadv.1602373

- 1019 BONGAERTS, P., RIGINOS, C., HAY, K. B., OPPEN, M. J. H. v. & HOEGH-GULDBERG, O. (2011). Adaptive
1020 divergence in a scleractinian coral: physiological adaptation of *Seriatopora hystrix* to
1021 shallow and deep reef habitats. *BMC Evolutionary Biology* **11**, 1-15. 10.1186/1471-
1022 2148-11-303
- 1023 BONGAERTS, P., RIGINOS, C., RIDGWAY, T., SAMPAYO, E. M., VAN OPPEN, M. J. H., ENGLEBERT, N.,
1024 VERMEULEN, F. & HOEGH-GULDBERG, O. (2010). Genetic divergence across habitats in the
1025 widespread coral *Seriatopora hystrix* and its associated *Symbiodinium*. *PloS ONE* **5**(5),
1026 e10871. 10.1371/journal.pone.0010871
- 1027 BONGAERTS, P. & SMITH, T. B. (2019). Beyond the “Deep Reef Refuge” hypothesis: a conceptual
1028 framework to characterize persistence at depth. In *Mesophotic coral ecosystems. Coral Reefs of the World* (Volume 12, eds Y. Loya, K. A. Puglise and T. C. L. Bridge), pp.
1029 881-895. 10.1007/978-3-319-92735-0_45
- 1030 BOULAY, J. N., HELLBERG, M. E., CORTES, J. & BAUMS, I. B. (2014). Unrecognized coral species
1031 diversity masks differences in functional ecology. *Proceedings of the Royal Society B*
1032 **281**(1776), 20131580. 10.1098/rspb.2013.1580
- 1033 BUITRAGO-LOPEZ, C., CARDENAS, A., HUME, B. C. C., GOSSELIN, T., STAUBACH, F., ARANDA, M., BARSHIS, D.
1034 J., SAWALL, Y. & VOOLSTRA, C. R. (2023). Disparate population and holobiont structure of
1035 pocilloporid corals across the Red Sea gradient demonstrate species-specific
1036 evolutionary trajectories. *Molecular Ecology* **32**(9), 2151-2173. 10.1111/mec.16871
- 1037 BURGESS, S. C., JOHNSTON, E. C., WYATT, A. S., LEICHTER, J. J. & EDMUNDS, P. J. (2021). Response
1038 diversity in corals: hidden differences in bleaching mortality among cryptic *Pocillopora*
1039 species. *Ecology* **102**(6), e03324. 10.1002/ecy.3324
- 1040 CHAN, W. Y., PEPLOW, L. M., MENÉNDEZ, P., HOFFMANN, A. A. & VAN OPPEN, M. J. H. (2018).
1041 Interspecific hybridization may provide novel opportunities for coral reef restoration.
1042 *Frontiers in Marine Science* **5**, 161. 10.3389/fmars.2018.00160
- 1043 CHAN, W. Y., PEPLOW, L. M. & VAN OPPEN, M. J. H. (2019). Interspecific gamete compatibility and
1044 hybrid larval fitness in reef-building corals: Implications for coral reef restoration.
1045 *Scientific Reports* **9**(1), 4757. 10.1038/s41598-019-41190-5
- 1046 COLTON, M. A., MCMANUS, L. C., SCHINDLER, D. E., MUMBY, P. J., PALUMBI, S. R., WEBSTER, M. M.,
1047 ESSINGTON, T. E., FOX, H. E., FORREST, D. L., SCHILL, S. R., POLLOCK, F. J., DEFILIPPO, L. B., TEKWA,
1048 E. W., WALSWORTH, T. E. & PINSKY, M. L. (2022). Coral conservation in a warming world
1049 must harness evolutionary adaptation. *Nature Ecology & Evolution*. 10.1038/s41559-
1050 022-01854-4
- 1051 COOKE, I., YING, H., FORÊT, S., BONGAERTS, P., STRUGNELL, J., SIMAKOV, O., ZHANG, J., FIELD, M. A.,
1052 RODRIGUEZ-LANETTY, M., BELL, S. C., BOURNE, D. G., VAN OPPEN, M. J. H., RAGAN, M. A. &
1053 MILLER, D. J. (2020). Signatures of selection in the coral holobiont reveal complex
1054 adaptations to inshore environments driven by Holocene climate change. *Science*
1055 *Advances* **6**, eabc6318. 10.1101/2020.02.25.951905
- 1056 COWMAN, P. F., QUATTRINI, A. M., BRIDGE, T. C. L., WATKINS-COLWELL, G. J., FADLI, N., GRINBLAT, M.,
1057 ROBERTS, T. E., MCFADDEN, C. S., MILLER, D. J. & BAIRD, A. H. (2020). An enhanced target-
1058 enrichment bait set for Hexacorallia provides phylogenomic resolution of the staghorn
1059 corals (Acroporidae) and close relatives. *Molecular Phylogenetics and Evolution* **153**,
1060 106944. 10.1016/j.ympev.2020.106944
- 1061 DAVIES, S. W., GAMACHE, M. H., HOWE-KERR, L. I., KRIEFALL, N. G., BAKER, A. C., BANASZAK, A. T., BAY, L.
1062 K., BELLANTUONO, A. J., BHATTACHARYA, D., CHAN, C. X., CLAAR, D. C., COFFROTH, M. A.,
1063 CUNNING, R., DAVY, S. K., DEL CAMPO, J., DIAZ-ALMEYDA, E. M., FROMMLET, J. C., FUESS, L. E.,
1064 GONZALEZ-PECH, R. A., GOULET, T. L., HOADLEY, K. D., HOWELLS, E. J., HUME, B. C. C., KEMP, D.
1065

1066 W., KENKEL, C. D., KITCHEN, S. A., LAJEUNESSE, T. C., LIN, S., MCLROY, S. E., MCMINDS, R.,
1067 NITSCHKE, M. R., OAKLEY, C. A., PEIXOTO, R. S., PRADA, C., PUTNAM, H. M., QUIGLEY, K., REICH,
1068 H. G., REIMER, J. D., RODRIGUEZ-LANETTY, M., ROSALES, S. M., SAAD, O. S., SAMPAYO, E. M.,
1069 SANTOS, S. R., SHOGUCHI, E., SMITH, E. G., STAT, M., STEPHENS, T. G., STRADER, M. E., SUGGETT,
1070 D. J., SWAIN, T. D., TRAN, C., TRAYLOR-KNOWLES, N., VOOLSTRA, C. R., WARNER, M. E., WEIS, V.
1071 M., WRIGHT, R. M., XIANG, T., YAMASHITA, H., ZIEGLER, M., CORREA, A. M. S. & PARKINSON, J. E.
1072 (2023). Building consensus around the assessment and interpretation of
1073 Symbiodiniaceae diversity. *PeerJ* **11**, e15023. 10.7717/peerj.15023

1074 DE QUEIROZ, K. (2005). Ernst Mayr and the modern concept of species. *Proceedings of the*
1075 *National Academy of Sciences of the United States of America* **102 Suppl 1**(Suppl 1),
1076 6600-7. 10.1073/pnas.0502030102

1077 DE-KAYNE, R., SELZ, O. M., MARQUES, D. A., FREI, D., SEEHAUSEN, O. & FEULNER, P. G. D. (2022).
1078 Genomic architecture of adaptive radiation and hybridization in Alpine whitefish.
1079 *Nature Communications* **13**(1), 4479. 10.1038/s41467-022-32181-8

1080 DIETZEL, A., BODE, M., CONNOLLY, S. R. & HUGHES, T. P. (2021). The population sizes and global
1081 extinction risk of reef-building coral species at biogeographic scales. *Nature Ecology &*
1082 *Evolution* **5**(5), 663-669. 10.1038/s41559-021-01393-4

1083 DRURY, C., PEREZ PORTELA, R., SERRANO, X. M., OLEKSIK, M. & BAKER, A. C. (2020). Fine-scale
1084 structure among mesophotic populations of the great star coral *Montastraea*
1085 *cavernosa* revealed by SNP genotyping. *Ecology and Evolution* **10**(12), 6009-6019.
1086 10.1002/ece3.6340

1087 FELDMAN, B., AFIQ-ROSLI, L., SIMON-BLECHER, N., BOLLATI, E., WAINWRIGHT, B. J., BONGAERTS, P., HUANG,
1088 D. & LEVY, O. (2021). Distinct lineages and population genomic structure of the coral
1089 *Pachyseris speciosa* in the small equatorial reef system of Singapore. *Coral Reefs* **41**(3),
1090 575-585. 10.1007/s00338-021-02160-4

1091 FIFER, J. E., YASUDA, N., YAMAKITA, T., BOVE, C. B. & DAVIES, S. W. (2021). Genetic divergence and
1092 range expansion in a western North Pacific coral. *Science of the Total Environment*
1093 **813**, 152423. 10.1016/j.scitotenv.2021.152423

1094 FISHER, R., O'LEARY, R. A., LOW-CHOY, S., MENGERSEN, K., KNOWLTON, N., BRAINARD, R. E. & CALEY, M.
1095 J. (2015). Species richness on coral reefs and the pursuit of convergent global
1096 estimates. *Current Biology* **25**(4), 500-5. 10.1016/j.cub.2014.12.022

1097 FORSMAN, Z. H., KNAPP, I. S. S., TISTHAMMER, K., EATON, D. A. R., BELCAID, M. & TOONEN, R. J. (2017).
1098 Coral hybridization or phenotypic variation? Genomic data reveal gene flow between
1099 *Porites lobata* and *P. compressa*. *Molecular Phylogenetics and Evolution* **111**, 132-148.
1100 10.1016/j.ympev.2017.03.023

1101 FRAÏSSE, C., POPOVIC, I., MAZOYER, C., ROMIGUIER, J., LOIRE, E., SIMON, A., GALTIER, N., DURET, L., BIERNE,
1102 N., VEKEMANS, X. & ROUX, C. (2021). DILS : Demographic inferences with linked selection.
1103 *Molecular Ecology Resources* **21**, 2629-2644. 10.1111/1755-0998.13323

1104 GÉLIN, P., FAUVELOT, C., BIGOT, L., BALLY, J. & MAGALON, H. (2017). From population connectivity to
1105 the art of striping Russian dolls: the lessons from *Pocillopora* corals. **8**(2), 1411-1426.
1106 10.1002/ece3.3747

1107 GISKA, I., FARELO, L., PIMENTA, J., SEIXAS, F. A., FERREIRA, M. S., MARQUES, J. P., MIRANDA, I., LETTY, J.,
1108 JENNY, H., HACKLANDER, K., MAGNUSSEN, E. & MELO-FERREIRA, J. (2019). Introgression drives
1109 repeated evolution of winter coat color polymorphism in hares. *Proceedings of the*
1110 *National Academy of Sciences of the United States of America* **116**(48), 1-7.
1111 10.1073/pnas.1910471116

- 1112 GOMEZ-CORRALES, M. & PRADA, C. (2020). Cryptic lineages respond differently to coral bleaching.
 1113 *Molecular Ecology* **29**, 4265-4273. 10.1111/mec.15631
- 1114 GONZÁLEZ, A. M., PRADA, C. A., ÁVILA, V. & MEDINA, M. (2020). Ecological speciation in corals. In
 1115 *Population Genomics: Marine Organisms. Population Genomics* (eds M. F. Oleksiak
 1116 and O. P. Rajora), pp. 303-324. 10.1007/13836_2018_35
- 1117 GONZÁLEZ, I. C. O., RIVERA-VICÉNS, R. E. & SCHIZAS, N. V. (2021). Description of four *Millepora* spp.
 1118 transcriptomes and their potential to delimit the Caribbean fire coral species. *Marine*
 1119 *Genomics* **59**, 100863. 10.1016/j.margen.2021.100863
- 1120 GONZALEZ-ZAPATA, F. L., BONGAERTS, P., RAMÍREZ-PORTILLA, C., ADU-OPPONG, B., WALLJASPER, G., REYES,
 1121 A. & SANCHEZ, J. A. (2018). Holobiont diversity in a reef-building coral over its entire
 1122 depth range in the mesophotic zone. *Frontiers in Marine Science* **5**.
 1123 10.3389/fmars.2018.00029
- 1124 GUTENKUNST, R. N., HERNANDEZ, R. D., WILLIAMSON, S. H. & BUSTAMANTE, C. D. (2009). Inferring the
 1125 joint demographic history of multiple populations from multidimensional SNP
 1126 frequency data. *PLoS Genetics* **5**(10), e1000695. 10.1371/journal.pgen.1000695
- 1127 HEIN, M. Y., MCLEOD, I. M., SHAVER, E. C., VARDI, T., PIOCH, S., BOSTRÖM-EINARSSON, L., AHMED, M. &
 1128 GRIMSDITCH, G. (2020). Coral Reef Restoration as a strategy to improve ecosystem
 1129 services A guide to coral restoration methods. United Nations Environment Program,
 1130 Nairobi, Kenya.
- 1131 HEWITT, G. M. (1988). Hybrid zones-natural laboratories for evolutionary studies. *Trends in*
 1132 *Ecology and Evolution* **3**, 158-167.
- 1133 HOBAN, S., ARCHER, F. I., BERTOLA, L. D., BRAGG, J. G., BREED, M. F., BRUFORD, M. W., COLEMAN, M. A.,
 1134 EKBLUM, R., FUNK, W. C., GRUEBER, C. E., HAND, B. K., JAFFE, R., JENSEN, E., JOHNSON, J. S.,
 1135 KERSHAW, F., LIGGINS, L., MACDONALD, A. J., MERGEAY, J., MILLER, J. M., MULLER-KARGER, F.,
 1136 O'BRIEN, D., PAZ-VINAS, I., POTTER, K. M., RAZGOUR, O., VERNESI, C. & HUNTER, M. E. (2022).
 1137 Global genetic diversity status and trends: towards a suite of Essential Biodiversity
 1138 Variables (EBVs) for genetic composition. *Biological Reviews* **97**(4), 1511-1538.
 1139 10.1111/brv.12852
- 1140 HOBBS, J.-P. A., RICHARDS, Z. T., POPOVIC, I., LEI, C., STAEUDLE, T. M., MONTANARI, S. R. & DIBATTISTA, J.
 1141 D. (2021). Hybridisation and the evolution of coral reef biodiversity. *Coral Reefs*, 1-15.
 1142 10.1007/s00338-021-02193-9
- 1143 HOWELLS, E. J., ABREGO, D., MEYER, E., KIRK, N. L. & BURT, J. A. (2016). Host adaptation and
 1144 unexpected symbiont partners enable reef-building corals to tolerate extreme
 1145 temperatures. *Global Change Biology* **22**(8), 2702-14. 10.1111/gcb.13250
- 1146 HOWELLS, E. J., WILLIS, B. L., BAY, L. K. & VAN OPPEN, M. J. H. (2013). Spatial and temporal genetic
 1147 structure of *Symbiodinium* populations within a common reef-building coral on the
 1148 Great Barrier Reef. *Molecular Ecology* **22**(14), 3693-3708. 10.1111/mec.12342
- 1149 HUERTA-SÁNCHEZ, E., JIN, X., ASAN, BIANBA, Z., PETER, B. M., VINCKENBOSCH, N., LIANG, Y., YI, X., HE, M.,
 1150 SOMEL, M., NI, P., WANG, B., OU, X., HUASANG, LUOSANG, J., CUO, Z. X. P., LI, K., GAO, G., YIN,
 1151 Y., WANG, W., ZHANG, X., XU, X., YANG, H., LI, Y., WANG, J., WANG, J. & NIELSEN, R. (2015).
 1152 Altitude adaptation in Tibetans caused by introgression of Denisovan-like DNA. *Nature*
 1153 **512**(7513), 194-197. 10.1038/nature13408
- 1154 HUGHES, T. P., KERRY, J. T., ÁLVAREZ-NORIEGA, M., ÁLVAREZ-ROMERO, J. G., ANDERSON, K. D., BAIRD, A.
 1155 H., BABCOCK, R. C., BEGER, M., BELLWOOD, D. R., BERKELMANS, R., BRIDGE, T. C., BUTLER, I. R.,
 1156 BYRNE, M., CANTIN, N. E., COMEAU, S., CONNOLLY, S. R., CUMMING, G. S., DALTON, S. J., DIAZ-
 1157 PULIDO, G., EAKIN, C. M., FIGUEIRA, W. F., GILMOUR, J. P., HARRISON, H. B., HERON, S. F., HOEY,
 1158 A. S., HOBBS, J.-P. A., HOOGENBOOM, M. O., KENNEDY, E. V., KUO, C.-Y., LOUGH, J. M., LOWE, R.

- 1159 J., LIU, G., MCCULLOCH, M. T., MALCOLM, H. A., MCWILLIAM, M. J., PANDOLFI, J. M., PEARS, R.
 1160 J., PRATCHETT, M. S., SCHOEPF, V., SIMPSON, T., SKIRVING, W. J., SOMMER, B., TORDA, G.,
 1161 WACHENFELD, D. R., WILLIS, B. L. & WILSON, S. K. (2017). Global warming and recurrent
 1162 mass bleaching of corals. *Nature* **543**(7645), 373-377. 10.1038/nature21707
- 1163 ISHIDA, H., JOHN, U., MURRAY, S. A., BHATTACHARYA, D. & CHAN, C. X. (2023). Developing model
 1164 systems for dinoflagellates in the post-genomic era. *Journal of Phycology* **59**(5), 799-
 1165 808. 10.1111/jpy.13386
- 1166 ISOMURA, N., IWAO, K., MORITA, M. & FUKAMI, H. (2016). Spawning and fertility of F1 hybrids of
 1167 the coral genus *Acropora* in the Indo-Pacific. *Coral Reefs* **35**(3), 851-855.
 1168 10.1007/s00338-016-1461-9
- 1169 JOHNSTON, E. C., CUNNING, R. & BURGESS, S. C. (2022). Cophylogeny and specificity between
 1170 cryptic coral species (*Pocillopora* spp.) at Mo'orea and their symbionts
 1171 (Symbiodiniaceae). *Molecular Ecology* **31**(20), 5368-5385. 10.1111/mec.16654
- 1172 KAHNG, S. E., AKKAYNAK, D., SHLESINGER, T., HOCHBERG, E. J., WIEDENMANN, J., TAMIR, R. & TCHERNOV,
 1173 D. (2019). Light, temperature, photosynthesis, heterotrophy, and the lower depth
 1174 limits of mesophotic coral ecosystems. In *Mesophotic coral ecosystems. Coral Reefs of*
 1175 *the World* (Volume 12, eds Y. Loya, K. A. Puglise and T. C. L. Bridge), pp. 801-828.
 1176 10.1007/978-3-319-92735-0_45
- 1177 KITAHARA, M. V., FUKAMI, H., BENZONI, F. & HUANG, D. (2016). The new systematics of Scleractinia:
 1178 Integrating molecular and morphological evidence. In *The Cnidaria, Past, Present and*
 1179 *Future*. (Volume 7, pp. 41-59. Springer, Cham. 10.1007/978-3-319-31305-4_4
- 1180 KITANOBO, S., IWAO, K., FUKAMI, H., ISOMURA, N. & MORITA, M. (2022). First evidence for
 1181 backcrossing of F(1) hybrids in *Acropora* corals under sperm competition. *Scientific*
 1182 *Reports* **12**(1), 5356. 10.1038/s41598-022-08989-1
- 1183 KITCHEN, S. A., VON KUSTER, G., KUNTZ, K. L. V., REICH, H. G., MILLER, W., GRIFFIN, S., FOGARTY, N. D. &
 1184 BAUMS, I. B. (2020). STAGdb: a 30K SNP genotyping array and Science Gateway for
 1185 *Acropora* corals and their dinoflagellate symbionts. *Scientific Reports* **10**(1), 12488.
 1186 10.1038/s41598-020-69101-z
- 1187 KLAUS, J. S., JANSE, I., HEIKOOP, J. M., SANFORD, R. A. & FOUKE, B. W. (2007). Coral microbial
 1188 communities, zooxanthellae and mucus along gradients of seawater depth and coastal
 1189 pollution. *Environmental Microbiology* **9**(5), 1291-1305. 10.1111/j.1462-
 1190 2920.2007.01249.x
- 1191 KNOWLTON, N. (1993). Sibling species in the sea. *Annual Review of Ecology and Systematics*,
 1192 189-216.
- 1193 KNOWLTON, N. (2001). The future of coral reefs. *Proc Natl Acad Sci U S A* **98**(10), 5419-25.
 1194 10.1073/pnas.091092998
- 1195 KNOWLTON, N., GROTTOLI, A. G., KLEYPAS, J., OBURA, D., CORCORAN, E., DE GOEIJ, J. M., FELIS, T.,
 1196 HARDING, S., MAYFIELD, A., MILLER, M., OSUKA, K., PEIXOTO, R. S., RANDALL, C. J., VOOLSTRA, C.
 1197 R., WELLS, S., WILD, C. & FERSE, S. (2021). Rebuilding coral reefs: A decadal grand
 1198 challenge. International coral reef society and future earth coasts pp. 56. International
 1199 Coral Reef Society and Future Earth Coasts
- 1200 LADNER, J. T. & PALUMBI, S. R. (2012). Extensive sympatry, cryptic diversity and introgression
 1201 throughout the geographic distribution of two coral species complexes. *Molecular*
 1202 *Ecology* **21**(9), 2224-2238. 10.1111/j.1365-294X.2012.05528.x
- 1203 LAJEUNESSE, T. C., PARKINSON, J. E., GABRIELSON, P. W., JEONG, H. J., REIMER, J. D., VOOLSTRA, C. R. &
 1204 SANTOS, S. R. (2018). Systematic revision of Symbiodiniaceae highlights the antiquity

1205 and diversity of coral endosymbionts. *Current Biology* **28**(16), 2570-2580 e6.
1206 10.1016/j.cub.2018.07.008

1207 LEGGAT, W. P., CAMP, E. F., SUGGETT, D. J., HERON, S. F., FORDYCE, A. J., GARDNER, S., DEAKIN, L., TURNER,
1208 M., BEECHING, L. J., KUZHIUMPARAMBIL, U., EAKIN, C. M. & AINSWORTH, T. D. (2019). Rapid
1209 coral decay is associated with marine heatwave mortality events on reefs. *Current*
1210 *Biology* **29**(16), 2723-2730 e4. 10.1016/j.cub.2019.06.077

1211 LEICHTER, J. J., STOKES, M. D. & GENOVESE, S. J. (2008). Deep water macroalgal communities
1212 adjacent to the Florida Keys reef tract. *Marine Ecology Progress Series* **356**, 123-138.
1213 10.3354/meps07230

1214 LEITWEIN, M., DURANTON, M., ROUGEMONT, Q., GAGNAIRE, P. A. & BERNATCHEZ, L. (2020). Using
1215 haplotype information for conservation genomics. *Trends in Ecology and Evolution*
1216 **35**(3), 245-258. 10.1016/j.tree.2019.10.012

1217 LESSER, M. P., SLATTERY, M. & LEICHTER, J. J. (2009). Ecology of mesophotic coral reefs. *Journal of*
1218 *Experimental Marine Biology and Ecology* **375**(1-2), 1-8. 10.1016/j.jembe.2009.05.009

1219 MALLET, J. (1995). A species definition for the Modern Synthesis. *Trends in Ecology and*
1220 *Evolution* **10**, 294-299.

1221 MAO, Y., ECONOMO, E. P. & SATOH, N. (2018). The roles of introgression and climate change in
1222 the rise to dominance of *Acropora* corals. *Current Biology* **28**(21), 3373-3382.e5.
1223 10.1016/j.cub.2018.08.061

1224 MAO, Y. & SATOH, N. (2019). A likely ancient genome duplication in the speciose reef-building
1225 coral genus, *Acropora*. *iScience* **13**, 20-32. 10.1016/j.isci.2019.02.001

1226 MARTIN, S. H. & JIGGINS, C. D. (2017). Interpreting the genomic landscape of introgression.
1227 *Current opinion in genetics & development* **47**, 69-74. 10.1016/j.gde.2017.08.007

1228 MATIAS, A. M. A., POPOVIC, I., THIA, J. A., COOKE, I. R., TORDA, G., LUKOSCHEK, V., BAY, L. K., KIM, S. W.
1229 & RIGINOS, C. (2023). Cryptic diversity and spatial genetic variation in the coral *Acropora*
1230 *tenuis* and its endosymbionts across the Great Barrier Reef. *Evolutionary Applications*
1231 **16**, 293-310. 10.1111/eva.13435

1232 McVEAN, G. (2009). A genealogical interpretation of principal components analysis. *PLoS*
1233 *Genetics* **5**(10), e1000686. 10.1371/journal.pgen.1000686

1234 MEIRMANS, P. G. (2015). Seven common mistakes in population genetics and how to avoid
1235 them. *Molecular Ecology* **24**(13), 3223-3231. 10.1111/mec.13243

1236 MEIRMANS, P. G., GOUDET, J., INTRABIODIV, C. & GAGGIOTTI, O. E. (2011). Ecology and life history
1237 affect different aspects of the population structure of 27 high-alpine plants. *Molecular*
1238 *Ecology* **20**(15), 3144-3155. 10.1111/j.1365-294X.2011.05164.x

1239 MEZIERE, Z., POPOVIC, I., PRATA, K., PANDOLFI, J. M. & RIGINOS, C. (2024). Exploring coral speciation:
1240 multiple *Stylophora pistillata* taxa on a divergence continuum on the Great Barrier
1241 Reef. *Evolutionary Applications* (17), e13644. 10.1111/eva.13644

1242 MOBERG, F. & FOLKE, C. (1999). Ecological goods and services of coral reef ecosystems.
1243 *Ecological economics* **29**(2), 215-233. 10.1016/S0921-8009(99)00009-9

1244 MUIR, P., WALLACE, C., BRIDGE, T. C. L. & BONGAERTS, P. (2015). Diverse staghorn coral fauna on
1245 the mesophotic reefs of north-east Australia. *PLoS ONE* **10**(2), e0117933.
1246 10.1371/journal.pone.0117933

1247 MUIR, P. R., OBURA, D. O., HOEKSEMA, B. W., SHEPPARD, C., PICHON, M. & RICHARDS, Z. T. (2022).
1248 Conclusions of low extinction risk for most species of reef-building corals are
1249 premature. *Nature Ecology & Evolution* **6**(4), 357-358. 10.1038/s41559-022-01659-5

- 1250 NATIONAL ACADEMIES OF SCIENCES, E., AND MEDICINE. (2018). A research review of interventions to
1251 increase the persistence and resilience of coral reefs. The National Academies Press,
1252 Washington, DC.
- 1253 ORTIZ, J. C., HUMANES, A. & SCHARFENSTEIN, H. (2023). Natural adaptation and assisted evolution
1254 of corals to heat stress (ed. Australian Institute of Marine Science).
- 1255 PANTE, E., PUILANDRE, N., VIRICEL, A., ARNAUD-HAOND, S., AURELLE, D., CASTELIN, M., CHENUIL, A.,
1256 DESTOMBE, C., FORCIOLI, D., VALERO, M., VIARD, F. & SAMADI, S. (2014). Species are
1257 hypotheses: avoid connectivity assessments based on pillars of sand. *Molecular*
1258 *Ecology* **24**(3), 525-544. 10.1111/mec.13048
- 1259 PATTERSON, N., PRICE, A. L. & REICH, D. E. (2006). Population structure and eigenanalysis. *PLoS*
1260 *Genetics* **2**(12), e190.
- 1261 PINSKY, M. L., CLARK, R. D. & BOS, J. T. (2023). Coral reef population genomics in an age of global
1262 change. *Annual Review of Genetics* (57), 87-115. 10.1146/annurev-genet-022123-
1263 102748
- 1264 PRADA, C. & HELLBERG, M. E. (2013). Long prereproductive selection and divergence by depth in
1265 a Caribbean candelabrum coral. *Proceedings of the National Academy of Sciences of*
1266 *the United States of America* **110**(10), 3961-3966. 10.1073/pnas.1208931110
- 1267 PRADA, C. & HELLBERG, M. E. (2014). Strong natural selection on juveniles maintains a narrow
1268 adult hybrid zone in a broadcast spawner. *American Naturalist* **184**(6), 702-13.
1269 10.1086/678403
- 1270 PRADA, C. & HELLBERG, M. E. (2021). Speciation-by-depth on coral reefs: Sympatric divergence
1271 with gene flow or cryptic transient isolation? *Journal of Evolutionary Biology* **34**, 128-
1272 137. 10.1111/jeb.13731
- 1273 PRADA, C., MCLROY, S. E., BELTRAN, D. M., VALINT, D. J., FORD, S. A., HELLBERG, M. E. & COFFROTH, M.
1274 A. (2014). Cryptic diversity hides host and habitat specialization in a gorgonian-algal
1275 symbiosis. *Molecular Ecology* **23**(13), 3330-40. 10.1111/mec.12808
- 1276 PRATA, K. E., RIGINOS, C., GUTENKUNST, R. N., LATIJNHOUWERS, K., SÁNCHEZ, J. A., ENGLEBERT, N., HAY, K.
1277 & BONGAERTS, P. (2022). Deep connections: divergence histories with gene flow in
1278 mesophotic *Agaricia* corals. *Molecular Ecology* **9**, 2511-2527. 10.1111/mec.16391
- 1279 PRITCHARD, J. K., STEPHENS, M. & DONNELLY, P. (2000). Inference of population structure using
1280 multilocus genotype data. *Genetics* **155**, 945-959.
- 1281 PUECHMAILLE, S. J. (2016). The program structure does not reliably recover the correct
1282 population structure when sampling is uneven: subsampling and new estimators
1283 alleviate the problem. *Molecular Ecology Resources* **16**(3), 608-27. 10.1111/1755-
1284 0998.12512
- 1285 QUIGLEY, K. M., RAMSBY, B., LAFFY, P., HARRIS, J., MOCELLIN, V. J. & BAY, L. K. (2022). Symbioses are
1286 restructured by repeated mass coral bleaching. *Science Advances* **8**(49), eabq8349.
1287 10.1126/sciadv.abq8349
- 1288 RAVINET, M., FARIA, R., BUTLIN, R. K., GALINDO, J., BIERNE, N., RAFAJLOVIC, M., NOOR, M. A. F., MEHLIG,
1289 B. & WESTRAM, A. M. (2017). Interpreting the genomic landscape of speciation: a road
1290 map for finding barriers to gene flow. *Journal of Evolutionary Biology* **30**(8), 1450-
1291 1477. 10.1111/jeb.13047
- 1292 REAKA-KUDLA, M. L. (1997). The global biodiversity of coral reefs: a comparison with rain
1293 forests. In *Biodiversity II: Understanding and Protecting Our Biological Resources*. (eds
1294 M. L. Reaka-Kudla, D. E. Wilson and E. O. Wilson), pp. 83–108. Joseph Henry Press,
1295 Washington, DC.

- 1296 RICHARDS, T. J., MCGUIGAN, K., AGUIRRE, J. D., HUMANES, A., BOZEC, Y.-M., MUMBY, P. J. & RIGINOS, C.
1297 (2023). Moving beyond heritability in the search for coral adaptive potential. *Global*
1298 *Change Biology* **29**(14), 3869-3882. 10.1111/gcb.16719
- 1299 RICHARDS, Z. T., BERRY, O. & VAN OPPEN, M. J. H. (2016). Cryptic genetic divergence within
1300 threatened species of *Acropora* coral from the Indian and Pacific Oceans. *Conservation*
1301 *Genetics* **17**(3), 577-591. 10.1007/s10592-015-0807-0
- 1302 RICHARDS, Z. T., VAN OPPEN, M. J. H., WALLACE, C. C., WILLIS, B. L. & MILLER, D. J. (2008). Some rare
1303 indo-pacific coral species are probable hybrids. *PloS ONE* **3**(9), e3240.
1304 10.1371/journal.pone.0003240.t001
- 1305 RICHARDSON, J. L., URBAN, M. C., BOLNICK, D. I. & SKELLY, D. K. (2014). Microgeographic adaptation
1306 and the spatial scale of evolution. *Trends Ecol Evol* **29**(3), 165-76.
1307 10.1016/j.tree.2014.01.002
- 1308 RIPPE, J. P., DIXON, G., FULLER, Z. L., LIAO, Y. & MATZ, M. (2021). Environmental specialization and
1309 cryptic genetic divergence in two massive coral species from the Florida Keys Reef
1310 Tract. *Molecular Ecology* **30**, 3468–3484. 10.1111/mec.15931
- 1311 RIVERA, H. E., COHEN, A. L., THOMPSON, J. R., BAUMS, I. B., FOX, M. D. & MEYER-KAISER, K. S. (2022).
1312 Palau's warmest reefs harbor thermally tolerant corals that thrive across different
1313 habitats. *Communications Biology* **5**(1), 1394. 10.1038/s42003-022-04315-7
- 1314 ROSE, N. H., BAY, R. A., MORIKAWA, M. K. & PALUMBI, S. R. (2017). Polygenic evolution drives
1315 species divergence and climate adaptation in corals. *Evolution* **72**(1), 82-94.
1316 10.1111/evo.13385
- 1317 ROSE, N. H., BAY, R. A., MORIKAWA, M. K., THOMAS, L., SHEETS, E. A. & PALUMBI, S. R. (2021). Genomic
1318 analysis of distinct bleaching tolerances among cryptic coral species. *Proceedings of*
1319 *the Royal Society B* **288**(1960), 20210678. 10.1098/rspb.2021.0678
- 1320 ROUGEUX, C., BERNATCHEZ, L. & GAGNAIRE, P. A. (2017). Modeling the multiple facets of speciation-
1321 with-gene-flow toward inferring the divergence history of lake whitefish species pairs
1322 (*Coregonus clupeaformis*). *Genome Biology and Evolution* **9**(8), 2057-2074.
1323 10.1093/gbe/evx150
- 1324 ROUX, C., FRAÏSSE, C., ROMIGUIER, J., ANCIAUX, Y., GALTIER, N. & BIERNE, N. (2016). Shedding light on
1325 the grey zone of speciation along a continuum of genomic divergence. *PLoS Biology*
1326 **14**(12), e2000234-22. 10.1371/journal.pbio.2000234
- 1327 RUIZ-JONES, L. J. & PALUMBI, S. R. (2017). Tidal heat pulses on a reef trigger a fine-tuned
1328 transcriptional response in corals to maintain homeostasis. *Science Advances* **3**,
1329 e1601298. 10.1126/sciadv.1601298
- 1330 RUNDLE, H. D. & NOSIL, P. (2005). Ecological speciation. *Ecology Letters* **8**(3), 336-352.
1331 10.1111/j.1461-0248.2004.00715.x
- 1332 SCHLUTER, D. (2001). Ecology and the origin of species. *Trends in ecology & evolution* **16**(7),
1333 372-380.
- 1334 SCHMIDT, T. L., THIA, J. A. & HOFFMANN, A. A. (2023). How can genomics help or hinder wildlife
1335 conservation? *Annual Review of Animal Biosciences* **12**. 10.1146/annurev-animal-
1336 021022-051810
- 1337 SCHMIDT-ROACH, S., LUNDGREN, P., MILLER, K. J., GERLACH, G., NOREEN, A. M. E. & ANDREAKIS, N. (2012).
1338 Assessing hidden species diversity in the coral *Pocillopora damicornis* from Eastern
1339 Australia. *Coral Reefs* **32**(1), 161-172. 10.1007/s00338-012-0959-z
- 1340 SCUCCHIA, F., ZASLANSKY, P., BOOTE, C., DOHENY, A., MASS, T. & CAMP, E. F. (2023). The role and risks
1341 of selective adaptation in extreme coral habitats. *Nature Communications* **14**(1), 4475.
1342 10.1038/s41467-023-39651-7

- 1343 SHEETS, E. A., WARNER, P. A. & PALUMBI, S. R. (2018). Accurate population genetic measurements
1344 require cryptic species identification in corals. *Coral Reefs* **37**(2), 549-563.
1345 10.1007/s00338-018-1679-9
- 1346 SHILLING, E. N., ECKERT, R. J., STURM, A. B. & VOSS, J. D. (2023). *Porites astreoides* coral populations
1347 demonstrate high clonality and connectivity in southeast Florida. *Coral Reefs* **42**,
1348 1131–1145. 10.1007/s00338-023-02417-0
- 1349 SHINZATO, C., MUNGPAKDEE, S., ARAKAKI, N. & SATOH, N. (2015). Genome-wide SNP analysis
1350 explains coral diversity and recovery in the Ryukyu Archipelago. *Scientific Reports* **5**,
1351 18211. 10.1038/srep18211
- 1352 SOUSA, V. C. & HEY, J. (2013). Understanding the origin of species with genome-scale data:
1353 modelling gene flow. *Nature* **14**(6), 404-414. 10.1038/nrg3446
- 1354 SOUTER, D., PLANES, S., WICQUART, J., OBURO, D. & STAUB, F. (2021). Status of coral reefs of the
1355 world: 2020 report. Global Coral Reef Monitoring Network (GCRMN) and International
1356 Coral Reef Initiative (ICRI).
- 1357 SOUTER, P. (2010). Hidden genetic diversity in a key model species of coral. *Marine Biology*
1358 **157**(4), 875-885. 10.1007/s00227-009-1370-3
- 1359 STARKO, S., FIFER, J. E., CLAAR, D. C., DAVIES, S. W., CUNNING, R., BAKER, A. C. & BAUM, J. K. (2023).
1360 Marine heatwaves threaten cryptic coral diversity and erode associations among
1361 coevolving partner. *Science Advances* **9**(32), eadf0954.
- 1362 STURM, A. B., ECKERT, R. J., CARREIRO, A. M., SIMÕES, N. & VOSS, J. D. (2022). Depth-dependent
1363 genetic structuring of a depth-generalist coral and its symbiodiniaceae algal
1364 communities at Campeche Bank, Mexico. *Frontiers in Marine Science* **9**, 835789.
1365 10.3389/fmars.2022.835789
- 1366 STURM, A. B., ECKERT, R. J., MENDEZ, J. G., GONZALEZ-DIAZ, P. & VOSS, J. D. (2020). Population genetic
1367 structure of the great star coral, *Montastraea cavernosa*, across the Cuban
1368 archipelago with comparisons between microsatellite and SNP markers. *Scientific*
1369 *Reports* **10**(1), 15432. 10.1038/s41598-020-72112-5
- 1370 TAYLOR, S. A. & LARSON, E. L. (2019). Insights from genomes into the evolutionary importance
1371 and prevalence of hybridization in nature. *Nature Ecology & Evolution* (3), 170-177.
1372 10.1038/s41559-018-0777-y
- 1373 THIA, J. A. (2022). Guidelines for standardizing the application of discriminant analysis of
1374 principal components to genotype data. *Molecular Ecology Resources* **23**, 523-538.
1375 10.1111/1755-0998.13706
- 1376 THOMAS, L., ROSE, N. H., BAY, R. A., LÓPEZ, E. H., MORIKAWA, M. K., RUIZ-JONES, L. & PALUMBI, S. R.
1377 (2018). Mechanisms of thermal tolerance in reef-building corals across a fine-grained
1378 environmental mosaic: lessons from Ofu, American Samoa. *Frontiers in Marine*
1379 *Science* **4**, 434. 10.3389/fmars.2017.00434
- 1380 THOMAS, L., UNDERWOOD, J. N., ADAM, A. A. S., RICHARDS, Z. T., DUGAL, L., MILLER, K. J. & GILMOUR, J.
1381 P. (2019). Contrasting patterns of genetic connectivity in brooding and spawning
1382 corals across a remote atoll system in northwest Australia. *Coral Reefs* **39**(1), 55-60.
1383 10.1007/s00338-019-01884-8
- 1384 TSUCHIYA, K., ZAYASU, Y., NAKAJIMA, Y., ARAKAKI, N., SUZUKI, G., SATOH, N. & SHINZATO, C. (2022).
1385 Genomic analysis of a reef-building coral, *Acropora digitifera*, reveals complex
1386 population structure and a migration network in the Nansei Islands, Japan. *Molecular*
1387 *Ecology* **31**(20), 5270-5284. 10.1111/mec.16665
- 1388 TURNHAM, K. E., ASCHAFFENBURG, M. D., PETTAY, D. T., PAZ-GARCIA, D. A., REYES-BONILLA, H., PINZON,
1389 J., TIMMINS, E., SMITH, R. T., MCGINLEY, M. P., WARNER, M. E. & LAJEUNESSE, T. C. (2023). High

1390 physiological function for corals with thermally tolerant, host-adapted symbionts.
1391 *Proc Biol Sci* **290**(2003), 20231021. 10.1098/rspb.2023.1021

1392 UNDERWOOD, J. N., RICHARDS, Z., BERRY, O., OADES, D., HOWARD, A. & GILMOUR, J. P. (2020). Extreme
1393 seascape drives local recruitment and genetic divergence in brooding and spawning
1394 corals in remote north-west Australia. *Evolutionary Applications* **13**, 2404-2421.
1395 10.1111/eva.13033

1396 VAN OPPEN, M. J. & GATES, R. D. (2006). Conservation genetics and the resilience of reef-building
1397 corals. *Molecular Ecology* **15**(13), 3863-83. 10.1111/j.1365-294X.2006.03026.x

1398 VAN OPPEN, M. J. H., BONGAERTS, P., FRADE, P. R., PELOW, L. M., BOYD, S. E., NIM, H. T. & BAY, L. K.
1399 (2018). Adaptation to reef habitats through selection on the coral animal and its
1400 associated microbiome. *Molecular Ecology* **27**(14), 2956-2971. 10.1111/mec.14763

1401 VAN OPPEN, M. J. H., OLIVER, J. K., PUTNAM, H. M. & GATES, R. D. (2015). Building coral reef
1402 resilience through assisted evolution. *Proceedings of the National Academy of
1403 Sciences of the United States of America* **112**(8), 2307-2313.
1404 10.1073/pnas.1422301112

1405 VERON, J. E. N. (1995). *Corals in Space and Time: The Biogeography and Evolution of the
1406 Scleractinia*. UNSW Press, Sydney.

1407 VOOLSTRA, C. R., HUME, B. C. C., ARMSTRONG, E. J., MITUSHASI, G., PORRO, B., OURY, N., AGOSTINI, S.,
1408 BOISSIN, E., POULAIN, J., CARRADEC, Q., PAZ-GARCÍA, D. A., ZOCCOLA, D., MAGALON, H., MOULIN,
1409 C., BOURDIN, G., IWANKOW, G., ROMAC, S., BANAIGS, B., BOSS, E., BOWLER, C., DE VARGAS, C.,
1410 DOUVILLE, E., FLORES, M., FURLA, P., GALAND, P. E., GILSON, E., LOMBARD, F., PESANT, S.,
1411 REYNAUD, S., SULLIVAN, M. B., SUNAGAWA, S., THOMAS, O. P., TROUBLÉ, R., THURBER, R. V.,
1412 WINCKER, P., PLANES, S., ALLEMAND, D. & FORCIOLI, D. (2023). Disparate genetic divergence
1413 patterns in three corals across a pan-Pacific environmental gradient highlight species-
1414 specific adaptation. *npj Biodiversity* **2**(1). 10.1038/s44185-023-00020-8

1415 WAPLES, R. S. & GAGGIOTTI, O. E. (2006). What is a population? An empirical evaluation of some
1416 genetic methods for identifying the number of gene pools and their degree of
1417 connectivity. *Molecular Ecology* **15**(6), 1419-1439. 10.1111/j.1365-294X.2006.02890.x

1418 WARNER, P. A., VAN OPPEN, M. J. H. & WILLIS, B. L. (2015). Unexpected cryptic species diversity in
1419 the widespread coral *Seriatopora hystrix* masks spatial-genetic patterns of
1420 connectivity. *Molecular Ecology* **24**, 2993–3008. 10.1111/mec.13225

1421 WEPFER, P. H., NAKAJIMA, Y., SUTTHACHEEP, M., RADICE, V. Z., RICHARDS, Z., ANG, P., TERRANE, T.,
1422 SUDEK, M., FUJIMURA, A., TOONEN, R. J., MIKHEYEV, A. S., ECONOMO, E. P. & MITARAI, S. (2020).
1423 Evolutionary biogeography of the reef-building coral genus *Galaxea* across the Indo-
1424 Pacific ocean. *Mol Phylogenet Evol* **151**, 106905. 10.1016/j.ympev.2020.106905

1425 WILLIS, B. L., VAN OPPEN, M. J. H., MILLER, D. J., VOLLMER, S. V. & AYRE, D. J. (2006). The role of
1426 hybridization in the evolution of reef corals. *Annual Review of Ecology, Evolution, and
1427 Systematics* **37**(1), 489-517. 10.1146/annurev.ecolsys.37.091305.110136

1428 ZAYASU, Y., TAKEUCHI, T., NAGATA, T., KANAI, M., FUJIE, M., KAWAMITSU, M., CHINEN, W., SHINZATO, C.
1429 & SATOH, N. (2021). Genome-wide SNP genotyping reveals hidden population structure
1430 of an acroporid species at a subtropical coral island: Implications for coral restoration.
1431 *Aquatic Conservation: Marine and Freshwater Ecosystems* **31**(9), 2429-2439.
1432 10.1002/aqc.3626

1433 ZHANG, J., RICHARDS, Z. T., ADAM, A. A. S., CHAN, C. X., SHINZATO, C., GILMOUR, J., THOMAS, L.,
1434 STRUGNELL, J. M., MILLER, D. J. & COOKE, I. (2023). Evolutionary responses of a reef-
1435 building coral to climate change at the end of the last glacial maximum. *Molecular
1436 Biology and Evolution* **39**, p.msac201. 10.1093/molbev/msac201

1437

1438

Appendix

1439 ***Structured review for population genomic studies***

1440 An initial search of Web of Science on 21/10/2022, using the search terms "(TI=(coral) OR
1441 TI=(scleractinia*) NOT TI=(fish)) AND (AB=(rad*) OR ALL=(snp*))". The search returned 802
1442 studies. Titles and abstracts were skimmed, and many irrelevant papers were excluded. Any
1443 manuscript that appeared to contain population genomic data of scleractinians or
1444 octocorallians was then retained for manual inspection.

1445

1446

1447 Each manuscript was read and evaluated by two people to ensure that the genomic data: i)
1448 pertained to the cnidarian coral host, ii) used many loci on a genomic scale (i.e., not
1449 microsatellites, not metabarcoding), iii) surveyed two or more sites, and iv) presented results
1450 that included ordination based on individual genotypes (principal components analysis,
1451 principal coordinates analysis or multidimensional scaling) and/or unsupervised model based
1452 clustering test (such as ADMIXTURE or STRUCTURE). We did not consider papers that only
1453 reported discriminant analysis of principal components, as DAPC finds the eigenvectors that
1454 best differentiate prespecified groups, whereas PCA, PCOA and MDS find eigenvalues that
1455 best capture total diversity regardless of group membership (see (Thia, 2022) for further
1456 discussion). If these four criteria were not met, the paper was excluded. For each retained
1457 paper, the two evaluators independently extracted various attributes from the study (see raw
1458 data) and reconciled any discrepancies between their scoring through discussion. Despite
1459 attempting to undertake a rigorous and inclusive search, the authors noticed that several
1460 suitable manuscripts were missing and therefore on July 18, 2023 we ran an ad hoc search in
1461 WOS based on authors that are known to be publishing on population genomics of corals
1462 (namely Barshis, DJ; Baums, IB; Bay, LK; Bongaerts, P; Cooke, I; Matz, MV; Palumbi, SR;
1463 Richards, ZT, Underwood, JN, van Oppen, MJH) and also repeated the the above search with
1464 exactly the same criteria for articles published since 21/10/2022. These new papers were
1465 evaluated as above.

1466

1467 The initial search yielded 853 papers that were reduced to 27 after skimming titles and
1468 abstracts, and 25 were found to be suitable for data extraction. The ad hoc search (combining
1469 authors and new publications) initially identified 897 papers in WOS that were reduced to 16
1470 papers once titles and abstracts were skimmed, and only 11 were suitable for data extraction.

1471

1472 ***Heat stress studies***

1473 The CORDAP database (Ortiz *et al.*, 2023) was downloaded and searched on 11/09/2023,
1474 searching for any records which used genome-wide data of the coral host (i.e., coral SNP data)
1475 and those which conducted clustering analyses (e.g., PCA, ADMIXTURE, or STRUCTURE). First,

1476 records were filtered based on whether the database columns “Host_genotype”,
1477 “Symbiodiniaceae_genotype”, and “Microbiome_genotype” were listed as “TRUE”. Next, the
1478 columns “Symbiodiniaceae_genotyping_approach”, “Host_genotyping_approach” and
1479 “Microbiome_genotyping_approach” were interrogated and only records where the
1480 genotyping method was listed as RADseq, WGS or RNAseq were kept for further checks. The
1481 titles and abstracts of the remaining records were skimmed, and only records for which at
1482 least host genotyping was performed were included. The remaining records were read and
1483 evaluated to ensure that they i) pertained to the cnidarian coral host, ii) used many loci on a
1484 genomic scale (i.e., not microsatellites, not metabarcoding, but single nucleotide
1485 polymorphisms), iii) presented results that included ordination based on individual genotypes
1486 (principal components analysis, principal coordinates analysis or multidimensional scaling)
1487 and/or an assignment test (such as ADMIXTURE or STRUCTURE). Additionally, the number of
1488 cryptic species assigned in each paper by the original authors was noted, as well as the
1489 evaluators’ interpretation of the number of cryptic species based on the plots and analyses
1490 included (where possible).

1491
1492 The original database consisted of 562 records, many of which did not include host,
1493 Symbiodiniaceae or microbiome genotyping. The initial filtering of the database for records
1494 that included some aspect of host genotyping yielded 223 results. Applying criteria i & ii by
1495 skimming abstracts and methods reduced the studies to 17 records. Of these 17 studies, nine
1496 studies included either ordination or model-based clustering analyses (e.g., PCA,
1497 ADMIXTURE). Of these, three (Rose *et al.*, 2017; Ruiz-Jones & Palumbi, 2017; Rose *et al.*, 2021)
1498 the criteria in Section 2.1. The remaining six studies were either ambiguous or showed no
1499 clear evidence for cryptic taxa.