Multi-model inference of non-random mating from an information theoretic approach

Running title: **Multi-model inference of non-random mating**

1 Keywords: Mate choice, mate competition, sexual selection, assortative mating,
2 information criteria, multi-model inference

3

4

Abstract

Non-random mating has a significant impact on the evolution of organisms. Here, I developed a modelling framework for discrete traits (with any number of phenotypes) to explore different models connecting the non-random mating causes (mate competition and/or mate choice) and their consequences (sexual selection and/or assortative mating).

I derived the formulas for the maximum likelihood estimates of each model and used information criteria to perform multimodel inference. Simulation results showed a good performance of both model selection and parameter estimation. The methodology was applied to ecotypes data of the marine gastropod *Littorina saxatilis* from Galicia (Spain), to show that the mating pattern is better described by models with two parameters that involve both mate choice and competition, generating positive assortative mating plus female sexual selection.

As far as I know, this is the first standardized methodology for model selection and multimodel inference of mating parameters for discrete traits. The advantages of this framework include the ability of setting up models from which the parameters connect causes, as mate competition and mate choice, with their outcome in the form of data patterns of sexual selection and assortative mating. For some models, the parameters may have a double effect i.e. they produce sexual selection and assortative mating, while for others there are separated parameters for one kind of pattern or another.

From an empirical point of view, it is much easier to study patterns than processes and, for this reason, the causal mechanisms of sexual selection are not so well known as the patterns

they produce. The goal of the present work is to propose a new tool that helps to distinguish among different alternative processes behind the observed mating pattern.

The full methodology was implemented in a software called InfoMating (available at http://acraaj.webs6.uvigo.es/InfoMating/Infomating.htm).

1. Introduction

The concept of sexual selection is a key piece of modern evolutionary theory as it explains a great range of evolutionary patterns and diversity. Darwin (1871, 1974) originally defined sexual selection as competition between individuals of one sex to achieve matings with the other sex. Yet Darwin distinguished two general biological mechanisms of sexual selection: mate competition and mate choice (see Ng et al. 2019 and references therein). However, the concept of sexual selection has been controversial since its very beginning (reviewed in Andersson 1994; Prum 2012; Parker 2014; Parker and Pizzari 2015) and there is still disagreement on the actual definition of sexual selection (Fitze and Galliard 2011), and even, its role as a key component of modern evolutionary biology has being confronted (Roughgarden et al. 2006; but see Shuker 2010; Parker and Pizzari 2015).

It seems that some of the disagreements and misunderstandings about sexual selection and related concepts, come from the distinct emphases that scientific fields (e.g. population genetics, speciation theory, behavioral ecology and sociology) put on the various aspects of the sexual selection theory. To make things worse, sexual selection is described sometimes as a process and sometimes as a pattern. Thus, the meaning and usage of the term sexual

selection may vary depending on the biological question we are asking, e.g. behavior or evolutionary agent, but it also depends on if we are viewing sexual selection as a process or as a pattern.

The distinction between patterns and processes is of key importance in science in general and in evolutionary biology in particular (Mahler et al. 2017). In the end, science is about understanding of processes that underlie the observed patterns (Swihart et al. 2002). The concept of pattern refers to the detection of regularity in nature, while the concept of process refers to the mechanisms creating and maintaining this order (Chapleau et al. 1988). Thus, recognition of patterns is the basis for developing and testing hypotheses about processes.

Sometimes, the distinction between pattern and process may be obscured because of the same biological concept can be meaningfully defined as both a process and a pattern (Armstrong 1977; Mahler et al. 2017). Consider for example, the classical definition of sexual selection, as arising from variation in reproductive success due to competition for access to mates (Andersson 1994; Shuker 2010). From such definition, sexual selection can be considered as the evolutionary agent (a process) that drives the evolution of some mating-related traits. However, from the same definition, if we put the emphasis on the pattern of evolutionary change that arises from the differences in the reproductive success, then we are viewing sexual selection as a pattern caused by some other biological process (competition).

In this work, I adhere to the definition used in population genetics, where sexual selection is caused by processes of mate competition that may produce intrasexual selection, and/or

processes of mate choice that may produce intersexual selection (Lewontin et al. 1968; Endler 1986; Casares et al. 1998; Rolán-Alvarez and Caballero 2000; Ng et al. 2019).

From a population genetics perspective, the causes and consequences of mating systems have often been analyzed separately. This approach facilitated the experimental handling and the study of the evolutionary causes (e.g. mate choice and mate competition, see Futuyma and Kirkpatrick 2017 and references therein) and consequences (e.g. assortative mating and sexual selection, see Arnold and Wade 1984; Rolán-Alvarez and Caballero 2000; Rolan-Alvarez et al. 2015b and references therein). Accordingly, sexual selection is considered an evolutionary pattern regardless of the process that gave rise to it, mate competition or mate choice.

More specifically, the process of mate competition refers in the broad sense to access to matings by courtship, intrasexual aggression and/or competition for limited breeding resources (Andersson 1994; Kokko et al. 2012; Wacker and Amundsen 2014). These processes may generate a pattern of sexual selection (a change in the frequency of the trait under study) in the same sex that competes (intrasexual selection Ng et al. 2019).

The process of mate choice occurs whenever the effects of traits expressed in one sex leads to non-random allocation of reproductive investment with members of the opposite sex (Edward 2015). Choice may be mediated by phenotypic (sensorial or behavioural) properties that affect the propensity of individuals to mate with certain phenotypes (Jennions and Petrie 1997). The observed pattern driven by mate choice can be a change in trait frequency in the other sex (intersexual selection) and/or a pattern of trait correlation between mates (assortative mating).

Still, the relationships among these concepts are complex and can be approached from different perspectives (the reader may consult Arnold and Wade 1984; Rolán-Alvarez and Caballero 2000; Edward 2015; Rolan-Alvarez et al. 2015b; Futuyma and Kirkpatrick 2017; Rosenthal 2017 for extended details and alternative definitions; Estévez et al. 2018; Ng et al. 2019).

Summarizing, the evolutionary consequences of mate competition and mate choice are sexual selection and assortative mating. When the traits under study are discrete, the patterns of sexual selection and assortative mating are defined in terms of change in the phenotype frequencies so that, sexual selection corresponds to the observed change in gene or phenotype frequencies in mated individuals with respect to population frequencies (Hartl and Clark 1997; Rolán-Alvarez and Caballero 2000). Similarly, assortative mating corresponds to the observed deviation from random mating when measured within matings (Rolán-Alvarez and Caballero 2000 and references therein).

In a given population, mating is random when an individual of one sex has an equal chance of mating with any other individual of the opposite sex. Thus, random mating implies that the overall frequency of mating between female and male types is equal to the product of frequencies of these types in the population (Gavrilets 2004). As mentioned, deviation from random mating can be observed in the form of sexual selection and/or assortative mating patterns and may occur as a consequence of mate competition and/or mate choice.

In a previous work (Carvajal-Rodríguez 2018b), the processes of mate competition and mate choice were modelled for discrete traits by means of the parameters m_{ij}, that represent the mutual mating propensity between a female of type *i* and a male *j*. Therefore, if A-type

females prefer A-type males, this mate choice is modelled as a higher mutual mating propensity between these types as compared with the mutual mating propensity of the A females with other male types ($m_{AA} > m_{AB}$). On the other hand, if B-type males mate more often than other males whatever the female, this mate competition is modelled by a higher marginal mating propensity of such males (see below).

By modelling the mating process as a differential mutual mating propensity among different types of mating pairs, it is possible to express the mean change in mating phenotypes as the information gained due to non-random mating (Carvajal-Rodríguez 2018b). Expressing the patterns obtained from mate choice and competition in terms of the information captured in the mutual mating propensity models, permitted to identify the necessary and sufficient conditions for describing random mating as the zero information model.

Thus, the mating information-based framework provides a formal approach for developing a set of hypotheses about the causes (mate competition and mate choice) and the patterns they may provoke (sexual selection and assortative mating). In addition, data-based evidence can be used for ranking each hypothesis and perform multi-model-based inference (Link and Barker 2006; Burnham et al. 2011; Aho et al. 2014).

In the following sections I proceed as follows:

1.- Given the population frequencies for some discrete trait I define the multinomial saturated mating model in terms of the mutual mating propensity parameters. The maximum-likelihood estimates of these parameters are the pair total indices (PTI) as defined in (Rolán-Alvarez and Caballero 2000). Once the saturated model is defined I

obtain the three necessary and sufficient conditions for random mating. Afterwards, by relaxing these conditions it is possible to generate models for which differential marginal mating propensity may produce female or male sexual selection without assortative mating, or on the contrary, models for which some mutual mating propensities represent mate choice that may produce assortative mating and frequency dependent sexual selection. I obtain the maximum likelihood estimates for the parameters of these models.

2.- Relying on the previous section, it is possible to generate several mutual mating propensity models and apply information criteria for selecting the best candidate ones and estimating the mating parameter values based on the most supported models. I developed a software called InfoMating to do so.

3.- Finally, I demonstrate the methodology by analysing simulated and real data.

2. Mutual mating propensity models

Consider a female trait with k_1 different phenotypes and a male trait with k_2 phenotypes, the total number of possible mating phenotypes is $K = k_1 \times k_2$. Let a sample have n' matings from which n'_{ij} correspond to i-type females that mated with j-type males. If the probability of the mating $i \times j$ is q'_{ij}, then the logarithm of the multinomial likelihood function of the sample is

$$lnL = C + \sum_{ij}^{K} n'_{ij} \ln(q'_{ij})$$

where C is the multinomial coefficient which is constant given the sample. As it is well-known, the maximum likelihood estimator of the multinomial probability of the mating $i \times j$ is n'_{ij} / n'.

2.1 Saturated non-random mating model M_{sat}

Let the population under study have n_{1i} females of type i from a total of n_1 females and n_{2j} males of type j from a total of n_2 males. Therefore, the population frequency of females of type i is $p_{1i} = n_{1i} / n_1$ and the population frequency of males of type j is $p_{2j} = n_{2j} / n_2$.

The mating probability between types i and j can be expressed as $q'_{ij} = m_{ij}q_{ij}$ (Carvajal-Rodríguez 2018b) where q_{ij} is the product of the female and male population frequencies of each type ($q_{ij} = p_{1i} \times p_{2j}$) and $m_{ij} = m'_{ij} / M$, where m'_{ij} is the mutual mating propensity, i.e. the expected number of matings given an encounter between females of type i and males of type j, and M is the mean mutual mating propensity $M = \sum_{i,j} q_{ij} m'_{ij}$, so that $\Sigma q'_{ij} = 1$.

Under this multinomial model, the log-likelihood of the sample is

$$lnL_{sat} = C + \sum_{ij}^{K} n'_{ij} \ln(m_{ij} q_{ij}) \qquad (1)$$

This model is saturated (M_{sat}) because it has as many parameters as independent mating-class frequencies, $P_{sat} = K - 1$. The female and male population frequencies, p_1 and p_2, are either known or they need to be estimated in all the models. Therefore, for model comparison, the population frequencies can be ignored when counting the number of parameters involved in each model.

The maximum likelihood estimate (MLE) of m_{ij} is $(n'_{ij} / n')/q_{ij}$ = $PTI_{i,j}$ where $PTI_{i,j}$ is the pair total index i.e. the frequency of the observed mating classes divided by their expected frequency under random mating (Rolán-Alvarez and Caballero 2000).

In this work I am interested in the estimation of the mutual mating propensity parameters (from hereafter mutual-propensity parameters) for various competition and mate choice models. From that point of view, it is convenient to express the maximum likelihood estimator in a different way which I call λ-notation.

2.2 λ-notation

Consider the non-normalized parameters m'_{ij} and recall that $m_{ij} = m'_{ij} / M$. The MLE of m'_{ij} under M_{sat} is simply $M \times PTI_{i,j}$ i.e. $M \times (n'_{ij} / n') / q_{ij}$ that can be conveniently rearranged as $(n'_{ij} / q_{ij}) / (n' / M)$. Because the mating parameters are normalized, it is possible, without loss of generality, to set just one of the m'_{ij} to an arbitrary value of 1. Thus, let set $m'_{k1k2} = 1$ and note (details in Appendix A) that in such case $n' / M = n'_{k1k2} / q_{k1k2}$. Therefore, the MLE of the parameters of the saturated model can be expressed as

$$\widehat{m}'_{ij} = \frac{\lambda\left(m'_{ij}\right)}{\lambda(1)}$$

where

$$\lambda(\theta) = \frac{\sum_{ij}^{A} n'_{ij}}{\sum_{ij}^{A} q_{ij}} \qquad (2)$$

i.e., the function λ of a mating parameter θ is the sum of the counts of all the mating classes in the set $A = \{(i_1, j_1), ...\}$ having mutual-propensity θ divided by the sum of their expected frequencies under random mating.

Thus, $\lambda(m'_{ij})$ expresses the sum of the observed matings with mutual-propensity m'_{ij} divided by the product of the population frequencies from each partner type. Similarly, $\lambda(1)$ corresponds to the sum of the observed matings having unity mating parameter divided by the corresponding products of population frequencies.

As already mentioned, the most parameterized model is the saturated model that has K-1 parameters so, when divided by the mean mutual-propensity M, the estimates $\lambda(m'_{ij}) / (M\lambda(1))$ are the corresponding pair total indices (PTI_{ij}).

The model M_{sat} is the most complex model that can be fitted to the available data. The principle of parsimony suggests to consider reduced special cases of this saturated model. Next, I computed the ML estimates of different classes of reduced models that require less parameters, beginning by the most reduced one which is the random mating model.

2.3 Random mating model M_0

The random model M_0 corresponds to the simplest, most reduced model, which is nested within all others (it is a particular case of any other model) while it is not possible to derive any simplified version from it. When random mating occurs, the mating probability between types i and j is $q'_{ij} = q_{ij} = p_{1i} \times p_{2j}$. Under this model, the information indices would be zero (Carvajal-Rodríguez 2018b). This zero-information model is a particular case of the

saturated one when the mutual-propensities are equal for every mating phenotype. The number of independent mating parameters is $P_0 = 0$.

The log-likelihood of the sample of mating is

$$lnL_0 = C + \sum_{ij}^{K} n'_{ij} \ln(q_{ij}) \qquad (3)$$

Now, let's define the marginal propensity m_{Fem_i} for a female of type i as

$$m_{Fem_i} = \sum_{j}^{males} p_{2j} \frac{m'_{ij}}{M} = \sum_{j}^{males} p_{2j} m_{ij} \qquad (4)$$

Similarly for a male of type j

$$m_{Male_j} = \sum_{i}^{females} p_{1i} \frac{m'_{ij}}{M} = \sum_{i}^{females} p_{1i} m_{ij}$$

Then, the M_0 model corresponds to M_{sat} subjected to the following restrictions:

i) Equal female marginals: $m_{Fem_i} = m_{Fem_j} \forall i, j$ (5)

ii) Equal male marginals: $m_{Male_i} = m_{Male_j} \forall i, j$

iii) Multiplicativity: $m_{ij} = m_{Fem_i} \times m_{Male_j} \forall i, j$

It is useful to express M_0 in terms of these three restrictions because by relaxing some of them it is possible to define different classes of models. For example, a model with equal female marginal propensities and multiplicative mutual-propensities (conditions i and iii hold) but different male marginal propensities (relaxing ii), corresponds to a case with competition among males that may provoke a (intra)sexual selection pattern (see below).

Therefore, by relaxing some of the conditions in (5), it is possible to control the kind of causes that produce the different non-random mating patterns. In fact, there are three general classes of models that can be combined. The two first classes correspond to relaxing the first or second condition and involve mate competition in females or males, provoking female or male (intra)sexual selection respectively. Provided that the third condition is maintained, these models cannot produce an assortative mating pattern (see below). The third class corresponds to relaxing the third condition and involves mate choice, which may provoke just assortative mating, or both assortative mating and sexual selection, the latter depending on the population phenotype frequencies (Fig. 1).



Fig 1. Mating models defined by mate competition or mate choice, and their effect after relaxing some of the conditions imposed to the random mating model M_0.

2.4 Mate competition models

These class of models correspond to relaxing the first and/or second conditions in M_0 while maintaining the condition of multiplicativity (5-iii). The maintenance of the third condition implies that the mutual-propensity of a mating pair (i,j) is the product of the marginal female (m_{Fem}) and male (m_{Male}) propensities. Under this condition there should be no deviation from random mating when comparing the observed and expected frequencies within matings and the assortative mating pattern should not be observed (Carvajal-Rodríguez 2018b). I distinguished models that generate a sexual selection pattern in just one sex or in both.

2.4.1 Intra-female competition

Relaxing condition (5-i) implies that at least one female marginal propensity, say female of type A, is different from the rest of female types i.e. $m_{Fem_A} \neq m_{Fem_B}$ with $A \neq B$. On the other side, the marginal propensity of males should be the same which means that there is no intra-male competition, all male types mate at an equal rate.

Therefore, a model with intra-female competition is obtained by defining every mutual-propensity involving a female of type i, by an absolute (unnormalized) mating parameter a_i as follows

$$m'_{11} = m'_{12} = \ldots = m'_{1k2} = a_1$$

$$m'_{21} = m'_{22} = \ldots = m'_{2k2} = a_2$$

.

.

.

$$m'_{(k1-1)1} = m'_{k12} = \ldots = m'_{k1k2} = a_{k1-1}$$

$$m'_{k11} = m'_{k12} = \ldots = m'_{k1k2} = 1 \qquad (6)$$

with $a_i > 0 \ \forall i$.

Note that the relationships among the parameters will not be altered when dividing them by a_{k1} so that $a_{k1} = 1$. Under this model, there can be as much as k_1-1 free mating parameters.

When computing the female and male marginal propensities (4) it is seen that

$$m_{Fem_1} = a_1/M;\ m_{Fem_2} = a_2/M \ldots;\ m_{Fem_k1} = 1/M$$

$$m_{Male_1} = m_{Male_2} = m_{Male_3} = \ldots = m_{Male_k2} = M/M = 1$$

where M is the mean mutual-propensity as defined above.

The model (6) has equal male marginal propensity and it is multiplicative. The MLE of the parameters is

$$\hat{a}_i = \frac{\lambda(a_i)}{\lambda(1)} \qquad (7)$$

where $\lambda(\theta)$ is defined as in (2). Thus, $\lambda(a_i)$ expresses the sum of the observed matings having mutual-propensity a_i, divided by the sum of the product of the population frequencies from each partner type. Similarly, $\lambda(1)$ corresponds to the sum of the observed matings having unity mating parameter divided by the sum of the corresponding products of population frequencies (details in Appendix A).

2.4.2 Intra-male competition

Relaxing condition (5-*ii*) implies that at least one male marginal propensity, say male of type *A*, is different from the rest of male types i.e. $m_{Male_A} \neq m_{Male_B}$ with $A \neq B$. On the other side, the marginal propensity of females should be the same which means that there is no intra-female competition, all female types mate at an equal rate. The corresponding model can be obtained just by interchanging rows with columns in (6). Noting the parameters as b_j instead of a_i, the maximum likelihood estimate is

$$\hat{b}_j = \frac{\lambda(b_j)}{\lambda(1)} \qquad (8).$$

2.4.3 Intra-female and male competition

By relaxing conditions (5-*i*) and (5-*ii*) the marginal propensities will be different within females and males. The corresponding model combines models (6) and (8) and has as much as $(k_1\text{-}1)\times(k_2\text{-}1)$ parameters in the most parameterized case, and a minimum of two (female and male) for the less parameterized one, in order to maintain the multiplicativity condition (5-*iii*). This type of model may produce a pattern of sexual selection in both sexes without assortative mating. By notational convenience, I fix the category k_1 in females and k_2 in males as having unitary parameters. Therefore

$$m'_{ij} = a_i b_j,\ i<k_1,\ j<k_2;\ m'_{ik2} = a_i,\ i<k_1;\ m'_{k1j} = b_j;\ j<k_2,\ m'_{k1k2} = 1;\ \text{with } a_i > 0,\ b_j > 0$$

$\forall i, j$.

This model is multiplicative (see Appendix A) and the parameters MLE are

$$\hat{a}_i = \left(\frac{p_{1k1}}{p_{1i}}\right)\frac{\sum_{j=1}^{k2} n'_{ij}}{\sum_{j=1}^{k2} n'_{k1j}} = \frac{\lambda(a_i+\sum_j a_i b_j)}{\lambda_{fem}(1)} \qquad (9)$$

303 $$\hat{b}_j = \left(\frac{p_{2k2}}{p_{2j}}\right)\frac{\sum_{i=1}^{k1} n'_{ij}}{\sum_{i=1}^{k1} n'_{ik2}} = \frac{\lambda(b_j+\sum_i a_i b_j)}{\lambda_{male}(1)}$$

304 where the lambda function $\lambda(a_i + a_ib_1 + ... a_ib_j + ...)$ is applied to the mutual-propensities
305 that depend on the parameter a_i. Thus, $\lambda(a_i + \sum_j a_ib_j)$ is the quotient between the sum of the
306 number of observed mating phenotypes that depend on the parameter a_i (i.e. $\sum_j n'_{ij}$) and the
307 sum of their expected random mating frequencies (which is simply p_{1i}). Correspondingly,
308 $\lambda_{sex}(1)$ is the quotient between the sum of cases that contribute with 1 to the mutual-
309 propensity by the given sex (i.e. $\sum_j n'_{k1j}$ for females) and the sum of the expected
310 frequencies (which is p_{1k1} for females). Formulae (9) is similar to (7) and (8). Note that the
311 model in (9) becomes (7) by fixing every b_j as 1 while it becomes (8) by fixing every a_i as
312 1. The percentage of sexual selection information corresponding to each sex (JS1 and JS2
313 in Carvajal-Rodríguez 2018b), would depend on the population frequencies and on the
314 mating parameter values.

315

316 *2.5 Mate choice models*

317 Mate choice models correspond to the class of non-multiplicative models, i.e. they can be
318 obtained by relaxing the condition (5-iii) and may produce assortative mating patterns
319 (positive or negative). If the female marginal propensities are equal and the same is true for
320 the males (conditions 5-i and 5-ii hold) there would not be sexual selection neither in
321 females nor males, and the model may produce only assortative mating patterns. However,
322 this cannot be guaranteed in general because the occurrence of the sexual selection pattern
323 is frequency dependent under non-multiplicative models (see below).

Consider a model where the unnormalized mutual-propensities are

$$m'_{ii} = a_i > 0 \ \forall i \in min\{k_1, k_2\} \text{ and } m'_{ij} = 1 \text{ for } i \neq j.$$

Thus, the homotype ($i\times i$) mutual-propensities are parameterized while the heterotype are not. This model is non-multiplicative in general, because the contribution of the type i to the mutual-propensity is distinct in m_{ii} that in m_{ij} or in m_{ji} (although with an even number of types a multiplicative model can be obtained by setting $m'_{ii} = 1 / m'_{jj}$).

By recalling the definition of marginal propensities in (4), the condition for equal female marginal $m_{Fem_i} = m_{Fem_j}$ is

$$p_{2i}(a_i - 1) = p_{2j}(a_j - 1) \qquad (10)$$

and in males

$$p_{1i}(a_i - 1) = p_{1j}(a_j - 1)$$

In general, depending on the conditions in (10), the mate choice models have double effect i.e. they produce assortative pattern jointly with sexual selection in at least one sex.

The maximum likelihood estimate for the model parameters is

$$\hat{a}_i = \frac{\lambda(a_i)}{\lambda(1)} \qquad (11)$$

Note that the homotype mating parameter may imply higher mutual-propensity than the heterotype ($a_i > 1$, positive assortative mating) or viceversa, the homotype has lower mutual-propensity ($a_i < 1$, negative assortative). The number of different parameters ranges from 1 ($a_1 = a_2 = ... = a_i$) to H; where $H = min\{k_1, k_2\}$ corresponds to the maximum possible number of different homotype matings.

It is also possible to define mate choice models with the heterotype mutual-propensities parameterized instead of the homotype ones (see Appendix A for details).

2.6 Models with mate competition and mate choice parameters

I have shown that mate choice models may generate both kind of patterns, assortative mating and sexual selection, depending on the within sex population frequencies. While it is not possible to assure that the mate choice model produces no sexual selection, it is possible to combine the previous models to ensure that there are parameters directly linked to mate competition and parameters directly linked to mate choice. These combined models have the property that when the mate choice parameter is set to 1, there is only a known sexual selection effect as caused by the competition parameter (female, male, or both). When the mate choice parameter is added, the assortative mating pattern appears and also, an extra effect of frequency-dependent sexual selection may be added to that of the original competition parameter.

2.6.1 Models with male competition and mate choice: independent parameters

Consider the model $m'_{i1} = \alpha$; $m'_{ii} = c$ for $i \neq 1$ and $m'_{ij} = 1$ otherwise; with $i \leq k_1$, $j \leq k_2$. An example of this kind of model can be seen in Fig. 2.



Fig 2. An example of male competition and mate choice independent parameters model with 5 × 5 mating phenotypes. α is the male competition parameter and *c* is the choice parameter. Rows are females, columns are males.

For the particular case of $\alpha \neq 1$, $c = 1$; the model has within male competition that corresponds to the marginal propensity α of the type-1 male compared with the other males, so, a male sexual selection pattern may be generated. On the contrary, the female marginal propensities are equal so there is no female competition. Considering mate choice and the assortative mating pattern, when $c = 1$ the model is multiplicative so assortative mating should not be produced, in fact, the pair sexual isolation statistics (PSI) are equal (see Appendix A for details) and the assortative mating is 0, i.e., as measured by the overall index of sexual isolation I_{PSI} ($I_{PSI} = (4\Sigma PSI_{ii} - \Sigma\, PSI_{ij})/(4\Sigma PSI_{ii} + \Sigma\, PSI_{ij})$) see also (Carvajal-Rodríguez 2018b).

However, by taking $c \neq 1$ a new component is added to the sexual selection pattern. The parameter *c* corresponds to mate choice and produces positive ($c > 1$) or negative ($c < 1$) assortative mating. The value of I_{PSI} is a function of the parameter *c* and the population

frequencies. Female sexual selection may also emerge depending on the value of *c* and the population frequencies.

The MLEs of both parameters are

$$\hat{\alpha} = \frac{\lambda(\alpha)}{\lambda(1)}$$

$$\hat{c} = \frac{\lambda(c)}{\lambda(1)}$$

A variant of the above model can be generated by changing the *c* parameter from the main diagonal to the anti-diagonal. Similarly female sexual competition linked to the α-parameter is obtained by transposing the matrix of the model.

2.6.2 Models with male competition and mate choice: compound parameters

Consider the model $m'_{11} = c\alpha$; $m'_{i1} = \alpha$ and $m'_{ii} = c$ for $i > 1$ and $m'_{ij} = 1$ otherwise; with $i \leq k_1$, $j \leq k_2$. An example of this model can be seen in Fig. 3.



Fig 3. An example of male competition and mate choice compound parameters model with 5 × 5 mating phenotypes. α is the male competition parameter and *c* is the choice parameter. Rows are females, columns are males.

When $c = 1$ the model is the same as the previous one. When $c \neq 1$, the mate choice parameter provokes an extra effect of sexual selection in males and females, plus assortative mating. The MLE of α and *c* are

$$\hat{\alpha} = \frac{\sum_{i=1}^{k1} n'_{i1} / (cq_{11} + \sum_{i>1} q_{i1})}{\lambda(1)} = \frac{\lambda_{c.11-}(\alpha)}{\lambda(1)}$$

where $\lambda_{c.11\text{-}}(\alpha)$ indicates that for matings with parameter α, the expected frequency indexed as 11 (i.e. q_{11}) is weighted by *c*. Similarly,

$$\hat{c} = \frac{\sum_{i=1}^{k} n'_{ii} / (\alpha q_{11} + \sum_{i>1} q_{ii})}{\lambda(1)} = \frac{\lambda_{\alpha.11-}(c)}{\lambda(1)}$$

where $k = \min\{k_1, k_2\}$ and $\lambda_{\alpha.11\text{-}}(c)$ indicates that for matings with mating parameter *c*, the expected frequency indexed as 11 is weighted by α.

The above estimates are dependent one on each other, so, for obtaining the estimates of this compound parameter model I have used a numerical bounded Nelder-Mead simplex algorithm, with restriction $\alpha > 0$, $c > 0$ (Press 2002; Singer and Singer 2004; Gao and Han 2012).

2.6.3 General model with male competition and mate choice parameters

The general model with male competition and mate choice parameters is $m'_{11} = c_1\alpha$; $m'_{i1} = \alpha$ and $m'_{ii} = c_k$ for $i > 1$ and $m'_{ij} = 1$ otherwise; with $i \leq k_1$, $j \leq k_2$. A particular case of this model can be seen in Fig. 4.

$$\begin{pmatrix} \alpha c_1 & 1 & 1 & 1 & 1 \\ \alpha & c_1 & 1 & 1 & 1 \\ \alpha & 1 & c_3 & 1 & 1 \\ \alpha & 1 & 1 & c_4 & 1 \\ \alpha & 1 & 1 & 1 & c_5 \end{pmatrix}$$

Fig 4. Male sexual selection and mate choice compound model with 5 × 5 mating phenotypes. α is the male sexual selection parameter and c_k's are the choice parameters with $c_2 = c_1$. Rows are females, columns are males.

Note that to distinguish the competition and mate choice parameters, it is necessary that at least one c_k parameter is equal to c_1 (as in Figs. 3 and 4) or that $c_1 = 1$ as in Fig. 2, otherwise the parameter for m_{11} does not distinguish competition and choice. Therefore, the

model in Fig. 4 has H parameters with $H = \min\{k_1,k_2\}$ from which, $H - 1$ are choice parameters (c) plus one male competition parameter α. The MLE are

$$\hat{\alpha} = \frac{\sum_{i=1}^{k1} n'_{i1} / (c_1 q_{11} + \sum_{i>1} q_{i1})}{\lambda(1)} = \frac{\lambda_{c1.11-}(\alpha)}{\lambda(1)}$$

$$\hat{c}_1 = \frac{n'_{11}/(\alpha q_{11})}{\lambda(1)} = \frac{\lambda_{\alpha.11-}(c_1)}{\lambda(1)}$$

$$\hat{c}_{k>1} = \frac{n'_{kk}/(q_{kk})}{\lambda(1)} = \frac{\lambda(c_k)}{\lambda(1)}$$

The model parameters $c_{k>1}$ can be estimated directly from the sample; on the contrary, the α and c_1 estimates are dependent on each other, so, for obtaining these estimates, I used a numerical bounded Nelder-Mead simplex algorithm with restriction $\alpha > 0$, $c_1 > 0$ (Press 2002; Singer and Singer 2004; Gao and Han 2012).

Previous models were simplified versions of the general model. For example, the model in Fig. 2 is the general model with restrictions $c_1 = 1$; $c_2 = c_3 = \ldots = c_k = c$. Also, the model in Fig. 3 corresponds to $c_1 = c_2 = c_3 = \ldots = c_k = c$. Another particular case that could be defined is $c_1 = c$; $c_2 = c_3 = \ldots = c_k = 1$. In the latter, the MLE of the parameters can again be expressed as a quotient of lambdas similar to the compound parameter case

$$\hat{\propto} = \frac{\lambda_{c1.11-}(\alpha)}{\lambda(1)}$$

$$\hat{c} = \frac{\lambda_{\alpha.11-}(c)}{\lambda(1)}$$

It is also possible to define other general model with the mate choice parameters in the anti-diagonal. Using the λ notation, the estimates follow the same formulae as defined for the

general model with the choice parameters in the main diagonal. Concerning models with female competition and mate choice, they are obtained just by transposing the matrices of the mating parameters.

2.7 General double effect models

The mating parameters $m_{ij} = \theta_{ij}$ with the restriction that at least some are equal to one, permit to generate any particular model. In general, these models produce patterns of sexual selection and assortative mating with each parameter possibly linked to the occurrence of both (see Appendix A). The MLE is

$$\hat{\theta}_{ij} = \frac{\lambda(\theta_{ij})}{\lambda(1)} \qquad (12)$$

The most parameterized model of this kind is the saturated, with K-1 parameters. In such case, as already mentioned, the estimates in (12) are the corresponding pair total indices (PTI).

All the above derived MLE formulae have been verified by numerical approximation using the bounded Nelder-Mead simplex algorithm (Press 2002; Singer and Singer 2004; Gao and Han 2012).The set of described models jointly with their expected effects are summarized in Table 1.

Table 1. Mutual mating propensity models as defined by different parameters in a case with two different phenotypic classes at each sex ($k_1 = k_2 = 2$). The unnormalized m'_{ij} values not explicitly given are assumed to be 1.

Name (abbreviation)	Model	MLE	Effect
Random (M_0)	$m'_{ij} = 1 \forall i,j$		Random mating
Competition Multiplicative Models			
Female competition (S_{Fem}-1P)	$m'_{11} = m'_{12} = a$	$\hat{a} = \lambda(a)/\lambda(1)$	Fem sexual selection
Male competition (S_{Male}-1P)	$m'_{11} = m'_{21} = b$	$\hat{b} = \lambda(b)/\lambda(1)$	Male sexual selection
Female and male competition (S2-2P)	$m'_{11} = ab$		2-sex sexual selection
	$m'_{12} = a$	$\hat{a} = \lambda_{fem}(a)/\lambda_{fem}(1)$	
	$m'_{21} = b$	$\hat{b} = \lambda_{male}(b)/\lambda_{male}(1)$	
Mate Choice Models			
One-parameter (C-1P)	$m'_{11} = m'_{22} = a$	$\hat{a} = \lambda(a)/\lambda(1)$	Assortative mating + sex sel (freqdep)
General mate choice (C-HP)	$m'_{11} = a_1$, $m'_{22} = a_2$	$\hat{a}_i = \lambda(a_i)/\lambda(1)$	Assortative mating + sex sel (freqdep)
Competition and Mate Choice			
2 independent parameters ($S_{Fem}C$-2P)	$m'_{1j} = \alpha$; $m'_{jj} = c$; $j > 1$	$\hat{\alpha} = \lambda(\alpha)/\lambda(1)$ $\hat{c} = \lambda(c)/\lambda(1)$	α-sexual selection in one sex + mate choice effect c-assortative mating
2 parameters (1 compound: $S_{Fem}C$-2Pc)	$m'_{11} = c\alpha$; $j > 1$: $m'_{1j} = \alpha$; $m'_{jj} = c$;	$\hat{\alpha} = \lambda_{c.11\text{-}}(\alpha)/\lambda(1)$ $\hat{c} = \lambda_{\alpha.11\text{-}}(c)/\lambda(1)$	α-sexual selection in one sex + mate choice effect c-assortative mating
H parameters (1 compound: $S_{Fem}C$-HPc)	$m'_{11} = c_1\alpha$; $j > 1$: $m'_{1j} = \alpha$; $m'_{jj} = c_k$;	$\hat{\alpha} = \lambda_{c1.11\text{-}}(\alpha)/\lambda(1)$ $\hat{c}_1 = \lambda_{\alpha.11\text{-}}(c_1)/\lambda(1)$ $\hat{c}_{k\neq 1} = \lambda(c_k)/\lambda(1)$	α-sexual selection in one sex + mate choice effect c_k-assortative mating
General Double Models (D-xP)			Assortative mating + sex sel (freq dep)
Saturated (M_{sat})	$m'_{ij} = a_{ij}$; $m'_{k1k2} = 1$	$\hat{a}_i = \lambda(a_i)/\lambda(1)$	

k_1: number of female categories; k_2: number of male categories; $H = min\{k_1,k_2\}$; sexsel (freqdep): frequency dependent sexual selection.

3. Model Selection and multimodel inference

Relying on the previous section, it would be possible to generate mate competition and mate choice models and, given a mating table, to apply some information criteria for selecting the best-fit candidates and estimating the mating parameter values based on the most supported models. Next, I briefly review the information criteria and model selection concepts and show how to apply them to perform model selection and multimodel inference among mate competition and mate choice models.

Information-based model selection and multi-model inference can be applied to describe uncertainty in a set of models to perform inference on the parameters of interest (Burnham et al. 2011; Grueber et al. 2011; Barker and Link 2015; Claeskens 2016). There are several information criteria at hand, although trusting on a single form of information criterion is unlikely to be universally successful (Liu and Yang 2011; Vrieze 2012; Brewer et al. 2016; Aho et al. 2017; Dziak et al. 2019). In the present work, two Kullback- Leibler divergence-based measures plus the so-called Bayesian information criterion are considered.

3.1. Information criteria

The Akaike information criterion (AIC) provides the link between the Kullback-Leibler divergence and the maximized log-likelihood of a given model (Akaike 1973). Here I use the sample-corrected version AICc, because it is asymptotically equivalent and may work better for small sample size

$$AIC_c = -2ln(L) + 2P_m + (2P_m(P_m+1)) / (n' - P_m - 1)$$

where L is the maximum likelihood of the model, P_m the total number of estimated mating parameters and n' is the number of matings.

There is also a version for the symmetric K-L (Jeffrey's) divergence, called the KICc criterion (Cavanaugh 2004; Keerativibool 2014). It seems adequate to consider the KICc criterion because the mating pattern obtained from the mutual-propensity models can be described by the informational flow from the mating frequencies, in the form of the Jeffrey's divergence (Carvajal-Rodríguez 2018b) so,

$$KIC_c = -2ln(L) + n'\ln(n' / (n' - P_m)) + P_2$$

with $P_2 = n'[(n' - P_m)(2P_m + 3)-2] / [(n' - P_m -2)(n' - P_m)]$

Finally, the Bayesian information criterion (BIC Schwarz 1978) permits an approximation to the Bayes factor applied for model comparison (Wagenmakers 2007)

$$BIC = -2\ln(L) + P_m\ln(n')$$

3.2 Overdispersion

In the context of model selection, data overdispersion, i.e. greater observed variance than expected, could generate the selection of overly complex models. The simplest approach to estimate overdispersion is by computing a single variance inflation factor (v). This inflation factor is the observed variation divided by the expected under the model with the highest likelihood (M_c), other than the saturated, among the proposed ones (Richards 2008; Symonds and Moussalli 2011). It can be asymptotically approximated by the deviance i.e. twice the difference between the log-likelihood of the saturated (M_{sat}) and the M_c model, divided by the difference in the number of parameters ($P_{Msat} - P_{Mc}$) between both models

$$v = 2[\ln(L_{Msat}) - \ln(L_{Mc})] / df$$

where $df = P_{Msat} - P_{Mc}$.

If $1 \leq v \leq 4$ this indicates overdispersion, while if higher than 4-6 this may indicate poor model structure and the construction of the set of models should be reconsidered (Burnham and Anderson 2002). For *v* values around 1 to 4, quasi-likelihood theory provides a way to analyse over dispersed data (Anderson et al. 1994; Richards 2008). The quasi-likelihood is the likelihood divided by an estimate of *v*. The quasi-likelihood version of the various information criteria, namely QAICc, $QKIC_c$ (Kim et al. 2014) and QBIC, is obtained simply by replacing the likelihood with the quasi-likelihood in the corresponding formula. In such cases, the number of parameters is increased by one and the model variance is multiplied by *v* (see below). When the quasi-likelihood version is used, it must be done for all models and criteria.

3.3. Model weights

Let IC be any information criterion. For a particular criterion and for any set of *R* models there is a minimum criterion value e.g. AIC_{cmin}, BIC_{min}, etc. Thus, the models can be ranked regarding the difference with that minimum

$$\Delta_i = IC_i - IC_{min} \text{ , for } i = 1, 2, \ldots, R$$

where IC_i refers to any specific information criterion for the model *i*.

Models can also be ranked by their weights from higher to lower. The weight w_i refers to the strength of evidence for that model (Burnham et al. 2011; Claeskens 2016)

525 $w_i = l_i / \Sigma l_j$ for $j = 1, 2, \ldots, R$

526 where $l_i = \exp(-0.5\Delta_i)$ is the relative likelihood of each model given the data.

527

528 *3.4 Multi-model inference*

529 Multi-model-based inference estimate the parameters of interest based on a group of the
530 most credible models instead of on a best-fit single model (Burnham and Anderson 2002;
531 Burnham et al. 2011; Symonds and Moussalli 2011). The multi-model inference is
532 performed as a model averaged prediction for the parameters that are variables in the best
533 model.

534 In our modelling framework and before performing the average of the estimated parameter
535 values, the different models should be translated to the same scale of mutual-propensity.
536 For example, a model like $m'_{11} = 2$, $m'_{12} = m'_{21} = m'_{22} = 1$, is not in the same scale that $m'_{11} =$
537 2, $m'_{12} = m'_{21} = m'_{22} = 0.5$. Without loss of generality, the latter can be transformed into an
538 equivalent model $m'_{11} = 4$, $m'_{12} = m'_{21} = m'_{22} = 1$, which is now in the same scale that the
539 first model.

540 The averaged parameter estimates were computed as a weighted mean where the weights
541 are the strength of evidence for each model as obtained under a given information criterion.
542 The parameters were averaged only over the models for which they appear as a variable.
543 Because the weights need to sum up to 1, it was necessary renormalize them by dividing by
544 the accumulated weight in the confidence subset.

Therefore, for each parameter *m* included in the confidence subset R_s, the average was computed as

$$\widehat{m} = \frac{\sum_i^{Rs} w_i \widehat{m}_i}{\sum_i^{Rs} w_i}$$

This way of performing the model averaged prediction is called natural averaging (Symonds and Moussalli 2011).

Finally, the reliability of each parameter estimate was measured as the unconditional standard error

$$Se(\widehat{m}) = \sum_i^{Rs} w_i \sqrt{vV(\widehat{m}_i) + (\widehat{m}_i - \widehat{m})^2}$$

where $V(\widehat{m}_i) = V(m_i \mid \text{model } i) = V(q') = q'(1-q') / n'$ is the model standard error squared and *v* is the variance inflation factor.

The use of the sum of weights to estimate variable importance in regression models has been criticized because of multicollinearity among the predictor variables and the imprecision of the weight measures (Galipaud et al. 2014; Cade 2015; Galipaud et al. 2017). However, the mutual-propensity parameters do not belong to a regression model and their average is performed in the same scale and with comparable units. Therefore, under the mutual mating propensity setting, the multimodel inference would work well as it was confirmed by Monte Carlo simulation (next section).

4. Simulations

4.1. Polygamous species (sampling with replacement)

565 To test how well the above methodology is able to distinguish among the different classes
566 of models and estimate the mating parameters, I used the sampling with replacement
567 algorithm in the program MateSim (Carvajal-Rodríguez 2018a) for generating mating
568 tables by Monte Carlo simulation (see Appendix B for detailed explanation).

569 The simulated cases correspond to one-sex competition and mate choice models. The
570 resulting mating tables were consequence of the mating system and the sampling process,
571 and consisted in two types of information (Fig. B1 in Appendix B). First, the population
572 frequencies (premating individuals) which were generated randomly for each simulation
573 run. Second, the sample of 500 mating pairs ($n' = 500$) for a hypothetical trait with two
574 classes at each sex. Because the simulated species had large population size ($n = 10\,000$)
575 the mating process was represented as a sampling with replacement, and the population
576 frequencies were constant over the mating season. The minimum phenotype frequency
577 (MPF) allowed was 0.1.

578 Five different model cases were simulated, namely random mating with mutual-
579 propensities $m'_{11}=m'_{22}=m'_{12}=m'_{21}= 1$ (M_0 in Table 2), female competition ($\alpha = 2$) and mate
580 choice ($c = 3$) with independent parameters $m'_{11}= m'_{12} = 2$, $m'_{22} = 3$, $m'_{21} = 1$ (SfC Table 2),
581 and with compound parameters $m'_{11}= 6$, $m'_{12} = 2$, $m'_{22} = 3$, $m'_{21} = 1$ (SfCc Table 2), and
582 male competition ($\alpha = 2$) and mate choice ($c = 3$) with independent parameters $m'_{11}= m'_{21} =$
583 2, $m'_{22} = 3$, $m'_{12} = 1$ (SmC Table 2), and with compound parameters $m'_{11}= 6$, $m'_{21} = 2$, m'_{22}
584 $= 3$, $m'_{12} = 1$ (SmCc Table 2). Each case was simulated 1 000 times.

585 For each simulation run, and given the normalized mutual-propensities m_{ij}, the number of
586 occurrences for each mating class $i \times j$ was obtained as

$$Q(i,j) = n' \times p_{1i} \times p_{2j} \times m_{ij}$$

where n' is the sample size, p_{1i} is the female population frequency for the phenotype i, p_{2j} is the male population frequency for the phenotype j.

Once the mating tables were obtained I proceeded with the multimodel inference analysis using InfoMating. Note that there were 1 000 different tables for each simulated case so, in the simulation study, it is better to consider the mean multimodel estimates instead of the full list of analysed models (which would imply 1 000 lists for each simulated case). Also, it is worth noting that with real data, the exactly true model is not necessarily included in the set of assayed models and so, it is important to evaluate the accuracy of the multimodel parameter estimates because, if the parameter estimates are correct, the model that would arise from that estimates and the set of most supported candidate models must be a good guess of the true one.

The sequence of analyses was as follows. For each mating table, InfoMating generates a set of 17 models, from the simplest random model M_0 to the saturated M_{sat}, including mate competition and choice models with one or two parameters (see all the types in Table 1). Then, the program computes the information criteria for each model and performs the multimodel inference as explained in the previous section. Thus, for each of the 5 simulated cases, 1 000 parameter estimates were obtained, and their average and standard error computed (Table 2).

Table 2. Average (standard error) parameter estimates under sample size 500 for a polygamous species with large population size (N = 10 000).

Model		m'_{11}	m'_{12}	m_{21}	m_{22}
M_0	**Expected**	**1**	**1**	**1**	**1**
	AICc	1.0 ± 0.0000	1.0 ± 0.0000	1.0 ± 0.0000	1.0 ± 0.0000
	KICc	1.0 ± 0.0000	1.0 ± 0.0000	1.0 ± 0.0000	1.0 ± 0.0000
	BIC	1.0 ± 0.0000	1.0 ± 0.0000	1.0 ± 0.0000	1.0 ± 0.0000
SfC	**Expected**	**2**	**2**	**1**	**3**
	AICc	1.94 ± 0.0017	1.99 ± 0.0009	1.0 ± 0.0000	2.95 ± 0.0025
	KICc	1.94 ± 0.0024	1.99±0.0020	1.0 ± 0.0000	2.95 ± 0.0029
	BIC	1.90 ± 0.0060	1.94 ± 0.006	1.0 ± 0.0000	2.90 ± 0.0074
SfCc	**Expected**	**6**	**2**	**1**	**3**
	AICc	5.93 ± 0.0044	2.0 ± 0.0001	1.0 ± 0.0000	2.97 ± 0.0027
	KICc	5.92 ± 0.0051	2.0 ± 0.0001	1.0 ± 0.0000	2.97 ± 0.0028
	BIC	5.87 ± 0.0086	2.0 ± 0.0017	1.0 ± 0.0000	2.96 ± 0.0044
SmC	**Expected**	**2**	**1**	**2**	**3**
	AICc	1.94 ± 0.0020	1.0 ± 0.0000	1.99 ± 0.0013	2.94 ± 0.0030
	KICc	1.93 ± 0.0032	1.0 ± 0.0000	1.98 ± 0.0029	2.93 ± 0.0037
	BIC	1.90 ± 0.0062	1.00 ± 0.0000	1.93 ± 0.0065	2.88 ± 0.0080
SmCc	**Expected**	**6**	**1**	**2**	**3**

AICc	5.93 ± 0.0046	1.0 ± 0.0000	2.0 ± 0.0001	2.97 ± 0.0029
KICc	5.92 ± 0.0052	1.0 ± 0.0000	2.0 ± 0.0001	2.97 ± 0.0029
BIC	5.87 ± 0.0085	1.0 ± 0.0000	2.0 ± 0.0010	2.97 ± 0.0037

M_0: Random mating model. SfC: female competition and mate choice with independent parameters. SfCc: female competition and mate choice with compound parameters. SmC: male competition and mate choice with independent parameters. SmCc: male competition and mate choice with compound parameters.

It can be appreciated that the random mating was perfectly estimated by the three IC methods. The competition plus mate choice parameter estimates were fairly good under the three criteria. The estimates were slightly better under AICc and slightly less accurate under BIC.

The whole simulation process was repeated using a small sample size (n' = 50 matings) and the results were qualitatively similar. However, the parameter estimates tended to be low-biased possibly because the power to detect deviations from random mating was low (see supplementary Table C1 in Appendix C).

4.2. Monogamous species (sampling without replacement)

For monogamous species, the mating process is without replacement (from the point of view of the available phenotypes) and can be represented via mass-encounters (Gimelfarb 1988; Carvajal-Rodríguez 2018a). The pattern obtained under the mass-encounter monogamous scenario (when the population size is large) was qualitatively similar to the polygamous species. However, there were less power to detect deviation from random

mating and so the estimates were low-biased, especially in the case of the compound parameter. Regarding sample size, it seems that the estimation was not very much affected (see supplementary Tables C2 and C3 in Appendix C).

Not surprisingly, the case of monogamous species with small population size (N = 200) was the worst scenario for multimodel estimation under the assumption of constant population phenotype frequencies (see Table C4 in Appendix C). Under this case and when most of the adults were involved in the mating process (mating sample size = 100), the change in the population phenotype frequencies during the breeding season significantly affected the observed non-random mating patterns. Only when the deviation from random mating is as large as with the compound effect of choice and competition, the estimated mutual-propensities provided some information (SfCc in Table C4).

5. Example of application

Littorina saxatilis is a marine gastropod mollusc adapted to different shore habitats in Galician rocky shores. There are two different ecotypes, an exposed-to-wave (smooth un-banded, SU), and a non-exposed (rough banded, RB) ecotype. Several experimental studies have shown that these ecotypes have evolved local adaptation at small spatial scale. For example, stronger waves on the lower shore may provoke that the SU ecotype becomes sexually mature at smaller size than the upper-shore (RB) ecotype. In addition, in some areas of the mid-shore habitat, the two ecotypes occasionally mate, producing apparently fertile intermediate morphological forms that are called hybrids (HY) (Rolan-Alvarez et al. 2015a).

649 Sexual isolation (positive assortative mating) between RB and SU morphs was observed in
650 wild mating pairs in the mid-shore zone, likewise within-morph size-assortative mating in
651 all shore levels (Cruz et al. 2001). It is assumed that the size is the key trait causing the
652 increase of sexual isolation in this model system, being the males the choosy sex in this
653 species (Rolan-Alvarez 2007).

654 Here, I reanalysed a *L. saxatilis* data set (Cruz et al. 2001) for estimating the mutual-
655 propensity parameters between the RB, SU and HY morphs in the mid-shore habitat. In the
656 original study, the authors analysed a hybrid zone encompassing 30 km of coast in Galicia
657 (NW Spain) with two sampling locations (Centinela and Senin) and seasons (autumn and
658 summer). Copulating pairs were collected jointly with the 15 nearest non-mating
659 individuals (see details in Cruz et al. 2001). In the present reanalysis, I considered the
660 pooled data of the two sampling locations and seasons (Table 3).

661

662 **Table 3.** The population frequencies by sex and the sample of matings from Cruz et al.
663 (2001) data.

	Total	**RB**	**HY**	**SU**
Female freqs	1254	0.22	0.11	0.67
Male freqs	1080	0.26	0.12	0.62
Matings			**Males**	
Females		**RB**	**HY**	**SU**
	RB	19	9	13
	HY	6	4	10
	SU	6	7	80

First, I computed the information partition (Carvajal-Rodríguez 2018b) that indicated significant assortative mating from the Chi-square test (J_{PSI} p-value < 0.0000001) while no significant sexual selection was detected. However, the randomization test was not significant in any case, possibly due to the low sample size within the mating classes.

Second, I proceeded with the model estimation and initially assayed only the subset of models with male and/or female mate competition plus the saturated (M_{sat}) and random mating (M_0) models. The estimate of overdispersion was high (7.20) indicating poor structure of the set of models regarding the data. The three information criteria gave similar output with the M_0 as the best fit model. The multimodel estimates of the mutual-propensities were just one in every case as expected from random mating. Because in the simulation study, the AICc criterion gave the best estimates I will rely on this criterion from now on.

The next step was to study only models with choice parameter plus the saturated (M_{sat}) and random mating (M_0) models. The overdispersion was 4.65 that still indicates somewhat poor model structure. The best fit model was a choice model with one parameter. The multimodel inference gave a clear pattern of positive assortative mating, that was higher for the RB × RB mating (m'_{RBRB} = 3), intermediate for HY × HY (m'_{HYHY} = 2.3) and slightly lower for SU × SU (m'_{SUSU} = 2).

Then, I considered jointly the previous competition and choice models and added new ones having separated competition and choice parameters. The overdispersion was 3.4 that is an acceptable value for multinomial models and can be corrected by using quasi-likelihoods

(see the overdispersion section above). Now, the best fit was a compound parameter model with female competition and choice. The estimates from this model were a RB female competition of $\alpha = 1.7$ and choice $c = 2.4$. The multi-model estimates gave positive asssortative mating, $m'_{RBRB} = 3$, $m'_{HYHY} = 3$, $m'_{SUSU} = 2$ and sexual selection favouring RB females.

Finally, I considered all the previous models plus models having parameters with double effect (i.e. one parameter may generate both sexual selection and assortative mating patterns). This implies a total of 35 models including M_0 and M_{sat}. The overdispersion was 2.5. The best model was the same for the three criteria and it was a double effect model with 2 parameters, $a = 0.2$ and $b = 2$, distributed as indicated in Fig. 5. Approximately, the same model was obtained using the multi-model estimates.

$$\begin{pmatrix} 2 & 2 & 1 \\ 1 & 2 & 1 \\ 0.2 & 1 & 1 \end{pmatrix}$$

Fig 5. Model D-2P-Rep3: Double two parameter model with three repetitions of the *b* parameter ($a = 0.2$, $b = 2$) producing female and male sexual selection plus positive assortative mating.

It is also possible to focus only on the models with separated parameters for competition and choice. The best fit model from this subgroup involves female competition. Recall that in *Littorina saxatilis* the choosy sex are the males, so I considered that the competitive advantage from the side of the females is explained by the males preferring a given kind of

females. The best fit model is $S_{Fem}C$-2Pc (see Table 1) with RB female competitive advantage of 1.7 more times matings than the other females and a choice parameter of 2.4. The qualitative pattern obtained from these models is similar to that in Fig.5; the RB females (first row) are preferred and there is a choice for within ecotype mating. The combination of competition and choice provokes that the mating RB×RB is the preferred by RB males; the matings RB×HY and HY×HY are preferred by HY males, and finally, it seems that the SU males do not discriminate between female ecotypes.

6. Discussion

6.1 Simulations

I have simulated mating tables corresponding to random mating, mate competition and mate choice models. The random mating pattern was perfectly assessed. For the other models, the competition and choice parameters were estimated quite accurately when the mating system resembles a sampling with replacement. Not surprisingly, BIC was slightly more conservative, while AICc presented slightly more accurate estimates in most cases. The KICc criterion performed similar to the best AICc and BIC cases. In general, the estimation was accurate and even in the cases with extreme phenotypic frequencies, the mean estimates were closer to the real value than to random mating.

The proposed approach does not require a particular experimental set-up other than the mating tables. However, to correctly identify the processes that produce the patterns of sexual selection and assortative mating, it is assumed that the encounters occur at random, i.e. the encounter between two phenotypes depends on the population phenotypic

distribution, and that the mating pattern is the product of the phenotypic distribution of the population and the individual preferences (Carvajal-Rodríguez 2018a). As a consequence, the availability of phenotypes should not be affected by the matings that have already occurred, as expected for polygamous species, or even for monogamous species, when the number of available individuals is higher than the mating pairs.

However, the above assumption is likely to be violated in the case of monogamous species with low population size, or even in large population sizes with local competition for mates (if the number of individuals in the patches is low) and/or space-temporal constraints. In such cases, the population phenotype frequencies may be altered during the reproductive season and the sexual selection and assortative mating patterns would be more difficult to detect (Carvajal-Rodríguez 2019). In fact, the simulations (see Appendix C) showed that the performance of the multimodel inference is affected by the sampling and the mating system (polygamous or monogamous) but it is still quite robust for detecting non-random mating deviation in the parameter values except in the worst scenario of monogamous species with small population size.

6.2 General

The advantages of model selection and multimodel inference in evolutionary ecology has been widely discussed, jointly with the pros and cons of applying any information criteria (Link and Barker 2006; Burnham et al. 2011; Aho et al. 2014; Barker and Link 2015; Aho et al. 2017; Dziak et al. 2019) or the reliability of the obtained estimates (Galipaud et al. 2014; Cade 2015; Giam and Olden 2016; Galipaud et al. 2017).

748 Multimodel inference has been however, rarely utilized to study the mating patterns that
749 may emerge from mate choice and mate competition. Here, by developing general models
750 that incorporate competition and mate choice, and providing their maximum likelihood
751 estimates, I am proposing a standardized methodology for model selection and multimodel
752 inference of the mating parameters producing the sexual selection and assortative mating
753 patterns.

754 The set of *a priori* models permits to perform an *a posteriori* quantification of the data-
755 based evidence and provide confidence sets on plausible non-trivial models while letting
756 multi-model inference of the parameter values. The approach was implemented by allowing
757 three different information criteria. Under the scenarios assayed, they performed similarly
758 for simulated and real data.

759 Regarding the methodology, it is worth noting that although the mating tables require at
760 least two phenotypes by sex (2×2 dimensions or higher) for fitting mate competition and
761 mate choice models, the proposed approach can still be applied if some sex, say females,
762 have only one phenotypic class. In this case, we just need to duplicate the row (see Fig. D1
763 in the Appendix D). Obviously, there cannot be any assortative pattern and sexual selection
764 can only be measured in the sex with more than one phenotypic class.

765 The statistical tools developed in this work have been also applied to empirical data.
766 Previous studies in the Galician *L. saxatilis* hybrid zone showed that mate choice favours
767 within-morph pairs (reviewed in Rolan-Alvarez 2007). The estimates obtained by
768 multimodel inference support the positive assortative mating for the ecotype. In addition,
769 another result emerged from the analysis: The RB females are preferred in general i.e. RB

male with SU female has less mutual-propensity than SU male with RB female ($m_{SURB} < m_{RBSU}$). This pattern may be favoured by the physical difficulty for the mating involving bigger RB males with the smaller SU females, and could be related with the somehow more frequent occurrence of mating pairs having females bigger than males (a typical trend in gastropods, E. Rolán-Alvarez personal communication). Besides the mating pattern depicted by the multi-model approach, the estimates of the mutual-propensities were also obtained. Testing the reliability of these estimates is, however, out of the scope of the present manuscript, and it was left for future work.

To conclude, I present a methodology to distinguish among several models of mate competition and choice behind the observed pattern of mating and the phenotypic frequencies in the population. From an empirical point of view it is much easier to study patterns than processes and this is why the causal mechanisms of natural and sexual selection are not so well known as the patterns they provoke. The goal of the present work is to propose a new tool that will help to distinguish among different alternative processes behind the observed mating pattern.

The developed methodology has been fully implemented in a program called InfoMating available at http://acraaj.webs6.uvigo.es/InfoMating/Infomating.htm or upon request to the author.

Acknowledgement

I would like to thank Carlos Canchaya, Emilio Rolán-Alvarez, Sara Magalhaes, Alexandre Courtiol and two anonymous reviewers for many valuable comments that significantly

improved this paper. E. R-A also provided the data for the example. This work was supported by Xunta de Galicia (Grupo de Referencia Competitiva, ED431C2016-037), Ministerio de Economía y Competitividad (CGL2016-75482-P) and by Fondos FEDER ("Unha maneira de facer Europa").

Appendix A) Mutual Mating Propensity Models

Saturated non-random mating model: λ notation.

Consider the total number of possible mating phenotypes $K = k_1 \times k_2$ and the saturated multinomial model for the K-1 free mating parameters m'_{ij}.

The log-likelihood function is

$$lnL_{sat} = C + \sum_{ij}^{K-1} n'_{ij} \ln\left(m'_{ij} q_{ij}\right) + n'_{k1k2} \ln\left(a\left(1 - \sum_{ij}^{K-1} q_{ij}\right)\right) - n' \ln(M)$$

where n' is the number of matings in the sample and n'_{ij} is the number of matings between i-type females and j-type males. I have fixed the parameter m'_{k1k2} to a.

Compute the first derivative of the likelihood with respect to a

$$\frac{dlnL_{sat}}{da} = \frac{n'_{k1k2}}{a} - \frac{n}{M} q_{k1k2}$$

then by taking $a = 1$ and equating to 0 we get

808 $n/M = n'_{k1k2} / q_{k1k2} \equiv \lambda(1)$

809 that corresponds to the number of observed matings having unity mating parameter divided
810 by the corresponding product of population frequencies. Under the saturated model there is
811 only one (for convenience m'_{k1k2}) mating parameter having unitary value and so the number
812 of observed matings is n'_{k1k2} and the product of the corresponding population frequencies is
813 $p_{1k1} \times p_{2k2} = q_{k1k2}$.

814 Now, let find the m'_{ij} parameter value that maximizes the likelihood

815 $$\frac{dlnL_{sat}}{dm'_{ij}} = \frac{n'_{ij}}{m'_{ij}} - \frac{n}{M} q_{ij} = 0$$

816 $$\widehat{m'_{ij}} = \frac{n'_{ij}/q_{ij}}{n/M} \equiv \frac{\lambda(m'_{ij})}{\lambda(1)}$$

817 The λ notation can be generalized for any set A of mating phenotypes having the same
818 value of propensity θ as follows

820 $$\lambda(\theta) = \frac{\sum_{ij}^{A} n'_{ij}}{\sum_{ij}^{A} q_{ij}}$$

819

821 where x_{ij} represents the number of mating pairs having absolute (non-normalized) mating
822 parameter θ and q_{ij} is the product of the population frequencies p_{1i} and p_{2j} i.e. the expected
823 frequency of the θ mating phenotypes under random mating.

824

825 *Intrafemale competition models*

The model is

$m'_{ij} = a_i$ with $a_i > 0 \; \forall \; i < k_1$ and $m'_{k1j} = a = 1 \; \forall \; j$.

There are k_1-1 independent parameters. Note that the parameters m'_{k1j} have been fixed to a = 1. The log-likelihood function is

$$lnL = C + \sum_{i<k1}^{k1-1} \sum_{j=1}^{k2} n'_{ij} \ln(a_i q_{ij}) + \sum_{j=1}^{k2} n'_{k1j} \ln(a q_{k1j}) - n ln(M)$$

Now, assume that the parameter a is not fixed and compute the first derivative of the likelihood with respect to a

$$\frac{dlnL}{da} = \frac{\sum_j^{k2} n'_{k1j}}{a} - \frac{n}{M} \sum_j^{k2} q_{k1j} \; = 0$$

then by taking a = 1 and equating to 0 we get

$n/M = \sum_j n'_{k1j} / \sum_j q_{k1j} \equiv \lambda(1)$

Now find the a_i parameter value that maximizes the likelihood

$$\frac{dlnL}{da_i} = \frac{\sum_j^{k2} n'_{ij}}{a_i} - \frac{n}{M} \sum_j^{k2} q_{ij} \; = 0$$

Solving for a_i

$$\widehat{a_i} = \frac{\sum_j^{k2} n'_{ij} \, / \sum_j^{k2} q_{ij}}{\frac{n}{M}} \equiv \frac{\lambda(a_i)}{\lambda(1)}$$

The formula expressed as the quotient of lambdas is valid for any number h of different parameters, $1 \leq h < k_1$. In the particular case of having only one parameter the sum of observed matings having propensity a_1, implies $\Sigma\Sigma n'_{ij}$ where the first summation is for all the female types except females of type k_1, and the second is over all male types. The sum of the product of frequencies is $1 - p_{1k1}$.

As before, $\lambda(1)$ also corresponds to the sum of the observed matings having expected propensity 1 divided by the sum of the corresponding products of population frequencies. The model for male sexual selection is solved in a similar way.

Intrasexual competition in both sexes

The model is

$$m'_{ij} = a_i b_j,\ i < k_1,\ j < k_2;\ m'_{ik2} = a_i,\ i < k_1;\ m'_{k1j} = b_j,\ j < k_2;\ m'_{k1k2} = 1$$

with $a_i > 0$, $b_j > 0\ \forall\ i, j$.

It is easy to see that is multiplicative. Let $A = \sum_i^{k1-1} a_i p_{1i} + a p_{1k1}$ and $B = \sum_j^{k2-1} b_j p_{2j} + b p_{2k2}$.

The mean mutual mating propensity is

$$M = \sum_{i,j} q_{ij} m'_{ij} = \sum_{\mathrm{i}}^{\mathrm{k1}} \sum_{\mathrm{j}}^{\mathrm{k2}} a_i b_j p_{1\mathrm{i}} p_{2\mathrm{j}} = AB$$

with $a_{k1} = a$ and $b_{k2} = b$.

The marginal propensity for i-type females is

$$m_{Fem_i} = a_{\mathrm{i}} \sum_j^{males} p_{2j} \frac{\mathrm{b_j}}{M} = a_{\mathrm{i}} \frac{B}{M}$$

Similarly, the marginal for *j*-type males

$$m_{Male_j} = b_{\mathrm{j}} \frac{\mathrm{A}}{M}$$

with $a_{k1} = a$ and $b_{k2} = b$.

The condition 5-iii) for a multiplicative model implies that $m_{ij} = M_{Fem_i} \times M_{Male_j}$. In addition, $m_{ij} = a_i b_j / M$ that jointly with the multiplicative condition requires $a_i b_j / M = M_{Fem_i} \times M_{Male_j} = a_i B b_j A / M^2$ solving for M we get $M = AB$ which we have already seen it is true.

The log-likelihood function

$$lnL = C + \sum_{i=1}^{k1} \sum_{j=1}^{k2} n'_{ij} \ln\left(a_i b_j q_{ij}\right) - nln(M)$$

with $a_{k1} = a = 1$ and $b_{k2} = b = 1$.

Consider the derivatives

$$\frac{dA}{da_i} = p_{1i} \; ; \frac{dB}{db_j} = p_{2j}; \frac{dM}{da_i} = p_{1i} B; \frac{dM}{db_j} = p_{2j} A$$

Now by taking the derivative of the log-likelihood with respect to a_i or b_j and equating to 0 we get the estimates

$$\widehat{a_\iota} = \frac{\sum_j^{k2} n'_{ij} / \sum_j^{k2} q_{ij}}{\frac{n}{M} B} = \frac{\lambda(a_i + \sum_j a_i b_j)}{\lambda_{fem}(1)}$$

874 $$\widehat{b_j} = \frac{\sum_i^{k1} n'_{ij} / \sum_i^{ki} q_{ij}}{\frac{n}{M}A} = \frac{\lambda(b_j + \sum_i a_i b_j)}{\lambda_{male}(1)}$$

875 Where

876 $$\frac{n}{M}B = \frac{\sum_j^{k2} n'_{k1j}}{p_{1k1}} \equiv \lambda_{fem}(1)$$

877 $$\frac{n}{M}A = \frac{\sum_i^{k1} n'_{ik2}}{p_{2k2}} \equiv \lambda_{male}(1)$$

878

879 *Mate choice models with parameterized heterotypes*

880 Consider models in which the homotype mating has absolute propensity of 1 while the
881 different heterotypes have absolute value of a_{ij}. The maximum likelihood estimate is

882 $$\hat{a}_{h1h2} = \frac{\lambda(a_{h1h2})}{\lambda(1)}$$

883 The number of parameters in this type of model is $K - min\{k_1, k_2\} - \Sigma_S(C_s\text{-}1)$ where the
884 sum is over the set of different heterotype matings and C_s is the cardinality of each set.

885

886 *Double effect models*

887 The following models generate a double pattern of sexual selection and assortative mating
888 even when the population frequencies are uniform.

889

Double effect models producing sexual selection in one sex under uniform frequencies

A simple approach consists in building a new model by setting $m'_{ii} = 1$ and $m'_{jj} = 1 + c$. Then, if we desire assortative mating jointly with sexual selection only in females we additionally set $m'_{ij} = 1 - c$; on the contrary, if we desire selection only in males we set $m'_{ji} = 1 - c$ with $-1 < c < 1$. If the frequencies are not uniform the model generates assortative mating jointly with sexual selection in both sexes.

In the case of the model with $m'_{ij} = 1 - c$ (female sexual selection if frequencies are uniform) the maximum likelihood estimate of c is one of the roots of the quadratic

$$(x_{jj} - x_{ij} + n'D) - c[x_{ij} + x_{jj} + D(x_{jj} - x_{ij})] - c^2 D[n' - (x_{ij} + x_{jj})] = 0$$

where $D = q_{ij} - q_{jj}$ and $n' = \sum x_{ij}$ is the number of matings (sample size).

If the frequencies are uniform and $k_1 = k_2$, i.e. $p_{1i} = p_{1j} = p_{2i} = p_{2j}\ \forall\ i, j$ then

$$\hat{c} = \frac{x_{jj} - x_{ij}}{x_{jj} + x_{ij}}$$

The case for male sexual selection is obtained simply by interchanging x_{ij} by x_{ji} and q_{ij} by q_{ji} in the formulas.

The above model has only one parameter c; we can introduce a more complex two parameter model, $M_{(a,c)}$ by setting $m'_{ii} = a$, $m'_{jj} = 1 + c$ and $m'_{ij} = 1 - c$, for female sexual selection (or $m'_{ji} = 1 - c$ for male sexual selection). For obtaining the MLE of this two parameter double model, with restrictions $a > 0$, $c < |1|$, I have used a numerical bounded Nelder-Mead simplex algorithm (Press 2002; Singer and Singer 2004; Gao and Han 2012).

Double effect models with sexual selection in both sexes under uniform frequencies

To get assortative mating jointly with sexual selection in both sexes under uniform frequencies, we just need to combine the above uniform one parameter models of each sex, so that $m'_{ii} = 1$, $m'_{jj} = 1 + c$ and $m'_{ij} = m'_{ji} = 1 - c$.

The maximum likelihood estimate of c involves the solution of the quadratic

$$[x_{jj} - x_s + n'D_2] - c[x_{jj} + x_s + D_2(x_{jj} - x_s)] + c^2 D_2[x_{jj} + x_s - n'] = 0$$

where $x_s = x_{ij} + x_{ji}$ and $D_2 = q_{ij} + q_{ji} - q_{jj}$.

General double effect models

We can also define a set of general models where any propensity m'_{ij} has parameter θ_{ij} with at least one propensity having value of 1. The MLE of the parameters of this kind of model is

$$\hat{\theta}_{ij} = \frac{\lambda(\theta_{ij})}{\lambda(1)}$$

where $\lambda(\theta_{ij})$ is defined as in (A2).

The simplest model defined in this way is

$$m'_{ii} = a \text{ and } m'_{ij} = m'_{ji} = m'_{jj} = 1,$$

which produces assortative mating and sexual selection in both sexes.

Consider as an example of this model, the case with $k_1 = k_2 = 2$ so that $0 < p_{11} < 1$; $0 < p_{21} < 1$; $m'_{11} = a$ and $m'_{12} = m'_{21} = m'_{22} = 1$. The mean mating propensity is $M = q_{11}(a - 1) + 1$. The absolute marginal propensity for the first female type $m'_{Fem_1} = a\, p_{21} + 1 - p_{21} = p_{21}(a - 1) + 1$, and for the second female type $m'_{Fem_2} = 1$. Similarly the male marginals are $m'_{Male_1} = p_{11}(a - 1) + 1$ and $m'_{Male_2} = 1$.

Recall that the condition for the sexual selection pattern within a given sex is that the marginal mating propensities are different which here is true for both sexes provided that $a \neq 1$. Regarding the assortative mating pattern it can be proved that the joint isolation index (I_{PSI}) is 0 only if $a = 1$. However, it is sufficient to prove that the model is not multiplicative (Carvajal-Rodríguez 2018b). Consider that the model is multiplicative, this implies, $m'_{12}/M = (m'_{Fem_1}/M) \times (m'_{Male_2}/M)$ that given the model values becomes

$$M = (m'_{Fem_1}) \times (m'_{Male_2})$$

which can be true only when $p_{11} = 1$ and so it is false by definition.

The estimate of a under this model is $\lambda(a) / \lambda(1)$.

The most parameterized model that can be defined in this way has K-1 free parameters and coincides with the saturated model so that the estimates are the corresponding pair total indices (PTI_{ij}).

Moreover, note that if no mutual propensity is fixed to 1 then $\lambda(1) = (n - A) / (1 - P) = n$ where A = number of observations having value 1 = 0 and P = product of population frequencies of the involved types having mutual propensity 1 = 0. Therefore the estimate of θ_{ij} can also expressed as $\lambda(\theta_{ij}) / n$ which is the observed frequency of mating pairs (i, j)

948 divided by the expected frequency by random mating which is the definition of the pair
949 total index PTI_{ij} (K-1 are free and one PTI is dependent on the others).

950 All the above derived MLE formulae have been checked by a numerical bounded Nelder-
951 Mead simplex algorithm (Press 2002; Singer and Singer 2004; Gao and Han 2012).

952

953 **Appendix B) Monte Carlo simulation of mating tables**

954 The mating tables for the simulation experiments were generated by the program MateSim
955 (Carvajal-Rodríguez 2018a) available at http://acraaj.webs.uvigo.es/MateSim/matesim.htm.

956 The number of replicates for each case was 1 000. For each run the program first generated
957 the number of premating males and females from a given population size. For example, if
958 the population size consisted in n_1 (= 5 000) females and n_2 (= 5 000) males, the program
959 got $n_{1A} = n_1 \times U$ females of the A type and $n_{1B} = n_1 - n_{1A}$ females of the B type. Where U is
960 a value sampled from the standard uniform distribution. The premating males were
961 obtained similarly. Then, the female population frequencies were $p_{1i} = n_{1i} / n_1$, and $p_{2i} = n_{2i}$
962 $/ n_2$ for the male ones. Finally, a sample of n' (= 500) matings was obtained, where the
963 number of counts for each mating phenotype $i \times j$ was

964 $$Q(i,j) = n' \times p_{1i} \times p_{2j} \times m'_{ij} \ / M$$

965 where m'_{ij} are the mutual-propensity parameters as defined for each kind of model, and $M =$
966 $\sum p_{1i} \times p_{2j} \times m'_{ij}$.

The format of the obtained tables was the same as the JMating (Carvajal-Rodriguez and Rolan-Alvarez 2006) input files (Fig. B1).

```
# format number
0
# num of types
2
#premating male numbers
1626        3374
#premating female numbers
917     4083
# matings by rows (females)
30      62
133     275
```

Fig. B1. Example of a table generated by the simulations. The format is the same as for the JMating software.

Appendix C) Polygamous species with low sample size and monogamous species

Table C1. Average (standard error) parameter estimates under sample size 50 for a polygamous species with large population size (N = 10 000).

Model		m'_{11}	m'_{12}	m_{21}	m_{22}
M_0	**Expected**	**1**	**1**	**1**	**1**
	AICc	1.0 ± 0.0000	1.0 ± 0.0000	1.0 ± 0.0000	1.0 ± 0.0000
	KICc	1.0 ± 0.0000	1.0 ± 0.0000	1.0 ± 0.0000	1.0 ± 0.0000

	BIC	1.0 ± 0.0000	1.0 ± 0.0000	1.0 ± 0.0000	1.0 ± 0.0000
SfC	**Expected**	**2**	**2**	**1**	**3**
	AICc	1.38 ± 0.0106	1.45 ± 0.0119	1.0 ± 0.0003	2.03 ± 0.0206
	KICc	1.32 ± 0.0104	1.35±0.0110	1.0 ± 0.0003	1.76 ± 0.0226
	BIC	1.29 ± 0.0106	1.29 ± 0.0105	1.0 ± 0.0003	1.64 ± 0.0225
SfCc	**Expected**	**6**	**2**	**1**	**3**
	AICc	4.71 ± 0.0204	1.86 ± 0.0201	1.0 ± 0.0000	2.36 ± 0.0286
	KICc	4.45 ± 0.0236	1.57 ± 0.0216	1.0 ± 0.0000	2.06 ± 0.0316
	BIC	4.28 ± 0.0283	1.42 ± 0.0198	1.0 ± 0.0000	1.99 ± 0.0324

M_0: Random mating model. SfC: female competition and mate choice with independent parameters. SfCc: female competition and mate choice with compound parameters.

Table C2. Average (standard error) parameter estimates under sample size 500 for a monogamous species (mass-encounter mating process) with large population size (*N* = 10 000).

Model		m'_{11}	m'_{12}	m_{21}	m_{22}
M_0	**Expected**	**1**	**1**	**1**	**1**
	AICc	1.07 ± 0.0053	1.07 ± 0.0053	1.07 ± 0.0059	1.07 ± 0.0056
	KICc	1.05 ± 0.0044	1.04 ± 0.0043	1.05 ± 0.0052	1.04 ± 0.0050

	BIC	1.01 ± 0.0028	1.01 ± 0.0020	1.01 ± 0.0033	1.01 ± 0.0026
SfC	**Expected**	**2**	**2**	**1**	**3**
	AICc	1.95 ± 0.0158	1.95 ± 0.0160	1.0 ± 0.0001	2.04 ± 0.0176
	KICc	1.93 ± 0.0162	1.93±0.0166	1.0 ± 0.0001	2.03 ± 0.0180
	BIC	1.82 ± 0.0183	1.82 ± 0.0189	1.0 ± 0.0000	1.93 ± 0.0195
SfCc	**Expected**	**6**	**2**	**1**	**3**
	AICc	3.02 ± 0.0341	2.07 ± 0.0210	1.0 ± 0.0001	2.84 ± 0.0298
	KICc	3.0 ± 0.0343	2.04 ± 0.0216	1.0 ± 0.0000	2.82 ± 0.0302
	BIC	2.86 ± 0.0353	1.92 ± 0.0244	1.0 ± 0.0000	2.73 ± 0.0313

M_0: Random mating model. SfC: female competition and mate choice with independent parameters. SfCc: female competition and mate choice with compound parameters.

Table C3. Average (standard error) parameter estimates under sample size 50 for a monogamous species (mass-encounter mating process) with large population size (N = 10 000).

Model		m'_{11}	m'_{12}	m_{21}	m_{22}
M_0	**Expected**	**1**	**1**	**1**	**1**
	AICc	1.0 ± 0.0362	1.04 ± 0.0414	1.0 ± 0.0365	1.02 ± 0.0406
	KICc	1.02 ± 0.0339	1.03 ± 0.0374	1.0 ± 0.0324	1.02 ± 0.0377

	BIC	1.03 ± 0.0312	1.03 ± 0.0340	1.0 ± 0.0283	1.03 ± 0.0361
SfC	**Expected**	**2**	**2**	**1**	**3**
	AICc	2.0 ± 0.0568	2.05 ± 0.0638	1.08 ± 0.0155	2.23 ± 0.0635
	KICc	1.76 ± 0.0539	1.78±0.0605	1.06 ± 0.0145	1.96 ± 0.0597
	BIC	1.62 ± 0.0515	1.66 ± 0.0578	1.05 ± 0.0143	1.79 ± 0.0555
SfCc	**Expected**	**6**	**2**	**1**	**3**
	AICc	3.36 ± 0.1004	2.19 ± 0.0731	1.03 ± 0.0073	3.11 ± 0.0997
	KICc	3.06 ± 0.0951	1.94 ± 0.0715	1.03 ± 0.0066	2.85 ± 0.0964
	BIC	2.89 ± 0.0919	1.79 ± 0.0687	1.02 ± 0.0048	2.68 ± 0.0929

M_0: Random mating model. SfC: female competition and mate choice with independent parameters. SfCc: female competition and mate choice with compound parameters.

Table C4. Average (standard error) parameter estimates under sample size 100 for a monogamous species (mass-encounter mating process) with small population size (N = 200).

Model		m'_{11}	m'_{12}	m_{21}	m_{22}
M_0	**Expected**	**1**	**1**	**1**	**1**
	AICc	1.07 ± 0.0083	1.10 ± 0.0141	1.09 ± 0.0141	1.07 ± 0.0090
	KICc	1.04 ± 0.0064	1.05 ± 0.0094	1.05 ± 0.010	1.04 ± 0.0070

	BIC	1.02 ± 0.0047	1.02 ± 0.0056	1.02 ± 0.0065	1.02 ± 0.0052
SfC	**Expected**	**2**	**2**	**1**	**3**
	AICc	1.15 ± 0.0173	1.11 ± 0.0157	1.07 ± 0.0118	1.11 ± 0.0145
	KICc	1.10 ± 0.0141	1.06±0.0125	1.04 ± 0.0091	1.06 ± 0.0108
	BIC	1.03 ± 0.0072	1.03 ± 0.0079	1.02 ± 0.0059	1.03 ± 0.0060
SfCc	**Expected**	**6**	**2**	**1**	**3**
	AICc	1.85 ± 0.0555	1.30 ± 0.0317	1.07 ± 0.0124	2.12 ± 0.0757
	KICc	1.66 ± 0.0515	1.22 ± 0.0289	1.04 ± 0.010	1.94 ± 0.070
	BIC	1.49 ± 0.0476	1.13 ± 0.0248	1.02 ± 0.0039	1.71 ± 0.0637

M_0: Random mating model. SfC: female competition and mate choice with independent parameters. SfCc: female competition and mate choice with compound parameters.

Appendix D) Incomplete set-up: toy example

The proposed modelling framework requires at least two phenotypes by sex (mating tables of 2×2 dimensions or higher) for measuring sexual competition and mate choice effects. However it still can be applied if some sex, say females, have only one phenotype. In this case we just need to duplicate the row (see Fig. D1). Obviously, only male sexual selection can be measured.

A

```
# format number
0
# num of types
2
#premating male numbers
277 133
#premating female numbers
269 269
# matings by rows (females)
70 39
70 39
```

B

```
# format number
0
# num of types
2
#premating male numbers
277 133
#premating female numbers
269 269
# matings by rows (females)
100 9
100 9
```

Fig. D1. Examples of two toy models with only one type of female and two types of males. Note that the rows of the mating table are duplicated (same female type). A: Random mating B: Male sexual selection.

The examples in Fig. D1 correspond to a population with only one female but two male phenotypes (phenotype-1 and phenotype-2). There were sampled 269 females plus 277 males with phenotype-1 and 133 males with phenotype-2. In the first example (Fig. D1-A) there were 70 matings involving the male phenotype-1 and 39 with male phenotype-2. In the second example (Fig. D1-B) the matings were 100 with phenotype-1 and 9 with phenotype-2.

The analysis of the first case indicated that there was no significant deviation from random mating ($J_{PTI} = 0.005$, $P = 0.78$). The best model was the random mating model M_0. As expected, the multimodel estimation of the mutual mating parameters was 1 for every

parameter. The results were the same for the three information indices (AICc, KICc and BIC).

The analysis of the second case detected a deviation from random mating (J_{PTI} = 0.405, $P < 10^{-7}$) due to male sexual selection (J_{PS2} = 0.405, $P < 10^{-7}$) see (Carvajal-Rodríguez 2018b) for details of the *J* indices. The best model was male sexual selection with one parameter (Smale-1P). The male sexual selection component indicated five times higher mating propensity of male phenotype-1 with respect to phenotype-2.

References

Aho, K., D. Derryberry, and T. Peterson. 2014. Model selection for ecologists: the worldviews of AIC and BIC. Ecology 95:631-636.

Aho, K., D. Derryberry, and T. Peterson. 2017. A graphical framework for model selection criteria and significance tests: refutation, confirmation and ecology. Methods in Ecology and Evolution 8:47-56.

Akaike, H. 1973. Information theory and an extension of the maximum likelihood principle. Pp. 267-281 *in* B. N. Petrov, and F. Csaki, eds. Second International Symposium on Information Theory, Budapest: Akademiai Kiado.

Anderson, D. R., K. P. Burnham, and G. C. White. 1994. AIC Model Selection in Overdispersed Capture-Recapture Data. Ecology 75:1780-1793.

Andersson, M. 1994. Sexual selection. Princeton University Press, Princeton, N.J.

Armstrong, D. M. 1977. Dispersal vs. Dispersion: Process vs. Pattern. Systematic Biology 26:210-211.

Arnold, S. J. and M. J. Wade. 1984. On the measurement of natural and sexual selection: applications. Evolution:720-734.

Barker, R. J. and W. A. Link. 2015. Truth, models, model sets, AIC, and multimodel inference: A Bayesian perspective. The Journal of Wildlife Management 79:730-738.

Brewer, M. J., A. Butler, and S. L. Cooksley. 2016. The relative performance of AIC, AICC and BIC in the presence of unobserved heterogeneity. Methods in Ecology and Evolution 7:679-692.

Burnham, K. P. and D. R. Anderson. 2002. Model selection and multimodel inference: a practical information-theoretic approach. Springer-Verlag, New York, NY.

Burnham, K. P., D. R. Anderson, and K. P. Huyvaert. 2011. AIC model selection and multimodel inference in behavioral ecology: some background, observations, and comparisons. Behavioral Ecology and Sociobiology 65:23-35.

Cade, B. S. 2015. Model averaging and muddled multimodel inferences. Ecology 96:2370-2382.
Carvajal-Rodríguez, A. 2018a. MateSim: Monte Carlo simulation for the generation of mating tables. Biosystems 171:26-30.
Carvajal-Rodríguez, A. 2018b. Non-random mating and information theory. Theoretical Population Biology 120:103-113.
Carvajal-Rodríguez, A. 2019. A generalization of the informational view of non-random mating: Models with variable population frequencies. Theoretical Population Biology 125:67-74.
Carvajal-Rodriguez, A. and E. Rolan-Alvarez. 2006. JMATING: a software for the analysis of sexual selection and sexual isolation effects from mating frequency data. BMC Evol Biol 6:40.
Casares, P., M. C. Carracedo, B. del Rio, R. Piñeiro, L. Garcia-Florez, and A. R. Barros. 1998. Disentangling the Effects of Mating Propensity and Mating Choice in Drosophila. Evolution 52:126-133.
Cavanaugh, J. E. 2004. Criteria for linear model selection based on Kullback's symmetric divergence. Australian & New Zealand Journal of Statistics 46:257-274.
Chapleau, F., P. H. Johansen, and M. Williamson. 1988. The distinction between pattern and process in evolutionary biology: the use and abuse of the term'strategy'. Oikos 53:136-138.
Claeskens, G. 2016. Statistical Model Choice. Annual Review of Statistics and Its Application 3:233-256.
Cruz, R., E. Rolán-Álvarez, and C. García. 2001. Sexual selection on phenotypic traits in a hybrid zone of Littorina saxatilis (Olivi). Journal of Evolutionary Biology 14:773-785.
Darwin, C. 1871. The descent of man, and selection in relation to sex. Murray.
Darwin, C. 1974. The Descent of Man, and Selection in Relation to Sex, London.
Dziak, J. J., D. L. Coffman, S. T. Lanza, R. Li, and L. S. Jermiin. 2019. Sensitivity and specificity of information criteria. bioRxiv:449751.
Edward, D. A. 2015. The description of mate choice. Behavioral Ecology 26:301-310.
Endler, J. A. 1986. Natural selection in the wild. Princeton University Press, Princeton, N.J.
Estévez, D., T. P. T. Ng, M. Fernández-Meirama, J. M. Voois, A. Carvajal-Rodríguez, G. A. Williams, J. Galindo, and E. Rolán-Alvarez. 2018. A novel method to estimate the spatial scale of mate choice in the wild. Behavioral Ecology and Sociobiology 72:195.
Fitze, P. S. and J.-F. L. Galliard. 2011. Inconsistency between Different Measures of Sexual Selection. The American Naturalist 178:256-268.
Futuyma, D. J. and M. Kirkpatrick. 2017. Evolution. Sunderland, Massachusetts U.S.A : Sinauer Associates, Inc. Publishers.
Galipaud, M., M. A. F. Gillingham, M. David, F. X. Dechaume-Moncharmont, and R. B. O'Hara. 2014. Ecologists overestimate the importance of predictor variables in model averaging: a plea for cautious interpretations. Methods in Ecology and Evolution 5:983-991.
Galipaud, M., M. A. F. Gillingham, and F.-X. Dechaume-Moncharmont. 2017. A farewell to the sum of Akaike weights: The benefits of alternative metrics for variable importance estimations in model selection. Methods in Ecology and Evolution 8:1668-1678.

Gao, F. and L. Han. 2012. Implementing the Nelder-Mead simplex algorithm with adaptive parameters. Computational Optimization and Applications 51:259-277.
Gavrilets, S. 2004. Fitness landscapes and the origin of species. Princeton University Press, Princeton, N.J.
Giam, X. and J. D. Olden. 2016. Quantifying variable importance in a multimodel inference framework. Methods in Ecology and Evolution 7:388-397.
Gimelfarb, A. 1988. Processes of Pair Formation Leading to Assortative Mating in Biological Populations: Encounter-Mating Model. The American Naturalist 131:865-884.
Grueber, C. E., S. Nakagawa, R. J. Laws, and I. G. Jamieson. 2011. Multimodel inference in ecology and evolution: challenges and solutions. Journal of Evolutionary Biology 24:699-711.
Hartl, D. L. and A. G. Clark. 1997. Principles of Population Genetics. Sinauer Associates, Inc., Sunderland, MA.
Jennions, M. D. and M. Petrie. 1997. Variation in mate choice and mating preferences: a review of causes and consequences. Biological Reviews 72:283.
Keerativibool, W. 2014. Unifying the Derivations of Kullback Information Criterion and Corrected Versions. Thailand Statistician 12:37-53.
Kim, H.-J., J. E. Cavanaugh, T. A. Dallas, and S. A. Foré. 2014. Model selection criteria for overdispersed data and their application to the characterization of a host-parasite relationship. Environmental and ecological statistics 21:329-350.
Kokko, H., H. Klug, and M. D. Jennions. 2012. Unifying cornerstones of sexual selection: operational sex ratio, Bateman gradient and the scope for competitive investment. Ecology Letters 15:1340-1351.
Lewontin, R., D. Kirk, and J. Crow. 1968. Selective mating, assortative mating, and inbreeding: definitions and implications. Eugen Q 15:141-143.
Link, W. A. and R. J. Barker. 2006. Model weights and the fundations of multimodel inference. Ecology 87:2626-2635.
Liu, W. and Y. Yang. 2011. Parametric or nonparametric? A parametricness index for model selection. The Annals of Statistics 39:2074-2102.
Mahler, D. L., M. G. Weber, C. E. Wagner, and T. Ingram. 2017. Pattern and Process in the Comparative Study of Convergent Evolution. The American Naturalist 190:S13-S28.
Ng, T. P. T., E. Rolán-Alvarez, S. S. Dahlén, M. S. Davies, D. Estévez, R. Stafford, and G. A. Williams. 2019. The causal relationship between sexual selection and sexual size dimorphism in marine gastropods. Animal Behaviour 148:53-62.
Parker, G. A. 2014. The sexual cascade and the rise of pre-ejaculatory (Darwinian) sexual selection, sex roles, and sexual conflict. Cold Spring Harbor perspectives in biology 6:a017509-a017509.
Parker, G. A. and T. Pizzari. 2015. Sexual Selection: The Logical Imperative. Pp. 119-163 *in* T. Hoquet, ed. Current Perspectives on Sexual Selection: What's left after Darwin? Springer Netherlands, Dordrecht.
Press, W. H. 2002. Numerical recipes in C++ : the art of scientific computing. Cambridge University Press, Cambridge.
Prum, R. O. 2012. Aesthetic evolution by mate choice: Darwin's really dangerous idea. Philosophical Transactions of the Royal Society of London B: Biological Sciences 367:2253-2265.

Richards, S. A. 2008. Dealing with overdispersed count data in applied ecology. Journal of Applied Ecology 45:218-227.

Rolan-Alvarez, E. 2007. Sympatric speciation as a by-product of ecological adaptation in the Galician Littorina saxatilis hybrid zone. Journal of Molluscan Studies 73:1-10.

Rolan-Alvarez, E., C. Austin, and E. G. Boulding. 2015a. The contribution of the genus Littorina to the field of evolutionary ecology. Oceanography and Marine Biology: an Annual Review 53:157-214.

Rolán-Alvarez, E. and A. Caballero. 2000. Estimating sexual selection and sexual isolation effects from mating frequencies. Evolution 54:30-36.

Rolan-Alvarez, E., A. Carvajal-Rodriguez, A. de Coo, B. Cortes, D. Estevez, M. Ferreira, R. Gonzalez, and A. D. Briscoe. 2015b. The scale-of-choice effect and how estimates of assortative mating in the wild can be biased due to heterogeneous samples. Evolution 69:1845-1857.

Rosenthal, G. G. 2017. Mate choice: the evolution of sexual decision making from microbes to humans. Princeton University Press.

Roughgarden, J., M. Oishi, and E. Akçay. 2006. Reproductive social behavior: cooperative games to replace sexual selection. Science 311:965-969.

Schwarz, G. 1978. Estimating the dimension of a model. The Annals of Statistics 6:461-464.

Shuker, D. M. 2010. Sexual selection: endless forms or tangled bank? Animal Behaviour 79:e11-e17.

Singer, S. and S. Singer. 2004. Efficient Implementation of the Nelder–Mead Search Algorithm. Applied Numerical Analysis & Computational Mathematics 1:524-534.

Swihart, R. K., J. B. Dunning, and P. M. Waser. 2002. Gray Matters in Ecology: Dynamics of Pattern, Process, and Scientific Progress. Bulletin of the Ecological Society of America 83:149-155.

Symonds, M. R. E. and A. Moussalli. 2011. A brief guide to model selection, multimodel inference and model averaging in behavioural ecology using Akaike's information criterion. Behavioral Ecology and Sociobiology 65:13-21.

Vrieze, S. I. 2012. Model selection and psychological theory: a discussion of the differences between the Akaike information criterion (AIC) and the Bayesian information criterion (BIC). Psychological methods 17:228.

Wacker, S. and T. Amundsen. 2014. Mate competition and resource competition are inter-related in sexual selection. Journal of evolutionary biology 27:466-477.

Wagenmakers, E.-J. 2007. A practical solution to the pervasive problems of p values. Psychonomic bulletin & review 14:779-804.