

Host phenology can drive the evolution of intermediate virulence strategies in some parasites

Hannelore MacDonald^{1,*}, Erol Akçay¹, Dustin Brisson¹

1. Department of Biology, University of Pennsylvania, Philadelphia, Pennsylvania 19104;

* Corresponding author; e-mail: haime@sas.upenn.edu

1 Abstract

Mechanistic trade-offs resulting in a negative correlation between transmission and virulence are the foundation of nearly all current theory on the evolution of parasite virulence. Although several ecological factors have been shown to modulate the optimal virulence strategies predicted from mechanistic trade-off models, none have been shown to be sufficient to explain the intermediate virulence strategies observed in any natural system. The timing of seasonal activity, or phenology, is a common factor that influences the types and impact of many ecological interactions but is rarely considered in virulence evolution studies. We develop a mathematical model of a disease system with seasonal host activity to study the evolutionary consequences of host phenology on parasite virulence. Results from this model demonstrated that seasonal host activity is sufficient to drive the evolution of intermediate parasite virulence, in the absence of traditional mechanistic trade-offs, in some types of natural disease systems. The optimal virulence strategy in these systems can be determined by both the duration of the host activity period as well as the variation in the host emergence timing. Parasites with low virulence strategies are favored in environments with long host activity periods and in environments in which hosts emerge synchronously. The results demonstrate that host phenology can be sufficient to select for intermediate optimal virulence strategies, providing an alternative mechanism to account for virulence evolution in some natural systems.

2 Introduction

The evolutionary causes and consequences of parasite virulence remain enigmatic despite decades of research. It was once thought that parasites continue to evolve ever lower levels of virulence to preserve their primary resource for future parasite generations.¹ However, the idea that parasites would limit damaging their host for the benefit of future generations violates multiple core principles of our modern understanding of evolutionary biology.^{2,3} Natural selection, as framed in the modern evolutionary synthesis, favors traits that improve short-term evolutionary fitness even if those traits negatively impact the environment for future generations.^{2,3} Thus, the level of virulence that maximizes parasite fitness is favored by natural selection despite its impact on the host population.⁴ Identifying environmental conditions and mechanistic constraints that drive the evolution of the vast diversity of parasite virulence strategies observed in nature has been an important research focus for decades.

Establishing that mechanistic trade-offs constrain parasite virulence strategies was a key breakthrough that propels virulence evolution research to this day. In the now classic paper, Anderson and May demonstrated that within-host parasite densities increase the probability of transmission to a new host - a component of parasite fitness - but also shorten infection duration resulting in fewer opportunities for transmission to naive hosts.⁴ That is, parasites can only produce more infectious progeny if they cause host damage by utilizing more host resources. This mechanistic trade-off that results in a negative correlation between virulence and transmission remains the foundational theoretical framework used to account for the evolution of all parasite virulence strategies observed in nature.^{5,6}

Many ecological factors and environmental conditions have been shown to alter the optimal virulence strategies driven by mechanistic trade-offs within models. For example, it is well-established that varying environmental conditions, such as the extrinsic host death rate, often shift the optimal virulence strategy governed by a mechanistic trade-off.^{4,7-9} However, no environmental condition, *in the absence* of mechanistic trade-offs, has been shown to select for intermediate virulence.

The timing of seasonal activity, or phenology, is an environmental condition affecting all aspects of life cycles, including reproduction, migration, and diapause, in most species.¹⁰⁻¹⁶ The phenology of host species also impacts the timing and prevalence of transmission opportunities for parasites which could alter optimal virulence strategies.¹⁷⁻²⁴ For example, host phenological patterns that extend the time between infection and transmission are expected to select for lower virulence, as observed in some malaria parasites (*Plasmodium vivax*). In this system, high virulence strains persist in regions where mosquitoes are present year-round while low virulence strains are more common in regions where mosquitoes are nearly absent during the dry season.²⁵ While host phenology likely impacts virulence evolution in parasites,²⁶⁻²⁹ it remains unclear whether this environmental condition can have a sufficiently large impact to select for an intermediate virulence phenotype in the absence of a mechanistic trade-off.

Here we investigate the impact of host phenology on the virulence evolution of an obligate-killer parasite. We demonstrate that intermediate virulence is adaptive when host activity patterns are highly seasonal in the absence of any explicit mechanistic trade-off, establishing that environmental context alone is sufficient to drive the evolution of intermediate virulence in disease systems that conform to the assumptions of the model. Further, multiple features of host seasonal activity, including season length and the synchronicity at which hosts first become active during the season, impact the optimal virulence level of parasites. These results provide an alternative framework that can account for virulence evolution in some natural systems.

Parameter	Description	Value
s	susceptible hosts	state variable
v_1	parasites that infect hosts in current season	state variable
v_2	parasites released in current season	state variable
t_l	length of host emergence period	time (varies)
T	season length	time (varies)
\hat{s}	emerging host cohort size	10^8 hosts
α	transmission rate	$10^{-8}/(\text{parasite} \times \text{time})$
β	number of parasites produced upon host death	parasites (varies)
δ	parasite decay rate in the environment	2 parasites/parasite/time
d	host death rate	0.5 hosts/host/time
τ	time between host infection and host death (1/virulence)	time (evolves)

Table 1: Model parameters and their respective values.

3 Model description

The model describes the transmission dynamics of a free-living, obligate-killer parasite that infects a seasonally available host (Figure 1). The host cohort, \hat{s} , enters the system at the beginning of the season over a period given by the function $g(t, t_l)$. Hosts, s , have non-overlapping generations and are alive for one season. The parasite, v , infects hosts while they are briefly susceptible early in their development (*e.g.* baculoviruses of forest *Lepidoptera*^{30–34} and univoltine insects parasitized by ichneumonids^{35–37}). The parasite must kill the host to release new infectious progeny. The parasite completes one round of infection per season because the incubation period of the parasite is longer than the duration of time the host spends in the susceptible developmental stage. This transmission scenario occurs in nature if all susceptible host stages emerge over a short period of time each season so that there are no susceptible host stages available when the parasite eventually kills its host. Parasites may also effectively complete only one round of infection per season if the second generation of parasites do not have enough time in the season to complete their life cycle in the short-lived host.

We ignore the progression of the susceptible stage, s , to later life stages as it does not impact transmission dynamics. To keep track of these dynamics, we refer to the generation of parasites that infect hosts in the beginning of the season as v_1 and the generation of parasites released from infected hosts upon parasite-induced death as v_2 . The initial conditions in the beginning of the season are thus $s(0) = 0, v_1(0^+) = v_2(0^-), v_2(\tau) = 0$. The transmission dynamics in season n are given by the following system of delay differential equations (all parameters are described in Table 1):

$$\frac{ds}{dt} = \hat{s}g(t, t_l) - ds(t) - \alpha s(t)v_1(t), \quad (1a)$$

$$\frac{dv_1}{dt} = -\delta v_1(t), \quad (1b)$$

$$\frac{dv_2}{dt} = \alpha\beta e^{-d\tau} s(t - \tau)v_1(t - \tau) - \delta v_2(t). \quad (1c)$$

where d is the host death rate, δ is the decay rate of parasites in the environment, α is the transmission rate, β is the number of parasites produced upon host death and τ is the delay between host infection and host death. τ is equivalent to virulence where low virulence parasites have long τ and high virulence parasites have short τ . We make the common assumption for free-living parasites that the removal of parasites through transmission (α) is negligible,^{34,38,39} *i.e.* (1b) ignores the term $-\alpha s(t)v_1(t)$.

60 The function $g(t, t_l)$ is a probability density function that captures the per-capita host emergence rate by
 61 specifying the timing and length of host emergence. We use a uniform distribution ($U(\bullet)$) for analytical
 62 tractability, but other distributions can be used.

$$g(t, t_l) = \begin{cases} \frac{1}{t_l} & 0 \leq t \leq t_l \\ 0 & t_l < t \leq T \end{cases}$$

63 t_l denotes the length of the host emergence period and T denotes the season length. The season begins
 64 ($t_0 = 0$) with the emergence of the susceptible host cohort, \hat{s} . The host cohort emerges from $0 \leq t \leq t_l$. v_2
 65 parasites remaining in the system at $t = T$ give rise to next season's initial parasite population ($\hat{v} = v_1(0)$).
 66 Parasites that have not killed their host by the end of the season do not release progeny. Background mortality
 67 arises from predation or some other natural cause. We assume that infected hosts that die from background
 68 mortality do not release parasites because the parasites are either consumed or the latency period corresponds
 69 to the time necessary to develop viable progeny.^{40,41} We ignore the impact of infection for host demography
 70 and assume \hat{s} is constant each year (*e.g.* a system where host regulation by parasites is negligible). We solve
 71 equations 1a-c analytically Appendix A.

72 3.0.1 Parasite fitness

A parasite introduced into a naive host population persists or goes extinct depending on the length of the
 host emergence period and season length. The stability of the parasite-free equilibrium is determined by the




Figure 1: **Infection diagram** The host cohort, \hat{s} , emerges from time $t = 0$ to $t = t_l$, all v_1 parasites emerge at $t = 0$. Hosts do not reproduce during the season. Infections generally occur early in the season when host density is high. Parasite-induced host death occurs after time τ , at which point new parasites, v_2 are released. v_2 decays in the environment from exposure. Parasites only have time to complete one round of infection per season. v_2 parasites in the environment at $t = T$ will carryover and emerge at the beginning of the next season.

production of v_2 resulting from infection of s given by

$$v_2(T) = e^{-\delta(T-t_l-\tau)}(v_2(t_l) + \alpha\beta e^{-d\tau} \hat{v}s(t_l) \int_0^{T-t_l-\tau} e^{-\frac{\alpha\hat{v}e^{-\delta(s+t_l)}(-1+e^{\delta s})}{\delta} - \delta t_l - ds} ds)$$

73 The parasite-free equilibrium is unstable and a single parasite introduced into the system at the beginning
 74 of the season will persist if the density of v_2 produced by time T is greater than or equal to $\hat{v} = v_1(0) = 1$
 75 (i.e. $v_2(T) \geq 1$, modulus is greater than unity). This expression is a measure of a parasite's fitness when rare
 76 given different host phenological patterns. See Appendix A for details of the analytical solution.

77 3.0.2 Parasite evolution

78 To study how parasite traits adapt given different seasonal host activity patterns, we use evolutionary invasion
 79 analysis.^{42,43} We first extend system (1) to follow the invasion dynamics a rare mutant parasite

$$\frac{ds}{dt} = \hat{s}g(t, t_l) - ds(t) - \alpha s(t)v_1(t) - \alpha_m s(t)v_{1m}(t), \quad (2a)$$

$$\frac{dv_1}{dt} = -\delta v_1(t), \quad (2b)$$

$$\frac{dv_{1m}}{dt} = -\delta_m v_{1m}(t), \quad (2c)$$

$$\frac{dv_2}{dt} = \alpha\beta s(t-\tau)v_1(t-\tau) - \delta v_2(t), \quad (2d)$$

$$\frac{dv_{2m}}{dt} = \alpha_m\beta_m s(t-\tau_m)v_{1m}(t-\tau_m) - \delta_m v_{2m}(t). \quad (2e)$$

80 where m subscripts refer to the invading mutant parasite and its corresponding traits. See Appendix B for
 81 details of the time-dependent solutions for equations (2a-2e).

82

83 The invasion fitness of a rare mutant parasite depends on the density of v_{2m} produced by the end of the
 84 season ($v_{2m}(T)$) in the environment set by the resident parasite at equilibrium density \hat{v}^* . The mutant
 85 parasite invades in a given host phenological scenario if the density of v_{2m} produced by time T is greater
 86 than or equal to the initial $v_{1m}(0) = 1$ introduced at the start of the season ($v_{2m}(T) \geq 1$).

$$v_{2m}(T) = e^{-\delta_m(T-t_l-\tau_m)}(v_{2m}(t_l) + \alpha_m\beta_m e^{-d\tau_m} v_{1m}(0)s(t_l) \int_0^{T-t_l-\tau_m} e^{-\frac{\alpha_m v_{1m}(0)e^{-\delta_m(s+t_l)}(-1+e^{\delta_m s})}{\delta_m} - \frac{\alpha\hat{v}^* e^{-\delta(s+t_l)}(-1+e^{\delta s})}{\delta} - \delta_m t_l - ds} ds) \quad (3)$$

87 To study the evolution of virulence traits, we first assume all other resident and mutant traits are identical
 88 (e.g. $\alpha = \alpha_m$). Note that when there is no trade-off between β and τ , the parasite growth rate in the host is
 89 essentially the trait under selection. That is, β is constant regardless of τ thus the time between infection and
 90 when the parasite kills the host and releases new parasites is the rate that β new parasites are assembled.
 91 To find optimal virulence for a given host phenological scenario, we find the uninvadable trait value that
 92 maximizes (3). That is, the virulence trait, τ^* , that satisfies

$$\frac{\partial v_{2m}(T)}{\partial \tau_m} \Big|_{\tau_m = \tau_r} = 0 \quad (4a)$$

$$\frac{\partial^2 v_{2m}(T)}{\partial \tau_m^2} \Big|_{\tau_m = \tau_r} < 0 \quad (4b)$$

93 Note that the measure in equation (3) incorporates the effect of the resident on the population state (number
 94 of susceptibles over one season), which means that it is not a measure of R_0 (which by definition assumes
 95 a non-disease environment). Thus, we can use $v_{2m}(T)$ as defined in (3) as a maximand in evolutionary
 96 dynamics.⁴⁴

97 To study the impact of mechanistic trade-offs between transmission and virulence on virulence evolution, we
 98 assume that the number of parasites produced at host death is a function of the time between infection and
 99 host death ($\beta(\tau)$)

$$v_{2m}(T) = e^{-\delta_m(T-t_i-\tau_m)}(v_{2m}(t_i) + \alpha_m \beta(\tau_m) e^{-d\tau_m} v_{1m}(0) s_1(t_i) \\ \int_0^{T-t_i-\tau_m} e^{-\frac{\alpha_m v_{1m}(0) e^{-\delta_m(s+t_i)}(-1+e^{-\delta_m s})}{\delta_m} - \frac{\alpha v^* e^{-\delta(s+t_i)}(-1+e^{-\delta s})}{\delta} - \delta_m t_i - ds} ds). \quad (5)$$

100 We then find τ^* that satisfies (4a) and (4b) using equation (5).

4 Results

Host phenology, in the absence of a mechanistic trade-off, is sufficient to drive the evolution of intermediate virulence in systems that conform to the assumptions of the model. Host phenology is composed of the duration of the activity period and the distribution of initial emergence times, both of which impact the optimal parasite virulence level. Temporally constrained host activity periods within each season can select against both extremely high and extremely low virulence levels resulting in an intermediate optimal level of virulence. Low virulence is selected against as parasites that do not kill the infected host prior to the end of the host activity period fail to produce progeny and thus have no evolutionary fitness. By contrast, highly-virulent parasites kill their hosts quickly and the released progeny decay in the environment for the remainder of the activity period. Thus, progeny released early in the host activity period are more likely to die in the environment prior to contacting a naive host in the following season. An intermediate virulence level that allows parasites to kill their host prior to the end of the activity period, but not so quickly that the progeny produced are likely to decay in the environment, result in the greatest evolutionary fitness.

The optimal virulence level increases linearly with decreases in the duration of host activity (Figure 2). Virulent parasites in environments where host activity periods are short minimize the cost of not producing progeny from infected hosts and do not incur the costs of progeny decaying in the environment. By contrast, environments where host activity periods are long favor parasites with a long incubation period to limit the cost of progeny decay due to environmental exposure while still killing hosts prior to the end of the season. The optimal level of virulence in all environmental scenarios results in parasite-induced host death just prior to the end of the seasonal activity period. The linear increase in optimal virulence as season length decreases suggests that parasite fitness is optimized when host death occurs at a fixed time before the end of the season.

Variation in the time at which each host first becomes active during the activity period also impacts the virulence levels that maximize parasite fitness (Figure 3). Synchronous host emergence results in a rapid and early spike in infection incidence due to the simultaneous availability of susceptible hosts and the abundance of free parasites. The long duration between host infection and the end of the activity period favors low virulence parasites that kill their host near the end of the season (Figure 3A, *i*). Variability in the time at which each susceptible host initially becomes active decreases the average time between infection and the end of the season, thus favoring more virulent parasites (Figure 3A, *ii*). That is, the large proportion of infections that occur later in the season require higher virulence to be able to release progeny before the activity period ends. This higher virulence level comes at the cost of progeny from hosts infected early in the season decaying in the environment. Thus, the number of progeny that survive to the next season decreases with increasing variation in host emergence times (Figure 3B).

High variability in host emergence timing results in an optimal virulence strategy that is much greater than in environments with synchronous host emergence, but lower than in environments with a moderate distribution (Figure 3). That is, increasing variation in host emergence timing favors parasites with higher virulence, but only when variation in host emergence timing is high. In environments where the variation in host emergence timing is high, increasing variation in host emergence timing favors parasites with slightly lower virulence (Figure 3A, *iii*). Lower virulence is favored in high emergence variability environments because the number of new infections occurring late in the season, where high virulence would be advantageous, are relatively rare due to small parasite population sizes at the beginning of the season and parasite decay during the season. Initial parasite population sizes are smaller in environments with broadly distributed host emergence timing as fewer total hosts are infected because infection probability is density dependent,

143 and thus fewer progeny are produced. Most parasites that find a susceptible host do so early in the season
144 resulting in additional decreases to the already small parasite population size. The optimal virulence strategy
145 allows parasites that infect hosts around the peak of new infections - occurring mid-season when susceptible
146 host densities are greatest and parasite populations have not decayed substantially - to release progeny while
147 limiting decay of these progeny. Parasites in environments where the distribution in host emergence times is
148 very broad suffer the costs of both decay of the progeny released by early-infected hosts and the cost of late
149 infected hosts not releasing progeny, collectively causing these environments to maintain low densities of
150 moderately virulent pathogens (Figure 3B, *iii*).

151 Mechanistic virulence-transmission trade-offs can modify the optimal virulence strategy in seasonal
152 environments but are not necessary for natural selection to favor intermediate virulence phenotypes. The
153 optimal virulence strategy is slightly lower in models that include a trade-off where duration of infection is
154 positively correlated with progeny production than in models with the same phenological parameters that do
155 not include the trade-off (Figure 4). Including this trade-off increases the fitness benefit of longer-duration
156 infections to a greater extent than the costs associated with infected host mortality not caused by the parasite.
157 By contrast, the optimal virulence strategy is greater in models that include a trade-off where duration of
158 infection is negatively correlated with progeny production than in similar models without the trade-off
159 (Figure 4). Including this trade-off increases the fitness benefit of shorter-duration infections despite the
160 added costs of greater parasite decay due to environmental exposure. Including mechanistic trade-offs
161 modifies the selection pressures on virulence strategies but are not essential for an intermediate virulence
162 strategy to be optimal in seasonal environments.




Figure 2: **Host seasonality is sufficient to select an intermediate virulence strategy.** **A.** The temporal duration between infection and host death (τ^*) always evolves to a value that is greater than 0 (extreme virulence) and less than the season length (extremely low virulence); the intermediate virulence strategy maximizes parasite fitness in environments where host activity is seasonal. The optimal parasite-induced host death rate results in host death and progeny release shortly before the end of the season ($t = T$). The release of progeny just prior to the end of the season limits the decay of progeny due to environmental exposure while avoiding progeny dying within their host at the end of the season. *i* and *ii* are representative examples of optimal virulence strategies in environments with shorter ($T = 3.2$) or longer ($T = 4$) host activity periods, respectively. τ^* is found using equation (4a) when there is no trade-off between transmission and virulence. **B.** Higher parasite virulence is favored in environments with limited host activity periods. Parasites with greater virulence produce more progeny that survive to the end of the season when seasons are short. That is, the density of the more virulent progeny (*i*) at $T = 3.2$ is greater than the density of the less virulent progeny (*ii*). The more virulent parasites kill their hosts quickly such that few infected hosts survive to the end of the season and the progeny released spend little time in the environment. By contrast, less virulent parasites (*ii*) often fail to kill their hosts and release progeny prior to the end of short activity periods ($T = 3.2$). Longer seasons ($T = 4$) favor less virulent parasites (*ii*) as they kill their hosts closer to the end of the season such that fewer of their released progeny decay in the environment (*ii*) than the progeny of the more virulent parasites that are released earlier in the season (*i*). The blue line represents the incidence rate of new infections; $t_l = 1$; all other parameters found in Table 1.

5 Discussion

Nearly all theory developed to explain parasite virulence evolution has utilized mechanistic trade-offs between virulence and other traits important to parasite fitness.^{5,6} The results of this study show that seasonal host activity, in the absence of a mechanistic trade-off, can account for **why intermediate virulence is adaptive** the evolution of intermediate virulence in some specific situations. Both aspects of phenology, the duration of the host activity period and host emergence synchronicity, impact the virulence strategy that maximizes the evolutionary fitness of parasites. Although mechanistic trade-offs between virulence and transmission can shift the optimal virulence level as predicted by prior theory, these trade-offs are not essential for intermediate virulence to evolve in this system. The current demonstration that an ecological context **alone** is sufficient to select for intermediate virulence broadens the scope of factors that can explain the diversity of parasite virulence strategies. Thus, the evolution of intermediate virulence in natural systems may be governed by a mechanistic trade-off or by ecological factors **alone** in some systems.

Seasonal host activity can select for intermediate virulence by generating conflicting costs for releasing progeny too early or too late in the season. Low virulence is maladaptive for parasites in this system as they do not kill their host before the end of the season and create no progeny. High virulence is also maladaptive as progeny released early are more likely to die due to environmental exposure. The conflicting costs of not releasing progeny before the end of the season and releasing progeny too early in the season selects for




Figure 3: **The variation in host emergence timing impacts the optimal virulence strategy.** **A.** Parasites with lower virulence are favored in environments where nearly all hosts emerge simultaneously (*i*). Progeny from the low virulence parasites are released nearly simultaneously just prior to the end of the season. High virulence parasites are favored in environments where host emergence period length is moderate (*ii*). Moderate variation in host emergence decreases the average time between infection and the end of the season and favors parasites with a high virulence strategy such that few infected hosts survive to the end of the season. Parasites in environments where host emergence variation is very high maximize the number of progeny that survive to the next season by using a moderate virulence strategy (*iii*). Parasites in these environments suffer the costs of hosts that are infected later in the season not releasing progeny as well as progeny decay in the environment when released from early-infected hosts. A moderate virulence strategy allows hosts infected around the mid-season peak in incidence to release progeny while limiting the decay of these progeny. τ^* is found using equation (4a) when there is no trade-off between transmission and virulence. **B.** Equilibrium density of parasites with the optimal virulence strategy for their environment decreases with increasing variation in host emergence timing. Optimal virulence results in peak equilibrium in new parasites density, indicated by the vertical lines. $T = 3$; other parameters found in Table 1.

180 intermediate virulence levels. Optimal virulence results in parasite-induced host death and the release of
 181 progeny slightly before the end of the host activity period. ~~These results suggest that seasonal host absence~~
 182 ~~can increase the~~

183 The result predicting adaptive evolution towards intermediate virulence stands in contrast to many
 184 prior theoretical investigations of obligate-killer parasites. Prior models of obligate-killer parasites predict
 185 ever-increasing virulence in the absence of mechanistic trade-offs^{45–48}. In simple obligate-killer models,
 186 killing infected hosts as quickly as possible is expected to maximize fitness as the early release of progeny
 187 permits infection of additional susceptible hosts resulting in a rapid exponential increase of parasites in the
 188 system. To date, only mechanistic trade-offs between virulence and transmission-associated factors as well
 189 as development time constraints have been demonstrated to constrain maximal virulence in obligate-killer
 190 parasite models^{40,46,48–50}. In contrast, our results indicate that host phenology - in the absence of an explicit
 191 mechanistic trade-off and development time constraints - can create the conditions that favor intermediate
 192 virulence in obligate-killer parasites. In the current model, intermediate virulence is favored as seasonal




Figure 4: **Mechanistic transmission-virulence trade-offs shift the optimal virulence strategy but are not necessary to favor intermediate virulence in environments with seasonal host activity.** The optimal virulence level for parasites in which longer durations of infection result in *more* progeny is slightly lower than for parasites that are not constrained with this mechanistic trade-off in the same environment (*i*). This mechanistic trade-off elevates the fitness benefit of longer duration infections by compensating for the cost of infected hosts dying without releasing progeny. The optimal virulence level for parasites in which longer infection durations result in *fewer* progeny is greater than for parasites without this trade-off in the same seasonal environments (*ii*). This mechanistic trade-off elevates the fitness benefit of shorter duration infections despite the cost of greater progeny decay in the environment. τ^* was found using equation (4a) when there is no trade-off between transmission and virulence and then compared to τ^* constrained by a trade-off with transmission. Trade-off for *i* : $\beta(\tau) = 99(\tau + 0.5)$, trade-off for *ii* : $\beta(\tau) = 99(-\tau + 4)$. All other parameters found in Table 1.

193 [host absence increases the](#) evolutionary benefit of remaining within hosts in order to reduce deaths in the
194 free-living stage caused by environmental exposure⁵¹.

195 Variation in host emergence synchronicity impacts the optimal virulence strategy of parasites in this
196 system. High parasite virulence is favored at low host emergence synchronicity. Low emergence synchronicity
197 slows incidence by decreasing both the rate hosts emerge and parasite equilibrium density. When more
198 infections occur later in the season, parasites have less time to release new parasites before the end of the
199 season. High parasite virulence is adaptive because hosts have a low expected life span at the time of infection.
200 This result is analogous to the prediction that high host mortality drives the evolution of high virulence.^{4,7-9}
201 The timing of host activity can thus lead to the evolution of high virulence in a similar manner to how host
202 demography impacts virulence.

203 The seasonal activity patterns of species with non-overlapping generations may have large impacts on the
204 virulence strategies of the parasites they host. For example, parasites and parasitoids of univoltine insects
205 that complete one round of infection per host generation may maximize their fitness by releasing progeny just
206 prior to the end of the season.³⁰⁻³⁷ The theoretical expectations presented here can be tested empirically by
207 measuring the virulence strategies of parasites across the natural diversity of phenological patterns observed
208 over the geographical range of many insect species. Similarly, experiments could rigorously assess the impact
209 of both season length and host emergence variability on the fitness of parasites with different levels of
210 virulence.

211 The prediction that shorter host activity periods can drive greater virulence is comparable to how the
212 virulence of different *Theileria parva* strains varies between regions. High within-host densities permit a
213 virulent *T. parva* strain to be reliably transmitted to feeding nymphal tick vectors shortly after being infected
214 by the adult stage in regions where the activity patterns of the two tick life stages overlap.⁵²⁻⁵⁴ In contrast, the
215 virulent strain is absent in regions where nymphal and adult activity is asynchronous while a less virulent
216 strain that persists in hosts longer is maintained.^{52,53} Thus, the prediction that the length of the host activity
217 period is inversely correlated with virulence coincides with empirical observations of the distribution of *T.*
218 *parva* strains.

219 Several features of the current model can be altered to investigate more complex impacts of phenology
220 on virulence evolution. For example, relaxing the assumption of a constant host population size may result
221 in a feedback between parasite fitness and host demography with consequences for population dynamics.
222 Additionally, parasite virulence evolution may select for alternative host phenological patterns that in turn
223 select for parasite traits with lower impacts on host fitness. We will extend the current model to address
224 these questions in future studies.

225 The model presented applies to obligate-killer parasites that complete one round of infection per season
226 (monocyclic) in hosts that have non-overlapping generations. Currently, there is no evidence that disease
227 systems that do not conform to these assumptions will not require a mechanistic trade-off to select for
228 intermediate virulence. Nevertheless, several prior models that included both host seasonality and mechanistic
229 trade-offs found qualitatively similar results as those presented here despite relaxing one or more of the
230 strict assumptions in this model,²⁶⁻²⁸ suggesting that phenology can have a large impact on virulence
231 outcomes. For example, longer seasons or longer periods between seasons have been shown to select for lower
232 virulence in polycyclic parasites in seasonal environments,^{27,28} similar to the results presented here. Similarly,
233 explicitly modeling parasite growth rates within hosts, which underlie the correlation between virulence and
234 instantaneous transmission rates, selects for intermediate virulence levels that maximize transmission rates
235 during host activity periods.²⁶ By contrast, assuming that virulence levels are mechanistically associated with

236 host density results in selection for higher virulence in seasonal environments.²⁹ Future studies incorporating
237 one or more of these competing forces with environmental decay of progeny could be sufficient to select for
238 intermediate virulence in the absence of an explicit mechanistic trade-off.

239 Some of the strict model assumptions can likely be relaxed without altering the result that phenology can
240 be sufficient to select for intermediate virulence strategies. Relaxing the obligate-killer assumption may result
241 in the same qualitative result that intermediate virulence is adaptive in some cases. For example, longer
242 latency periods that result in progeny release near the end of the season would still be adaptive for parasites
243 that reduce host fecundity or increase host death rate, even if there is no correlation between the virulence
244 level and instantaneous or life-time transmission. Longer latency periods are equivalent to lower virulence in
245 this type of system as infected hosts have more time to reproduce and thus higher fitness. This extension is
246 not expected to qualitatively alter the results if the parasite transmission period is short relative to the season
247 length. Many parasite-host systems conform to the assumptions of this model extension such as soil-borne
248 plant pathogens, demicyclic rusts, post-harvest diseases, and many diseases systems infecting univoltine
249 insects.^{55–58}

250 The importance of parasite virulence to both host-parasite interactions and public health policy has
251 resulted in a concentrated research effort on virulence evolution. Nearly all theoretical research to date
252 has incorporated a mechanistic trade-off between virulence and transmission rates or infection duration, a
253 hypothesis which is still essential to explain the evolution of intermediate virulence in most disease systems.
254 However, ecological factors such as seasonal host activity or spatial structuring provide alternative theoretical
255 frameworks that may account for virulence strategies in some natural systems.^{59,60} Future work that identifies
256 and empirically validates ecological factors that influence virulence evolution would be useful for predicting
257 outbreaks of highly virulent parasites.

258 6 Code and data availability

259 Code is available on the Github repository: [https://github.com/hanneloremac/Host-phenology-drives-the-
260 evolution-of-intermediate-parasite-virulence](https://github.com/hanneloremac/Host-phenology-drives-the-evolution-of-intermediate-parasite-virulence)

261 References

- 262 ¹ Theobald Smith. Some problems in the life history of pathogenic microorganisms. *Science*, 20(520):817–832,
263 1904.
- 264 ² WD Hamilton. The genetical evolution of social behaviour. i. *Journal of theoretical biology.*, 7(1).
- 265 ³ WD Hamilton. The genetical evolution of social behaviour. ii. *Journal of theoretical biology.*, 7(1).
- 266 ⁴ R M Anderson and R M May. Coevolution of hosts and parasites. *Parasitology*, 85(Pt 2):411–426, 1982.
- 267 ⁵ Clayton E Cressler, David V McLEOD, Carly Rozins, Josée van den Hoogen, and Troy Day. The adaptive
268 evolution of virulence: a review of theoretical predictions and empirical tests. *Parasitology*, 143(7):915–930,
269 2016.
- 270 ⁶ S Alizon, A Hurford, N Mideo, and M Van Baalen. Virulence evolution and the trade-off hypothesis:
271 history, current state of affairs and the future. *J Evol Biol*, 22(2):245–259, 2009.

- 272 ⁷ Richard E Lenski and Robert M May. The evolution of virulence in parasites and pathogens: Reconciliation
273 between two competing hypotheses. Journal of Theoretical Biology, 169:253–265, 1994.
- 274 ⁸ S Gandon, V A Jansen, and M Van Baalen. Host life history and the evolution of parasite virulence.
275 Evolution, 55(5):1056–1062, 2001.
- 276 ⁹ VS Cooper, MH Reiskind, JA Miller, KA Shelton, BA Walther, JS Elkinton, and PW Ewald. Timing of
277 transmission and the evolution of virulence of an insect virus. Proc Biol Sci, 269(1496):1161–1165, 2002.
- 278 ¹⁰ Jelmer A. Elzinga and et al. Time after time: flowering phenology and biotic interactions. Trends in Ecology
279 Evolution, 2007.
- 280 ¹¹ Jessica Forrest and Abraham J Miller-Rushing. Toward a synthetic understanding of the role of phenology
281 in ecology and evolution. Philosophical transactions of the Royal Society of London. Series B, Biological
282 sciences, 365(1555):3101–3112, 2010.
- 283 ¹² Stephanie Pau, Elizabeth M. Wolkovich, Benjamin I. Cook, T. Jonathan Davies, Nathan J. B. Kraft, Kjell
284 Bolmgren, Julio L. Betancourt, and Elsa E. Cleland. Predicting phenology by integrating ecology, evolution
285 and climate science. Global Change Biology, 17(12):3633–3643, 2011.
- 286 ¹³ JT Anderson, DW Inouye, AM McKinney, RI Colautti, and T Mitchell-Olds. Phenotypic plasticity and
287 adaptive evolution contribute to advancing flowering phenology in response to climate change. Proc Biol
288 Sci, 279(1743):3843–3852, 2012.
- 289 ¹⁴ A Novy, SL Flory, and JM Hartman. Evidence for rapid evolution of phenology in an invasive grass. J Evol
290 Biol, 26(2):443–450, 2013.
- 291 ¹⁵ N Lustenhouwer, RA Wilschut, JL Williams, WH van der Putten, and JM Levine. Rapid evolution of
292 phenology during range expansion with recent climate change. Glob Chang Biol, 24(2):e534–e544, 2018.
- 293 ¹⁶ JS Park. Cyclical environments drive variation in life-history strategies: a general theory of cyclical
294 phenology. Proc Biol Sci, 286(1898):20190214, 2019.
- 295 ¹⁷ Arjen Biere and Sonja J. Honders. Impact of flowering phenology of silene alba and s. dioica on susceptibility
296 to fungal infection and seed predation. Oikos, 77(3):467–480, 1996.
- 297 ¹⁸ Sonia Altizer, Andrew Dobson, Parvize Hosseini, Peter Hudson, Mercedes Pascual, and Pejman Rohani.
298 Seasonality and the dynamics of infectious diseases. Ecology Letters, 9(4):467–484, 2006.
- 299 ¹⁹ Sarah A. Hamer, Graham J. Hickling, Jennifer L. Sidge, Edward D. Walker, and Jean I. Tsao. Synchronous
300 phenology of juvenile ixodes scapularis, vertebrate host relationships, and associated patterns of borrelia
301 burgdorferi ribotypes in the midwestern united states. Ticks and Tick-borne Diseases, 3(2):65–74, 2012.
- 302 ²⁰ OJ Gethings, H Rose, S Mitchell, J Van Dijk, and ER Morgan. Asynchrony in host and parasite phenology
303 may decrease disease risk in livestock under climate warming: Nematodirus battus in lambs as a case
304 study. Parasitology, 142(10):1306–1317, 2015.
- 305 ²¹ Nicholas H Ogden, Genevieve Pang, Howard S Ginsberg, Graham J Hickling, Russell L Burke, Lorenza
306 Beati, and Jean I Tsao. Evidence for geographic variation in life-cycle processes affecting phenology of the
307 lyme disease vector ixodes scapularis (acari: Ixodidae) in the united states. Journal of Medical Entomology,
308 55(6):1386–1401, 2018.

- 309 ²² Micaela Elvira Martinez. The calendar of epidemics: Seasonal cycles of infectious diseases. PLOS Pathogens,
310 14(11):e1007327–15, 2018.
- 311 ²³ Travis McDevitt-Galles, Wynne E Moss, Dana M Calhoun, and Pieter TJ Johnson. Phenological synchrony
312 shapes pathology in host–parasite systems. Proceedings of the Royal Society B, 287(1919):20192597, 2020.
- 313 ²⁴ Hannelore MacDonald, Erol Akçay, and Dustin Brisson. The role of host phenology for parasite transmission.
314 Theoretical Ecology, 2020.
- 315 ²⁵ Michael T White, George Shirreff, Stephan Karl, Azra C Ghani, and Ivo Mueller. Variation in relapse
316 frequency and the transmission potential of Plasmodium vivax malaria. Proceedings of the Royal Society
317 B: Biological Sciences, 283(1827):20160048–9, March 2016.
- 318 ²⁶ Aaron A King, Sourya Shrestha, Eric T Harvill, and Ottar N Bjørnstad. Evolution of acute infections and
319 the invasion-persistence trade-off. The American Naturalist, 173(4):446–455, 2009.
- 320 ²⁷ Ian Sorrell, Andrew White, Amy B Pedersen, Rosemary S Hails, and Mike Boots. The evolution of
321 covert, silent infection as a parasite strategy. Proceedings of the Royal Society B: Biological Sciences,
322 276(1665):2217–2226, 2009.
- 323 ²⁸ Femke van den Berg, Nicolas Bacaer, Johan AJ Metz, Christian Lannou, and Frank van den Bosch. Periodic
324 host absence can select for higher or lower parasite transmission rates. Evolutionary Ecology, 25(1):121–137,
325 2011.
- 326 ²⁹ Ruairi Donnelly, Alex Best, Andrew White, and Mike Boots. Seasonality selects for more acutely virulent
327 parasites when virulence is density dependent. Proceedings of the Royal Society B: Biological Sciences,
328 280(1751):20122464, 2013.
- 329 ³⁰ W Baltensweiler, G Benz, P Bovey, and V Delucchi. Dynamics of larch bud moth populations. Annual
330 Review of Entomology, 22(1):79–100, 1977.
- 331 ³¹ SA Woods and JS Elkinton. Biomodal patterns of mortality from nuclear polyhedrosis virus in gypsy moth
332 (*lymantria dispar*) populations. Journal of Invertebrate Pathology, 50(2):151–157, 1987.
- 333 ³² Shan L Bilimoria. The biology of nuclear polyhedrosis viruses. Viruses of invertebrates, pages 1–72, 1991.
- 334 ³³ Greg Dwyer and Joseph S Elkinton. Using simple models to predict virus epizootics in gypsy moth
335 populations. Journal of Animal Ecology, pages 1–11, 1993.
- 336 ³⁴ Greg Dwyer. Density dependence and spatial structure in the dynamics of insect pathogens. The American
337 Naturalist, 143(4):533–562, 1994.
- 338 ³⁵ Robert W Campbell. The gypsy moth and its natural enemies. Number 381. US Department of Agriculture,
339 Forest Service, 1975.
- 340 ³⁶ V Delucchi. Parasitoids and hyperparasitoids of *zeiraphera diniana* [lep., tortricidae] and their pole in
341 population control in outbreak areas. Entomophaga, 27(1):77–92, 1982.
- 342 ³⁷ M Kenis and J Hilszczanski. Natural enemies of cerambycidae and buprestidae infesting living trees. In
343 Bark and wood boring insects in living trees in Europe, a synthesis, pages 475–498. Springer, 2007.

- 344 ³⁸ Roy Malcolm Anderson and Robert Mccredie May. The population dynamics of microparasites and
345 their invertebrate hosts. Philosophical Transactions of the Royal Society of London. B, Biological Sciences,
346 291(1054):451–524, 1981.
- 347 ³⁹ Thomas Caraco and Ing-Nang Wang. Free-living pathogens: life-history constraints and strain competition.
348 Journal of Theoretical Biology, 250(3):569–579, 2008.
- 349 ⁴⁰ Ing-Nang Wang. Lysis timing and bacteriophage fitness. Genetics, 172(1):17–26, 2006.
- 350 ⁴¹ Nicholas J White. Determinants of relapse periodicity in plasmodium vivax malaria. Malaria journal,
351 10(1):1–36, 2011.
- 352 ⁴² Johan AJ Metz, Roger M Nisbet, and Stefan AH Geritz. How should we define ‘fitness’ for general ecological
353 scenarios? Trends in ecology & evolution, 7(6):198–202, 1992.
- 354 ⁴³ Stefan AH Geritz, Géza Mesze, Johan AJ Metz, et al. Evolutionarily singular strategies and the adaptive
355 growth and branching of the evolutionary tree. Evolutionary ecology, 12(1):35–57, 1998.
- 356 ⁴⁴ Sébastien Lion and Johan AJ Metz. Beyond r_0 maximisation: on pathogen evolution and environmental
357 dimensions. Trends in ecology & evolution, 33(6):458–473, 2018.
- 358 ⁴⁵ BR Levin. Coevolution in bacteria and their viruses and plasmids. Coevolution, pages 99–127, 1983.
- 359 ⁴⁶ Dieter Ebert and Wolfgang W Weisser. Optimal killing for obligate killers: the evolution of life histories
360 and virulence of semelparous parasites. Proceedings of the Royal Society of London. Series B: Biological
361 Sciences, 264(1384):985–991, 1997.
- 362 ⁴⁷ A Sasaki and HCJ Godfray. A model for the coevolution of resistance and virulence in coupled
363 host–parasitoid interactions. Proceedings of the Royal Society of London. Series B: Biological Sciences,
364 266(1418):455–463, 1999.
- 365 ⁴⁸ Thomas Caraco and Nang Wang. Free-living pathogens: life-history constraints and strain competition.
366 Journal of theoretical biology, 250(3):569–579, 2008.
- 367 ⁴⁹ Knut Helge Jensen, Tom Little, Arne Skorping, and Dieter Ebert. Empirical support for optimal virulence
368 in a castrating parasite. PLoS Biol, 4(7):e197, 2006.
- 369 ⁵⁰ Frida Ben-Ami. The virulence–transmission relationship in an obligate killer holds under diverse
370 epidemiological and ecological conditions, but where is the tradeoff? Ecology and evolution, 7(24):11157–
371 11166, 2017.
- 372 ⁵¹ Sarah E Reece, Kimberley F Prior, and Nicole Mideo. The life and times of parasites: rhythms in strategies
373 for within-host survival and between-host transmission. Journal of biological rhythms, 32(6):516–533,
374 2017.
- 375 ⁵² RAI Norval, JA Lawrence, AS Young, Brian D Perry, TT Dolan, and J Scott. Theileria parva: influence of
376 vector, parasite and host relationships on the epidemiology of theileriosis in southern africa. Parasitology,
377 102(3):347–356, 1991.
- 378 ⁵³ S E Randolph. Tick ecology: processes and patterns behind the epidemiological risk posed by ixodid ticks
379 as vectors. Parasitology, 129(7):S37–S65, 1999.

- 380 ⁵⁴ H Ochanda, AS Young, C Wells, GF Medley, and Brian D Perry. Comparison of the transmission of theileria
381 parva between different instars of rhipicephalus appendiculatus. Parasitology, 113(3):243–253, 1996.
- 382 ⁵⁵ Elodie Gaulin, Christophe Jacquet, Arnaud Bottin, and Bernard Dumas. Root rot disease of legumes caused
383 by aphanomyces euteiches. Molecular Plant Pathology, 8(5):539–548, 2007.
- 384 ⁵⁶ Eldon I Zehr et al. Control of brown rot in peach orchards. Plant disease, 66(12):1101–1105, 1982.
- 385 ⁵⁷ Ivan H Crowell. The hosts, life history and control of the cedar-apple rust fungus gymnosporangium
386 juniperi-virginianae schw. Journal of the Arnold Arboretum, 15(3):163–232, 1934.
- 387 ⁵⁸ J Holuša and K Lukášová. Pathogen's level and parasitism rate in ips typographus at high population
388 densities: importance of time. Journal of Applied Entomology, 141(9):768–779, 2017.
- 389 ⁵⁹ Michael Boots and Akira Sasaki. 'small worlds' and the evolution of virulence: infection occurs locally and at
390 a distance. Proceedings of the Royal Society of London. Series B: Biological Sciences, 266(1432):1933–1938,
391 1999.
- 392 ⁶⁰ Benjamin Kerr, Claudia Neuhauser, Brendan J M Bohannan, and Antony M Dean. Local migration promotes
393 competitive restraint in a host–pathogen 'tragedy of the commons'. Nature, 442(7098):75–78, 2006.

Appendix A

In Appendix A we find analytical solutions for equations 1a-c from the main text to study parasite fitness given different host phenological patterns.

$$\frac{ds}{dt} = \hat{s}g(t, t_l) - ds(t) - \alpha s(t)v_1(t), \quad (\text{A.1a})$$

$$\frac{dv_1}{dt} = -\delta v_1(t), \quad (\text{A.1b})$$

$$\frac{dv_2}{dt} = \alpha\beta e^{-d\tau} s(t-\tau)v_1(t-\tau) - \delta v_2(t). \quad (\text{A.1c})$$

with initial conditions: $s(0) = 0, v_1(0^+) = v_2(0^-), v_2(\tau) = 0$.

(A.1a-c) is solved analytically by describing host emergence using a uniform distribution

$$g(t, t_l) = \begin{cases} \frac{1}{t_l} & 0 \leq t \leq t_l \\ 0 & t_l < t \end{cases}$$

To solve the dynamics during the host's activity period, we first find the analytical solution for $v_1(t)$:

$$v_1(t) = \hat{v}e^{-\delta t}$$

We then use $v_1(t)$ to find the time-dependent solutions for both $s(t)$ and $v_2(t)$:

$$s(t) = \begin{cases} \frac{\hat{s}}{t_l} e^{(-dt + \frac{\alpha\hat{v}e^{-\delta t}}{\delta})} \int_0^t e^{(ds - \frac{\alpha\hat{v}e^{-\delta s}}{\delta})} ds & 0 < t < t_l \\ s(t_l) e^{(-d(t-t_l) - \frac{\alpha\hat{v}e^{-\delta t}(1-e^{\delta(t-t_l)})}{\delta})} & t_l \leq t < T \end{cases}$$

$$v_2(t) = \begin{cases} \frac{\alpha\beta e^{-d\tau} \hat{v} \hat{s}}{t_l} e^{-\delta t} \int_0^t e^{(-ds + \frac{\alpha\hat{v}e^{-\delta s}}{\delta})} \int_0^s e^{(dx - \frac{\alpha\hat{v}e^{-\delta x}}{\delta})} dx ds & \tau < t < t_l \\ e^{-\delta(t-t_l-\tau)} (v_2(t_l) + \alpha\beta e^{-d\tau} \hat{v} s(t_l) \int_0^{t-t_l-\tau} e^{-\frac{\alpha\hat{v}e^{-\delta(s+t_l)(-1+e^{\delta s})}{\delta} - \delta t_l - ds} ds) & t_l \leq t < T \end{cases}$$

where $s(t_l)$ and $v_2(t_l)$ are the densities of s and v_2 when the emergence period of s ends.

A parasite introduced into a naive host population persists or goes extinct depending on the host emergence period length and season length. The stability of the parasite-free equilibrium is determined by the production of v_2 resulting from infection of s given by

$$v_2(T) = e^{-\delta(T-t_l-\tau)} (v_2(t_l) + \alpha\beta e^{-d\tau} \hat{v} s(t_l) \int_0^{T-t_l-\tau} e^{-\frac{\alpha\hat{v}e^{-\delta(s+t_l)(-1+e^{\delta s})}{\delta} - \delta t_l - ds} ds)$$

The parasite-free equilibrium is unstable and the parasite will persist in the system if the density of v_2 produced by time T is greater than or equal to $\hat{v} = v_1(0) = 1$ introduced at the beginning of the activity period of s (*i.e.* $v_2(T) \geq 1$, modulus is greater than unity). This expression is a measure of a parasite's fitness when rare given different host phenological patterns.

407 Appendix B

408 In Appendix B we find analytical solutions for equations 2a-e from the main text to study the evolution of
409 parasite virulence given different host phenological patterns.

$$\frac{ds}{dt} = \hat{s}g(t, t_l) - ds(t) - \alpha s(t)v_1(t) - \alpha_m s(t)v_{1m}(t), \quad (\text{B.1.a})$$

$$\frac{dv_{1m}}{dt} = -\delta_m v_{1m}(t), \quad (\text{B.1.b})$$

$$\frac{dv_{2m}}{dt} = \alpha_m \beta_m s(t - \tau_m)v_{1m}(t - \tau_m) - \delta_m v_{2m}(t). \quad (\text{B.1.c})$$

$$\frac{dv_1}{dt} = -\delta v_1(t), \quad (\text{B.1.d})$$

$$\frac{dv_2}{dt} = \alpha \beta s(t - \tau)v_1(t - \tau) - \delta v_2(t), \quad (\text{B.1.e})$$

410 with initial conditions: $s(0) = 0, v_{1m}(0^+) = v_{2m}(0^-), v_{2m}(\tau) = 0, v_1(0^+) = v_2(0^-), v_2(\tau) = 0$. m subscripts
411 refer to the invading mutant parasite and its corresponding traits.

412

(B.1.a-c) has the following time-dependent solution:

$$\begin{aligned} v_{1m}(t) &= v_{1m}(0)e^{-\delta_m t} && 0 < t < T \\ s_1(t) &= \begin{cases} \frac{\hat{s}_1}{t_l} e^{(-dt + \frac{\alpha \hat{v}^* e^{-\delta t}}{\delta} + \frac{\alpha_m v_{1m}(0) e^{-\delta_m t}}{\delta_m})} \int_0^t e^{(ds - \frac{\alpha \hat{v}^* e^{-\delta s}}{\delta} + \frac{\alpha_m v_{1m}(0) e^{-\delta_m s}}{\delta_m})} ds & 0 < t < t_l \\ s(t_l) e^{(-d(t-t_l) - (\frac{\alpha \hat{v}^* e^{-\delta(t-t_l)}(1-e^{\delta t})}{\delta} + \frac{\alpha_m v_{1m}(0) e^{-\delta_m(t-t_l)}(1-e^{\delta_m t})}{\delta_m})} & t_l \leq t < T \end{cases} \\ v_{2m}(t) &= \begin{cases} \frac{\alpha_m \beta_m e^{-d\tau_m} v_{1m}(0) \hat{s}}{t_l} e^{-\delta_m t} \int_0^t e^{(-ds + \frac{\alpha \hat{v}^* e^{-\delta s}}{\delta} + \frac{\alpha_m v_{1m}(0) e^{-\delta_m s}}{\delta_m})} \int_0^s e^{(dx + \frac{\alpha \hat{v}^* e^{-\delta x}}{\delta} + \frac{\alpha_m v_{1m}(0) e^{-\delta_m x}}{\delta_m})} dx ds & \tau_m < t < t_l \\ e^{-\delta_m(t-t_l-\tau_m)}(v_2(t_l) + \alpha_m \beta_m e^{-d\tau_m} v_{1m}(0) s_1(t_l) \\ \int_0^{t-t_l-\tau_m} e^{-\frac{\alpha_m v_{1m}(0) e^{-\delta_m(s+t_l)}(-1+e^{-\delta_m s})}{\delta_m} - \frac{\alpha \hat{v}^* e^{-\delta(s+t_l)}(-1+e^{-\delta s})}{\delta} - \delta_m t_l - ds} ds) & t_l \leq t \leq T \end{cases} \end{aligned}$$

413 The invasion fitness of a rare mutant parasite is given by the density of v_{2m} produced by the end of the season.

414 The mutant parasite invades in a given host phenological scenario if the density of v_{2m} produced by time T

415 is greater than or equal to the initial $v_{1m}(0) = 1$ introduced at the start of the season ($v_{2m}(T) \geq 1$).

$$\begin{aligned} v_{2m}(T) &= e^{-\delta_m(T-t_l-\tau_m)}(v_2(t_l) + \alpha_m \beta_m e^{-d\tau_m} v_{1m}(0) s_1(t_l) \\ &\quad \int_0^{T-t_l-\tau_m} e^{-\frac{\alpha_m v_{1m}(0) e^{-\delta_m(s+t_l)}(-1+e^{-\delta_m s})}{\delta_m} - \frac{\alpha \hat{v}^* e^{-\delta(s+t_l)}(-1+e^{-\delta s})}{\delta} - \delta_m t_l - ds} ds) \end{aligned}$$

416 We use $v_{2m}(T)$ to find optimal virulence for a given host phenological scenario by finding the trait value that

417 maximizes $v_{2m}(T)$. That is, the virulence trait, τ^* , that satisfies

$$\frac{dv_{2m}(T)}{d\tau_m} \Big|_{\tau_m=\tau_r} = 0 \quad (\text{B.2})$$

$$\frac{d^2v_{2m}(T)}{d\tau_m^2} \Big|_{\tau_m=\tau_r} < 0 \quad (\text{B.3})$$

418 We use (B.2) to find τ^* in Figures 1A, 2A, 3A and 3B in the main text. For all phenological patterns, τ^* is
 419 uninvadable *i.e.* condition (B.3) is satisfied.

420 To study the impact of mechanistic trade-offs between transmission and virulence on virulence evolution,
 421 we assume that the number of parasites produced at host death is a function of the time between infection
 422 and host death ($\beta(\tau)$). This is done in Figures 3A and 3B where a mechanistic trade-off is assumed to exist
 423 between τ and β in (B.2).