

1 **Deceptive combined effects of short allele dominance and stuttering: an example**
2 **with *Ixodes scapularis*, the main vector of Lyme disease in the U.S.A.**

3
4 Thierry De Meeûs^{1,*}, Cynthia T. Chan^{2,3}, John M. Ludwig^{2,4}, Jean I. Tsao⁵, Jaymin Patel^{2,6},
5 Jigar Bhagatwala^{2,7}, and Lorenza Beati².

6
7 1. Intertryp, IRD, Cirad, Univ Montpellier, Montpellier, France.

8 2. The U.S. National Tick Collection, Institute for Coastal Plain Science, Georgia Southern
9 University, Statesboro, GA, USA.

10 3. College of Agricultural & Environmental Sciences, University of Georgia, Athens, GA,
11 USA.

12 4. Department of Microbiology, University of Georgia, Athens, GA, USA

13 5. Department of Fisheries and Wildlife, Michigan State University, East Lansing, MI, USA

14 6. Division of Hospital Medicine, College of Medicine, University of Florida, Gainesville, FL,
15 USA.

16 7. Medical College of Georgia, Augusta University, Augusta, GA, USA.

17
18 * Corresponding author: thierry.demeeus@ird.fr

19
20 Keywords: Microsatellite loci, short allele dominance, stuttering, heterozygote deficit,
21 linkage disequilibrium, curing microsatellite data.

22
23 Running title: Short allele dominance and stuttering
24
25
26

27 **Abstract**

28 Null alleles, short allele dominance (SAD), and stuttering increase the perceived
29 relative inbreeding of individuals and subpopulations as measured by Wright's F_{IS} and F_{ST} .
30 **Ascertainment bias, due to** such amplifying problems are usually caused by inaccurate
31 primer design (if developed from a different species or a distant population), poor DNA
32 quality, low DNA concentration, or a combination of some or all these sources of
33 inaccuracy. When combined, these issues can increase the correlation between
34 polymorphism at concerned loci and, consequently, of linkage disequilibrium (LD) between
35 those. In this note, we studied an original microsatellite data set generated by analyzing
36 nine loci in *Ixodes scapularis* ticks from the eastern U.S.A. To detect null alleles and SAD
37 we used correlation methods and variation measures. To detect stuttering, we evaluated
38 heterozygote deficit between alleles displaying a single repeat difference. We
39 demonstrated that an important proportion of loci affected by amplification problems (one
40 with null alleles, two with SAD and three with stuttering) lead to highly significant
41 heterozygote deficits ($F_{IS}=0.1$, p -value<0.0001). This occurred together with a prohibitive
42 proportion (22%) of pairs of loci in significant LD, **two of which were still significant after a**
43 **false discovery rate (FDR) correction, and some** variation in the **measurement** of
44 population subdivision across loci (Wright's F_{ST}). This suggested a strong Wahlund effect
45 and/or **selection** at several loci. By finding small peaks corresponding to previously
46 disregarded larger alleles in some homozygous profiles for loci with SAD and by pooling
47 alleles close in size for loci with stuttering, we generated an amended dataset. Except for
48 one locus with null alleles **and another still displaying a modest SAD**, the analyses of the
49 corrected dataset revealed a significant excess of heterozygotes ($F_{IS}=-0.07$ as expected in
50 dioecious and strongly subdivided populations, with a more reasonable proportion **(19%)**
51 **of pairs of loci characterized by significant LD, none of which stayed significant after the**
52 **FDR procedure**. Strong subdivision was also confirmed by the standardized F_{ST}' corrected
53 for null alleles ($F_{ST}'=0.19$) **and small effective subpopulation sizes ($N_e=7$)**.

54

55

56 Introduction

57 Null alleles, short allele dominance (SAD) and stuttering are frequent consequences
58 of poor PCR amplifications, in particular for microsatellite markers. Amplification problems
59 usually arise when primers are designed by using DNA of a different species or a distant
60 population, when DNA is degraded, at too low of a concentration (Chapuis and Estoup,
61 2007), or any combination of these listed causes.

62 Null alleles occur when a mutation on the flanking sequence of the targeted locus
63 affects the hybridization of the corresponding primer, resulting in amplification failure.
64 Heterozygous individuals with one null allele falsely appear to be homozygous, while
65 homozygous individuals for null alleles are considered to be missing data.

66 SAD, also called large allele dropout (Van Oosterhout et al., 2004), known from
67 minisatellite markers, was also discovered to occur in microsatellite loci (Wattier et al.,
68 1998). In heterozygous samples, longer alleles are less successfully amplified than shorter
69 alleles through competition for Taq polymerase. This can lead to misinterpreting
70 heterozygous individuals as homozygous for the shortest allele (De Meeûs et al., 2007).

71 Stuttering is the result of inaccurate PCR amplification through Taq slippage of a
72 specific DNA strand. This generates several PCR products that differ from each other by
73 one repeat and can cause difficulties when discriminating between fake and true
74 homozygotes, such as heterozygous individuals for dinucleotide microsatellite allele
75 sequences with a single repeat difference.

76 Allelic dropout is akin to SAD, but occurs randomly to any allele irrespective of its
77 size, and can affect both alleles of heterozygous individuals.

78 The consequence of these issues is a homozygous excess when compared to the
79 expected Castle-Weinberg proportions (Castle, 1903; Weinberg, 1908) (classically known
80 as Hardy-Weinberg, however, more accurately depicted as Castle-Weinberg; because the
81 former was the first to derive it for two equipotent alleles in 1903 and the latter
82 generalized the concept in January 1908, prior to Hardy in April 1908 (Hardy, 1908)), as
83 measured by Wright's F_{IS} (Wright, 1965). These problems, like all others associated with
84 amplification, are locus specific (Van Oosterhout et al., 2004; De Meeûs et al., 2007; De
85 Meeûs, 2018a) and thus lead to locus specific variation (namely, an increase) of F_{IS} . A less
86 well known, though well documented (Chapuis and Estoup, 2007; Séré et al., 2017b;
87 Manangwa et al., 2019) effect of such amplification problems consists of an increase of
88 Wright's F_{ST} (Wright, 1965) that is commonly used to measure the degree of genetic
89 differentiation between subpopulations.

90 While an analytical cure exists for null alleles (Chapuis and Estoup, 2007; Séré et
91 al., 2017a), such remediation is unavailable for SAD and stuttering. To the best of our
92 knowledge, the impact of amplification problems on linkage disequilibrium (LD) between
93 locus pairs has yet to be investigated. Problems with amplification can be expected to
94 more commonly occur in individuals that display some kind of deviating DNA in terms of
95 quantity or quality: flanking sequences that have accumulated mutations, samples
96 containing weak DNA concentration, badly preserved DNA extracts, or a combination of
97 these different problems. When combined, the effect of the occurrence of null alleles, SAD,
98 and stuttering may artificially generate a positive correlation between allele occurrences at
99 affected loci and then increase the perceived LD between them.

100 In this note, we utilize an original data set generated through the analysis of nine
101 microsatellite loci in *Ixodes scapularis*, sampled across the eastern U.S. to show that the
102 combined effect of SAD, stuttering, and null alleles can induce an increase in the number
103 of locus pairs in significant LD. We then propose and test an efficient way to amend such
104 data.

105

106 **Material and Methods**

107 *Sampling and DNA extraction*

108 Larvae, nymphs, and adults of *I. scapularis* were sampled indiscriminately from
109 different sites across the eastern U.S. on different occasions, extending from November
110 2001 to May 2014, by means of dragging and flagging the vegetation (Figure 1 and Table
111 1) (Rulison et al., 2013).

112 Gravid females fall on the ground where they lay thousands of eggs at the same
113 place that hatch as weakly mobile larvae. Larval collections can thus be composed of
114 clusters of thousands of sisters and brothers within the same subsample (Kempf et al.,
115 2011). Consequently, to avoid possible Wahlund effects, where a heterozygote deficit
116 results from the admixture of individuals from genetically distant subpopulations (e.g. see
117 (De Meeûs, 2018a)) (here families), we removed immature stages from the present study.
118 The remaining 387 adult ticks were subdivided into cohorts, with each cohort comprised
119 of samples collected across two consecutive years in the fall, the following winter and spring
120 across the tick distribution range. This subdivision was based on observations showing
121 that northeastern adults active in Fall can undergo winter quiescence and resume activity
122 in spring (Yuval and Spielman, 1990).

123 Many publications have emphasized the importance of mitochondrial clades in
124 different populations of *I. scapularis* across the U.S. (Norris et al., 1996; Qiu et al., 2002;

125 Sakamoto et al., 2014). Thus, to account for the mitochondrial clade representation and to
126 (again) avoid possible Wahlund effects, all ticks were assigned clades by phylogenetic
127 analysis of their 12S rDNA gene sequences. We identified 6 main clades in our dataset,
128 the previously identified American clade was subdivided in two lineages (AMI and AMII),
129 and the so-called southern clade was subdivided in 4 lineages (SOI, SOII, SOIII and SOIV)
130 (Table 1).

131 In conclusion, the combination of Site-Clade-Cohort data defined 61 subsamples
132 within the 387 individual adult ticks. Overall, 35 subsamples included a small number of
133 individuals (1-4), 12 subsamples contained at least 10 individuals and 5 subsamples
134 contained at least 20 individuals (Table 1). Because the smallest subsamples were
135 expected to exert a negligible weight on our analyses, they were not eliminated.

136 Procedures for all DNA extractions followed modified published protocols (Beati and
137 Keirans, 2001) with a DNeasy Blood & Tissue Kit (Qiagen, Valencia, CA).

138

139 *Selection and characterization of microsatellite markers*


140 Thirteen of the first batches of genome sequences of *I. scapularis* that were
141 accessioned by VectorBase (www.vectorbase.org; Giraldo-Calderón *et al.* 2015) in
142 GenBank (AC205653.1, AC205652.1, AC205650.1, AC205647.1, AC205646.1,
143 AC205643.1, AC205642.1, AC205641.1, AC205638.1, AC205635.1, AC205634.1,
144 AC205632.1, AC205630.1) were used to manually detect motifs with at least 6 repeats of
145 AG, AT, CA, TA, TG, CT, GC, ACG, GTT, TTA, CAC, GAT, and AAAC. Primer pairs were
146 selected in the flanking regions by using Oligo v.5 (Molecular Biology Insights, Colorado
147 Springs, CO). DNA, extracted and pooled from six ticks from Connecticut, was used to test
148 whether the selected primer sets successfully amplified fragments of the expected size.
149 PCRs were performed using the 5-Prime Master PCR kit (5-Prime, Gaithersburg, MD) and
150 a single touch-down amplification protocol consisting of 5 min. denaturation at 93°C; 5
151 cycles: 20 sec. denaturation at 93°C, 20 sec. annealing at 55°C-1.5°C/cycle, 30 sec.
152 elongation at 72°C; 30 cycles: 20 sec. denaturation at 93°C, 25 sec. annealing at 47°C, 30
153 sec. elongation at 70°C; final extension at 70°C for 5 min. Amplicons were run on 4% E-
154 gels (Life Technologies Co., Carlsbad, CA). The risk of having selected primers within
155 repeated portions of the genome had to be considered due to the fact that large repeated
156 genomic fragments are known to occur abundantly in *I. scapularis* (Gulia-Nuss et al.,
157 2016). In order to confirm that the primers were amplifying the targeted loci, the amplicon
158 of one randomly chosen tick was cloned with a TOPO-TA PCR cloning kit (Life
159 Technologies Co, Carlsbad, CA) for each locus. Five cloned colonies were picked

160 randomly for each tick and the insert amplified and sequenced (DNA Analysis Facility on
161 Science Hill, Yale University). Finally, as microsatellite primers are known to amplify
162 sometimes more than one closely related species, the same set of primers was tested on
163 DNA samples of *Ixodes ricinus*, *Ixodes pacificus*, and *Ixodes persulcatus* (LB, personal
164 collection), all taxa belonging to the *I. ricinus* complex of ticks (Keirans et al., 1999).

165 The primer pairs that yielded amplicons of the expected size were then used to
166 individually amplify a subset of 67 DNA samples from ticks, representative of the
167 distribution **area of *I. scapularis* in USA, and** collected by flagging or dragging in Alabama
168 (10 **ticks**), Georgia (15), Connecticut (16), Massachusetts (14), New York (2),
169 Pennsylvania (2), and South Carolina (8). For these amplifications, forward primers were
170 labeled with fluorescent dyes (Applied Biosystems, Thermo Fisher Scientific, CA) as listed
171 in Table 2. The amplicons were sent to the DNA Analysis Facility on Science Hill (Yale
172 University, New Haven, CT) for genotyping. The allele peaks were scored using
173 GeneMarker (SoftGenetics, State College, PA). All data were recorded in an Excel
174 spreadsheet for further ease of conversion.

175

176 Figure 1: Sampling sites for *Ixodes scapularis* from the eastern U.S.A. (State codes as in
177 Table 1).



178

179

180

181
182
183

Table 1: State, site, GPS coordinates (decimal degrees), 12S clade membership, date of sampling, corresponding cohort membership and size of *Ixodes scapularis* adult subsamples (N) from the eastern U.S.A.

State	Site	Latitude	Longitude	Clade	Date	Cohort	N
Alabama	AL1	33.24	-86.13	AMI	2011 Jan	C7	21
	AL1	33.24	-86.13	SOI	2011 Jan	C7	6
	AL1	33.24	-86.13	AMII	2011 Jan	C7	18
	AL1	33.24	-86.13	SOII	2011 Jan	C7	6
	AL2	32.95	-87.14	AMI	2012 Dec	C9	3
	AL2	32.95	-87.14	AMII	2012 Dec	C9	2
Connecticut	CT1	41.35	-72.76	AMI	2001 Nov	C1	3
	CT1	41.35	-72.76	AMI	2003 Jun	C2	21
Florida	FL1	30.65	-84.21	SOII	2012 Dec	C9	4
	FL2	30.06	-81.37	AMI	2011 Jan	C7	2
	FL2	30.06	-81.37	AMII	2011 Jan	C7	3
	FL2	30.06	-81.37	SOI	2011 Jan	C7	1
	FL2	30.06	-81.37	SOII	2011 Jan	C7	4
Georgia	GA1	32.45	-81.78	AMI	2009 Dec	C5	11
	GA1	32.45	-81.78	SOI	2009 Dec	C5	5
	GA1	32.45	-81.78	SOII	2009 Dec	C5	17
	GA1	32.45	-81.78	SOIII	2009 Dec	C5	1
Iowa	IA1	42.67	-91.59	AMI	2007 May	C3	3
Louisiana	LA1	30.94	-91.46	AMI	2012 Dec	C9	3
	LA1	30.94	-91.46	AMII	2012 Dec	C9	2
	LA1	30.94	-91.46	SOIII	2012 Dec	C9	1
	LA1	30.94	-91.46	SOIV	2012 Dec	C9	2
Massachusetts	MA1	41.71	-69.92	AMI	2010 May	C5	6
Maryland	MD1	39.29	-76.88	AMI	2010 Oct	C7	11
North-Carolina	NC1	33.91	-78.39	AMI	2009 Jan	C4	22
	NC1	33.91	-78.39	SOI	2009 Jan	C4	4
	NC1	33.91	-78.39	SOIII	2009 Jan	C4	2
New-York	NY1	40.76	-72.83	AMI	2009 Oct	C5	10
Pennsylvania	PA1	40.67	-75.96	AMI	2010 Oct	C7	5
	PA2	41.06	-79.48	AMI	2014 May	C10	33
South-Carolina	SC1	33.33	-81.66	AMI	2011 Apr	C7	8
	SC1	33.33	-81.66	AMII	2011 Apr	C7	1
	SC1	33.33	-81.66	SOI	2011 Apr	C7	3
	SC1	33.33	-81.66	SOII	2011 Apr	C7	1
	SC1	33.33	-81.66	SOIII	2011 Apr	C7	1
	SC2	33.23	-81.73	AMI	2010 Dec	C7	4
	SC2	33.23	-81.73	AMII	2010 Dec	C7	8
	SC2	33.23	-81.73	SOI	2010 Dec	C7	4
	SC2	33.23	-81.73	SOII	2010 Dec	C7	10
	SC3	33.15	-81.61	SOII	2010 Dec	C7	2
	SC5	34.29	-79.87	AMI	2012 Dec	C9	3
	SC5	34.29	-79.87	SOII	2012 Dec	C9	2
Tennessee	TN1	35.37	-86.07	AMI	2010 Dec	C7	23
Texas	TX1	31.80	-96.23	AMI	2012 Dec	C9	5
	TX1	31.80	-96.23	SOI	2012 Dec	C9	2
	TX1	31.80	-96.23	SOIV	2012 Dec	C9	3
	TX2	30.24	-100.72	AMI	2012 Dec	C9	1
	TX2	30.24	-100.72	SOIV	2012 Dec	C9	5
	TX3	31.91	-95.9	AMI	2012 Dec	C9	3
	TX3	31.91	-95.9	SOI	2012 Dec	C9	2
	TX3	31.91	-95.9	SOIV	2012 Dec	C9	6
	TX4	31.59	-95.61	AMI	2012 Dec	C9	2
	TX4	31.59	-95.61	SOI	2012 Dec	C9	2
Virginia	VA1	38.29	-78.29	AMI	2010 Oct	C7	7
	VA2	37.32	-80.73	AMI	2012 Dec	C9	4
Wisconsin	WI1	43.95	-90.70	AMI	2011 Oct	C8	8
	WI1	43.95	-90.70	AMI	2012 Oct	C9	4
	WI2	44.04	-90.65	AMI	2011 Oct	C8	8
	WI2	44.04	-90.65	AMI	2012 Oct	C9	12
	WI3	44.02	-90.64	AMI	2011 Oct	C8	8
WI3	44.02	-90.64	AMI	2012 Oct	C9	3	

184

Table 1: List of primer sets used or developed for this study. PAC = *Ixodes pacificus*, PER = *I. persulcatus*, RIC = *I. ricinus*.

Locus	Repeat	Primer name	Primer sequences (5'→3')	Ref	Dye	Size)	Nal	Sp
IS1	(AG) ₁₀	Amy1-IsAG25a Amy2-IsAG25b	AAATGTCCGAACAGCCTTAT GCCCTTGAGTCTACCCACTA	Fagerberg <i>et al</i> (2001)	6 FAM	93-193	17	PAC/PER/RIC
IS3	(GTT) ₅	bac1d_a bac1d_b	GCAGATCTCTTTGGGCTAG AAGCTAAGGCGTTTCGTTG	AC205653	VIC	76-100	7	none
IS4	(AT) ₂₁	bac1m_a bac1m_b	TGTCGGTTTGATGCCAA GGCTCCATTCACCAGTC	AC205653	VIC	88-126	17	PAC/PER/RIC
IS5	(CA) ₉	bac3dh_a bac3dh_b	TGCCTGTGACGAAACCA TCTCCAAGAGATCTAGGTA	AC205650	NED	62-140	17	none
IS6	(TA) ₁₀	bac1j_a bac1j_b	TCTCCAAGAGATCTAGGTA ATCTGTTTCAGTGGGCACA	AC205653	VIC	100-186	13	PER
IS7	(TA) ₁₁	bac1k_a bac1k_b	GGGACTGGACACACGA CTAGGTGGCGCAAGTC	AC205653	VIC	48-170	26	none
IS8	(CA) ₁₄	bac3s_a bac3s_b	CGTTTCAAAGTCGGAGA GATGTGAGGGCGTGGT	AC205650	PET	96-194	11	PER
IS9	(AAAC) ₅	bac4cef_a bac4cef_b	CGCCTTTTGTCCCAACC GACTAACAGCATTGGAGCA	AC205647	6 FAM	85-125	12	PER
IS10	(TTA) ₉	bac5cf_a bac5cf_b	TCCCCAACCAAGATTGATG GAGACGACGTAGATTCTTG	AC205646	6 FAM	77-137	15	none
IS11	(TTA) ₆	bac5g_a bac5g_b	GCTTTAGCGGGCTGGT TACGTGAATACGTCCTTGG	AC205646	PET	81-165	12	PER
IS12	(TA) ₄₃	bac6a_a bac6a_b	GCAAGCTTCGCTATTCTC CAGTAATTTCCGATCGGTT	AC205643	6 FAM	111-229	26	none
IS13	(TA) ₂₂	bac6c_a bac6c_b	TAGGTACAAGAAAACGTGCT CAAGGTAATTGTTCTCGTCA	AC205643	NED	37-91	17	none
IS14	(TA) ₅	bac6d_a bac6d_b	CCTTGCTTACATGGTT CGTACCAAACCAAAGCAAG	AC205643	HEX	57-105	13	PAC/PER/RIC
IS15	(AT) ₈	bac6e_a bac6e_b	TATTGTAACCGACGCTAGG GACAATCTCTACGCAAATCC	AC205643	NED	79-125	18	none
IS16	(CA) ₈	bac6f_a bac6f_b	CCCCAACACGCACA TTGCTTCATGCAGGGAAC	AC205643	VIC	80-106	12	RIC
IS17	(CA) ₆	bac7e_a bac7e_b	CCAGCATTTAACCCCTCAAG TAGTGGGGTATGGCACTG	AC205642	HEX	139-197	12	PER/RIC
IS18	(TG) ₆	bac8a_a bac8a_b	GTAGGTACCCTAAGAAGGAT TTGAGGAAGCAGAATGTAGG	AC205641	6 FAM	75-195	16	PER/RIC
IS19	(CT) ₇	bac9a_a bac9a_b	AGAACCAGTTCAGCATTCC GAACATTTTCACGTTTGC	AC205638	PET	94-166	6	PER
IS20	(GC) ₉	bac11a_a bac11a_b	CGCTCCCTTCGAAGTTC GAGAAGACAGTTTCCATCG	This study	HEX	76-106	13	PAC/PER/RIC
IS21	(ACG) ₆	bac11c_a bac11c_b	CGAATCGCGCACACTAG GCTGTGTTGCTGGTCC	This study	NED	109-251	14	PAC/PER
IR27	(AC) ₉		ATACCCGTAGAACGAGAG GTTTTCAAGATTTCCGCC	Delaye <i>et al</i> (2008)	6 FAM			RIC

Ref: Reference or GenBank accession number; Size: Approximate size range (bp, 67 individuals); Nal: Number of alleles (67 individuals); Sp: Cross-species amplification

188 *Genotyping*

189 Based on their degree of polymorphism, nine microsatellite loci (IR27, IS1, IS3,
190 IS11, IS15, IS16, IS17, IS18, and IS19) were used for genotyping at the continental scale
191 (Table 2). Of these, IR27 (Delaye et al., 1998) and IS1 (Fagerberg et al., 2001), were
192 drawn from previously published studies. The loci were amplified and genotyped using the
193 procedures described above, although PCR conditions had to be slightly optimized for
194 markers IS11 and IS15 (touchdown annealing temperature decreased from 58°C to 50°C)
195 and IR27 (touchdown annealing temperature decreased from 56°C to 53°C) (Table 2).

196

197 *Population genetics analyses*

198 The raw data set was coded and converted into all required formats using Create
199 (Coombs et al., 2008).

200 To test for LD, we used the *G*-based test first described by Goudet et al. (Goudet et
201 al., 1996) and adapted for contingency tables of locus pairs, with 15000 reshuffling of
202 genotypes to get maximum precision and minimize possible *p*-values before false
203 discovery rate corrections (see below). The *G* statistics obtained for each subsample were
204 then summed over all subsamples to get a single statistic and hence, a single test across
205 subsamples. This procedure was shown to be the most powerful (De Meeûs et al., 2009)
206 and was implemented within Fstat 2.9.4 (Goudet, 2003) an updated version of the original
207 1995 Fstat software (Goudet, 1995). There are as many tests as locus pairs and these
208 tests are correlated (one locus is used as many times as there is any other locus). To take
209 into account this repetition of correlated tests, we used Benjamini and Yekutieli (BY) false
210 discovery rate procedure (Benjamini and Yekutieli, 2001) with R version 3.5.1 (R-Core-
211 Team, 2018). To check if some loci were involved in a significant LD pair more often than
212 by chance, as compared to the other loci, we also undertook a Fisher exact test with
213 Rcmdr version 2.3-1 (Fox, 2005; Fox, 2007).

214 For a hierarchy with three levels (individuals, subsamples, and total sample), three
215 *F*-statistics can be defined (Wright, 1965). *F_{IS}* measures inbreeding of individuals relative
216 to inbreeding of subsamples or relative deviation of observed genotypic proportions from
217 local random mating proportions. *F_{ST}* measures inbreeding of subsamples relative to total
218 inbreeding or relative inbreeding due to the subdivision of the total population into several
219 isolated subpopulations. *F_{IT}* measures inbreeding of individuals relative to total inbreeding.
220 Under the null hypothesis (panmixia and no subdivision), all these statistics are expected
221 to be null. Otherwise, *F_{IS}* and *F_{IT}* can vary from -1 (one heterozygote class) to +1 (all
222 individuals homozygous) and *F_{ST}* from 0 (all subsamples share similar allele frequencies)

223 to +1 (all subsamples fixed for one or the other allele). These statistics were estimated with
224 Weir and Cockerham's unbiased estimators (Weir and Cockerham, 1984) with Fstat.

225 In dioecious species (like ticks), heterozygote excess occurs over all loci (e.g. (De
226 Meeûs et al. 2007)) and is proportional to subpopulation size ($N_e = -1/(2 \times F_{IS}) - F_{IS}/(1 + F_{IS})$)
227 (Balloux, 2004). Therefore, the finding of homozygous excesses really represents a strong
228 deviation from random mating expectations. Technical problems, like null alleles,
229 stuttering, SAD or allele dropouts unevenly affects some loci, producing a positive F_{IS} with
230 an important variation across loci and significant outliers (De Meeûs 2018). Significant
231 departure from 0 of these F -statistics was tested with 10000 randomizations of alleles
232 between individuals within subsample (deviation from local random mating test) or of
233 individuals between subsamples within the total sample (population subdivision test). For
234 F_{IS} , the statistic used was f (Weir and Cockerham's F_{IS} estimator). To test for subdivision,
235 we used the G -based test (Goudet et al. 1996) over all loci, which is the most powerful
236 procedure when combining tests across loci (De Meeûs et al. 2009).

237 To compute 95% confidence intervals (95%CI) of F -statistics, we used the standard
238 error of F_{IS} (StrdErrFIS) and F_{ST} (StrdErrFST) computed by jackknifing over populations, or
239 5000 bootstraps over loci as described elsewhere (De Meeûs et al. 2007). For jackknives,
240 the number of usable subsamples was restricted to subsamples with at least 5 ticks (23
241 subsamples) (e.g. (De Meeûs, 2012) p 73).

242 In case of significant homozygote excess and LD we tried to discriminate
243 demographic from technical causes with the determination key proposed by De Meeûs (De
244 Meeûs 2018). Null alleles better explain the data if the StrdErrFIS becomes at least twice
245 as high as StrdErrFST; F_{IS} and F_{ST} are positively correlated; and a positive correlation
246 links F_{IS} and the number of missing data (putative null homozygotes). The significance of
247 correlations was tested with a unilateral ($\rho > 0$) Spearman's rank correlation test with R. The
248 presence of null alleles was also verified with MicroChecker v 2.2.3 (Van Oosterhout et al.
249 2004) and null allele frequencies estimated with Bookfield's second method (Bookfield
250 1996). The adjustment between observed and expected numbers of missing data was
251 tested with a unilateral exact binomial test in R with the alternative hypothesis being "there
252 is not enough missing data as expected if heterozygote deficits were entirely explained by
253 null alleles under panmixia". MicroChecker also seeks stuttering and SAD. Stuttering is
254 detected when MicroChecker reveals an observed proportion of heterozygous individuals
255 for alleles with one repeat difference significantly smaller than the expected value. The
256 presence of stuttering was detected with the graphic output of MicroChecker (we ignored
257 the comments panel that happened to contradict the graphic in some instances). We

258 considered that the observed deficit of heterozygous individuals for one repeat difference
259 was a likely consequence of stuttering. Due to the small population sample sizes and
260 bootstrapping procedure in MicroChecker, the statistical support (p -value) of this result
261 was not always significant for all runs. Hence, we set the randomization at the maximum
262 value (10000) and repeated the analysis three times to check for consistency. Stuttering
263 was admitted when two out of three tests supported it. The occurrence of SAD was also
264 checked with an unilateral ($\rho < 0$) Spearman's rank correlation test between allele size and
265 F_{IT} as proposed by Manangwa et al. (Manangwa et al., 2019). This test is more powerful
266 than with F_{IS} as was proposed earlier (Wattier et al., 1998; De Meeûs et al., 2004). If
267 previous tests are not significant and if $StrdErrFIS > StrdErrFST$, then a Wahlund effect
268 better explains the data (De Meeûs, 2018a), this is especially true in instances of
269 significant LD (Manangwa et al., 2019). In these cases, a positive correlation between the
270 number of times a locus is found in significant LD (NLD) and its total genetic diversity as
271 measured by Nei's H_T (Nei and Chesser, 1983) (Spearman's correlation above 0.1)
272 suggests an admixture of individuals from several subpopulations of the same species but
273 with an important degree of genetic differentiation between admixed subpopulations (i.e.
274 number of immigrants $N_e m = 2$). If the correlation is negative and the proportion of
275 significant LDs is above 40%, an admixture of different strongly divergent entities (e.g.
276 species) better explains the data (Manangwa et al., 2019). We thus undertook a bilateral
277 Spearman's test.

278 In some instances, repetition of independent tests were submitted to Benjamini and
279 Hochberg (BH) correction (Benjamini and Hochberg, 2000) (computed with R), to check for
280 robustness of significant p -values.

281 To our knowledge, when diagnosed, there is no analytical remedy for SAD or
282 stuttering as there is for null alleles (Chapuis and Estoup, 2007; Séré et al., 2017a).
283 Nevertheless, SAD can be cured by going back to the chromatograms of homozygous
284 individuals and trying to find a micro-peak (see the Results and discussion section), with a
285 larger size, ignored in the first reading. If enough profiles can be corrected this might
286 salvage the incriminated locus.

287 Stuttering can be addressed by pooling alleles close in size. However, this
288 procedure requires that none of the pooled allele groups is constituted of rare alleles only.
289 Indeed, pooling rare alleles, usually found in heterozygosity with a more frequent allele,
290 will tend to artificially generate heterozygous excesses. In order to avoid this
291 consequence, each pooled group should contain at least one frequent allele (e.g. with
292 $p \geq 0.05$).

293 In dioecious small populations, a heterozygote excess is expected. However, loci
294 with null alleles may display heterozygote deficits. In such situations a bilateral test (F_{IS} is
295 not different from 0) is needed and obtained as $p_{bilateral}=p_{\min}+1-p_{\max}$, where p_{\min} is the
296 minimum unilateral p -value (for heterozygote deficit or excess) and p_{\max} is the maximum
297 one.

298 Finally, a more accurate estimate of F_{ST} can be made for datasets with null alleles
299 after recoding missing genotypes as homozygous for allele 999 (null allele) with the ENA
300 method as implemented in FreeNA (Chapuis and Estoup, 2007). This estimate can then
301 be corrected for polymorphism with the formula $F_{ST}'=F_{ST}/(1-H_s)$ (Hedrick, 1999).

302

303 **Results and discussion**

304 *Primer selection and characterization of loci*

305 The inspection of the GenBank genomic sequences revealed the presence of 67
306 short tandem repeated motifs. The program Oligo v.5 did not find suitable primers for 17 of
307 them. Of the remaining 50 primer pairs, 22 amplified the pooled DNA sample and the sizes
308 of the amplicons were approximately as expected. The cloned amplified inserts all
309 contained the expected microsatellite repeats and flanking regions. The 22 primer sets
310 consistently amplified DNA from the 67 *I. scapularis* ticks and some of them also amplified
311 DNA of the other *Ixodes* species (Table 2). Finally, based on their polymorphism and ease
312 of interpretation, nine loci were retained for the population genetics analyses: IR27 (from
313 (Delaye et al., 1998)), IS1, IS3, IS11, IS15, IS16, IS17, IS18, and IS19 (from the present
314 study), with 9.8, 3.4, 1.3, 8, 8.3, 4.9, 2.8, 5.7 and 4.1 % missing genotypes (blanks)
315 respectively.

316

317 *Raw data analyses*


318 All data are available in the supplementary File S1.

319 There was a very important proportion of locus pairs in significant LD (~22%), with a
320 negative correlation between NLD and H_T ($\rho=-0.04$, p -value=0.9242). Two locus pairs
321 remained in significant LD after BY correction: IR27-IS3 (p -value=0.0301) and IS11-IS16
322 (p -value=0.0451). No single locus was found more often in significant LD than the others
323 (p -value=0.09).

324 There was a highly significant heterozygote deficit ($F_{IS}=0.111$, in 95%CI=[-0.02,
325 0.236, p -value<0.0002), with a considerable variation across loci (Figure 2).

326

327 Figure 2: F_{IS} values for each locus and averaged across those (All) of *Ixodes scapularis*
 328 from the eastern U.S.A. with 95% jackknife confidence intervals over subsamples
 329 (for each locus) and bootstraps over loci (All). Results of tests for panmixia, short
 330 allele dominance, number of subsamples with stuttering for each locus, and null
 331 allele frequencies are also indicated.



332
 333
 334 StdrdErrFIS (0.07) was almost four times higher than StrdErrFST (0.019); the
 335 correlation between F_{IS} and F_{ST} was negative ($\rho=-0.6166$, p -value=0.962) and the
 336 correlation between F_{IS} and the number of blanks (missing genotypes) was positive but not
 337 significant ($\rho=0.1833$, p -value=0.3218). These results suggested that locus-specific effects
 338 were involved. Nevertheless, null alleles only partly explained the observed pattern at best.
 339 The substantial proportion of significant LDs suggested the existence of a possible strong
 340 Wahlund effect, though a negative correlation between NLD and H_T with less than 40%
 341 significant tests observed here would refute this interpretation (Manangwa et al., 2019).


342 Small subsampling due to the partitioning of the data into cohorts and 12S clades
 343 was expected to considerably lower the power of these tests, especially so after the rather
 344 stringent BY procedure for the LD test series.

345 Subdivision was substantial ($F_{ST}=0.101$, 95%CI=[0.066, 0.136], p -value<0.0001),
 346 and variable across loci and subsamples (Figure 3), but not meaningfully more variable
 347 than expected (see Figure 6 in (De Meeûs, 2018b)). Some loci (IS1, IS15 and IS17)

348 displayed very low values (Figure 3). Selection might have produced the pattern observed,
 349 though evidence for this is weak.

350

351 Figure 3: F_{ST} values for each locus and averaged across those (All) of *Ixodes scapularis*
 352 from the eastern U.S.A. with 95% jackknife confidence intervals over subsamples
 353 (for each locus) and bootstrap over loci (All). Results of tests for significant
 354 subdivision (p -value) are also indicated.



355

356

357 Two loci displayed highly significant SAD (Figure 2): loci IS1 ($\rho=-0.57$, p -
 358 value=0.0015) and IS15 ($\rho=-0.46$, p -value=0.008), which stayed significant after BH
 359 correction (0.0135 and 0.036 respectively). Stuttering was diagnosed for four loci (IS1,
 360 IS15, IS16, IS17 and IS19). According to Brookfield's second method, null allele
 361 frequencies could range between 0.06 and 0.28 on average (Figure 2), but these
 362 estimates do not correct for other errors and, as discussed above, null alleles only partly
 363 explain the observed F_{IS} and its variation across loci.

364 These heterozygote deficits and high proportion of significant LD may come from
 365 amplification problems detected as SAD, stuttering and null alleles. Amplification problems
 366 are expected to occur in individuals presenting an apostate DNA. This can happen if these
 367 individuals belong to lineages that are significantly divergent from the majority; when their
 368 DNA is partly degraded, at low concentration; or a combination of these different causes.
 369 This may amplify a preexisting correlation between allele occurrences at different loci. One
 370 way to test for this is to study the correlation between the number of missing genotypes

371 (blanks) of an individual and its number of heterozygous sites. This correlation is expected
372 to be negative if amplification problems occur more often in some individuals than in others
373 (Kaboré et al., 2011). A Spearman's rank correlation test with Rcmdr outputted $\rho=-0.26$ (p -
374 value <0.0001).

375


376 *Cured data set*

377 To prevent the omission of five (out of nine) loci, due to stuttering, SAD and/or
378 possible selection, we went back to the data. We first scanned the chromatograms for
379 previously ignored micro-peaks in homozygous individuals at loci displaying SAD (IS1 and
380 IS15) (Figure 4). For IS1 and IS15, SAD might well have explained why we also found
381 stuttering at these locus (see below). We then tried to pool alleles close in size as
382 described above for loci IS16, IS17 and IS19. For IS16, alleles 88, 90 and 92 were
383 recoded as 94; and allele 96 as 98. For locus IS17, alleles 170 and 172 were recoded as
384 174; alleles 178, 180, 182 and 184 were recoded as 186; alleles 188, 190 and 192 were
385 recoded as 194; and alleles 196 and 198 as 200. Finally, for locus IS19, allele 91 was
386 recoded as 93; alleles 97, 99, 101 and 103 as 105; and alleles 107 and 109 as 111. The
387 obtained amended dataset was called "Cured dataset" (Supplementary file S1). Pooling
388 alleles only increases homoplasy. Everything being equal, the effect of homoplasy on F_{IS}
389 or F_{ST} is equivalent to a mutation rate increase by a factor $K/(K-1)$ (Rousset, 1996), where
390 K is the number of possible alleles. Thus, for microsatellite loci with many possible alleles,
391 the effect is deemed negligible, especially for F_{IS} . Here, the resulting number of alleles per
392 locus was 17 on average, with 5 and 27 as the outermost-lying values. The resulting
393 homoplasy effect on F -statistics can thus be safely ignored.

394

395
396

Figure 4: Examples of an initially dismissed micro-peak that produced SAD at locus IS15 and of stuttering at locus IS19




397
398

399 With the cured dataset, the proportion of locus pairs in significant LD dropped to
400 19% and the smallest p -value after BY correction was 0.0797. The correlation between
401 NLD and H_T became positive ($\rho=0.57$, p -value=0.1056). This change may point towards a
402 Wahlund effect (Manangwa et al., 2019). Nevertheless, this would be incompatible with the
403 F_{IS} observed (see below) and with the fact that none of these tests remained significant
404 after BY adjustment. A heterozygote deficit was no longer observed ($F_{IS}=-0.004$ in
405 95%CI=[-0.105, 0.124], bilateral p -value=0.7034) (Figure 5). StrdErrFIS=0.064 was three
406 times higher than StrdErrFST=0.022. There was no correlation between F_{IS} and F_{ST} ($\rho=-$
407 0.27, p -value=0.77). The correlation between number of missing genotypes and F_{IS} was
408 positive, though marginally not significant ($\rho=0.52$, p -value=0.0809). MicroChecker
409 diagnosed null alleles in 11 subsamples for locus IS11 and in one subsample for loci IR39
410 and IS19, which correspondingly displayed relatively high and variable F_{IS} and substantial
411 proportions of missing genotypes (see above and Figure 5). IS15 still displayed SAD ($\rho=-$
412 0.3814, p -value=0.0248), though with much less intensity. In addition, after BH correction,
413 the test was not significant any longer (p -value=0.2232). When IS11 and IS15 were
414 removed from the dataset, global F_{IS} became significantly negative ($F_{IS}=-0.074$ in
415 95%CI=[-0.142, -0.017], p -value=0.0002) as expected for small subpopulations in

416 dioecious species (Balloux, 2004). As for subdivision, F_{ST} remained almost unaffected,
 417 even with the ENA method ($F_{ST}=0.103$ in $95\%CI=[0.067..0.14]$, $p\text{-value}<0.0001$).
 418

419 Figure 5: F_{IS} values for each locus, averaged across those (All), or over 7 loci without
 420 amplification problems (IS11 and IS15) for *Ixodes scapularis* cured data from the
 421 eastern U.S.A. with 95% jackknife confidence intervals over subsamples (for each
 422 locus) and bootstraps over loci (All). Results of tests for panmixia, short allele
 423 dominance, number of subsamples with stuttering for each locus, and null allele
 424 frequencies are also indicated.



425


426

427 **Conclusion**

428 Combinations of amplification errors manifesting as null alleles, SAD or stuttering
 429 lead not only to heterozygote deficits, but an overall increase in LD. In order to correct for
 430 SAD, it may be useful to hunt for small and hard-to-detect peaks in chromatograms for
 431 homozygous individuals and pool alleles close in size to correct for stuttering. Uncorrected,
 432 these problems have the potential to lead to the unnecessary withdrawal of the “flawed”
 433 loci. These proposed amendments, together with null allele management with the ENA
 434 algorithm (Chapuis and Estoup, 2007), resulted in analyses revealing heterozygote
 435 excesses, as expected in small dioecious subpopulations. These changes also reduced
 436 the proportion of significant LD tests which, more importantly, were no longer significant
 437 after BY correction. Such corrections can then lead to more accurate estimates of the

438 degree of population subdivision, as shown elsewhere for null alleles (Chapuis and
 439 Estoup, 2007; Séré et al., 2017b). The different steps to be followed during population
 440 genetics data analyses of such datasets are summarized as a flowchart in Figure 6.
 441

442 Figure 6: Flowchart of the recommended steps in population genetics data analyses.



443
 444
 445 In our case, the relatively important global LD across loci is probably due to small
 446 effective sizes of the *I. scapularis* subpopulations (Waples, 2006; Waples and Do, 2010).
 447 Additionally, null alleles are still influencing the cured data, and predominantly affect
 448 individuals that display some kind of deviating DNA (as explained in the introduction) and
 449 may also contribute to inflate the perceived LD. In fact, the correlation between the number
 450 of missing genotypes and the number of heterozygous sites was still significantly negative
 451 in the cured data set ($\rho = -0.32$, p -value < 0.0001).

452 It is worthy to note that a reanalysis of the raw data set using MicroDrop 1.01 (Wang
 453 et al., 2012) resulted in smaller F_{IS} and F_{ST} values and only three significant LD tests, none
 454 of which stayed significant after BY correction (though the smaller p -value = 0.0556 was
 455 marginal). However, locus IS15 still displayed a significant SAD, (p -value = 0.0085, and
 456 corrected p -value = 0.0765) and stuttering still was detected once, twice and three times for
 457 loci IS1, IS16 and IS17, respectively, and appeared once for locus IR27. MicroDrop takes
 458 into consideration that heterozygote deficits and missing data are entirely due to allelic
 459 dropouts. It is difficult to thoroughly understand these results since here we have
 460 convincing evidence that the amplification problems were due to SAD, stuttering and null

461 alleles. This issue would require a full simulation study with different scenarios and a
462 comparison of methods. That MicroDrop may efficiently cure not only allelic dropouts but
463 any kind of amplification problem without any bias offers the potential for an extremely
464 valuable tool for the scientific community, however, this remains to be ascertained.
465 Nevertheless, MicroDrop did not entirely cure the data from SAD and stuttering, which
466 does not advocate for the efficacy of the algorithm used in this software. As far as our *I.*
467 *scapularis* data set is concerned, the cures proposed provided satisfactory results and
468 additional useful tools to those already proposed in recently published papers on the
469 detection and identification of causes of heterozygote deficits (Waples, 2015; De Meeûs,
470 2018b; Waples, 2018; Manangwa et al., 2019).

471 Subdivision into small and isolated subpopulations was confirmed by the relatively
472 small effective population size estimated from F_S (without IS11 and IS15) with Balloux's
473 method (Balloux, 2004) ($N_e \approx 7$ in 95%CI=[4, 29] individuals), and the relatively important
474 subdivision measurements between contemporaneous subsamples (to avoid temporal
475 effects) in cohorts where these were possible (i.e. cohorts 5, 7, 8 and 9) (average
476 $F_{ST} \approx 0.19$ in 95%CI=[0.1, 0.29]). These results contrast with those obtained from the
477 closely related European *I. ricinus*, with no or weak LD in adults, displaying Wahlund
478 effects at small scales, and with much weaker subdivisions (Delaye et al., 1997; De Meeûs
479 et al., 2002; Kempf et al., 2010).

480 Finally, the correlation between mitochondrial clade allocation and genetic structure
481 of *I. scapularis* is not within the scope of this work and will be treated in detail in a further
482 study that will also include immature stages.

483

484 **Data availability**

485 The raw and cured datasets are available as “supplementary file S1” at:
486 [http://www.t-de-meeus.fr/Data/DeMeeus-et-al-SAD&StutteringI-scapularisUSA-PCI-](http://www.t-de-meeus.fr/Data/DeMeeus-et-al-SAD&StutteringI-scapularisUSA-PCI-EvolBiol-TableS1)
487 [EvolBiol-TableS1](http://www.t-de-meeus.fr/Data/DeMeeus-et-al-SAD&StutteringI-scapularisUSA-PCI-EvolBiol-TableS1)

488

489 **Acknowledgements**

490 This work was funded by NSF Grant # EF0914390 to L.B. and EEID EF-0914476 to
491 J.T. We thank **the** members of the Lyme Gradient Consortium and many individuals who
492 provided ticks. We are also grateful to Heather Walker, Jenny Dickson, Keely Duff, Laquita
493 Burton, Alysha Benn and Nina Griffin, and the **several other** undergraduate students who
494 provided field and laboratory assistance. **We would like to thank one anonymous referee**

495 and Eric Petit for their comments and suggestions that helped with improving the present
496 manuscript.

497

498

499 **References**

500 Balloux, F., 2004. Heterozygote excess in small populations and the heterozygote-excess
501 effective population size. *Evolution* 58, 1891-1900.

502 Beati, L., Keirans, J.E., 2001. Analysis of the systematic relationships among ticks of the
503 genera *Rhipicephalus* and *Boophilus* (Acari : Ixodidae) based on mitochondrial 12S
504 ribosomal DNA gene sequences and morphological characters. *J. Parasitol.* 87, 32-48.

505 Benjamini, Y., Hochberg, Y., 2000. On the adaptive control of the false discovery rate in
506 multiple testing with independent statistics. *Journal of Educational and Behavioral*
507 *Statistics* 25, 60-83.

508 Benjamini, Y., Yekutieli, D., 2001. The control of the false discovery rate in multiple testing
509 under dependency. *Ann. Stat.* 29, 1165–1188.

510 Castle, W.E., 1903. The laws of heredity of Galton and Mendel, and some laws governing
511 race improvement by selection. *Proc Am Acad Arts Sci* 39, 223-242.

512 Chapuis, M.P., Estoup, A., 2007. Microsatellite null alleles and estimation of population
513 differentiation. *Mol. Biol. Evol.* 24, 621-631.

514 Coombs, J.A., Letcher, B.H., Nislow, K.H., 2008. CREATE: a software to create input files
515 from diploid genotypic data for 52 genetic software programs. *Mol. Ecol. Res.* 8, 578–580.

516 De Meeûs, T., 2012. *Initiation à la génétique des populations naturelles: Applications aux*
517 *parasites et à leurs vecteurs*. IRD Editions, Marseille.

518 De Meeûs, T., 2018a. Revisiting F_{IS} , F_{ST} , Wahlund effects and null alleles. *J. Hered.* 109,
519 446-456.

520 De Meeûs, T., 2018b. Revisiting F_{IS} , F_{ST} , Wahlund effects, and Null alleles. *J. Hered.* 109,
521 446-456.

522 De Meeûs, T., Beati, L., Delaye, C., Aeschlimann, A., Renaud, F., 2002. Sex-biased
523 genetic structure in the vector of Lyme disease, *Ixodes ricinus*. *Evolution* 56, 1802-1807.

524 De Meeûs, T., Guégan, J.F., Teriokhin, A.T., 2009. MultiTest V.1.2, a program to
525 binomially combine independent tests and performance comparison with other related
526 methods on proportional data. *BMC Bioinformatics* 10, 443.

527 De Meeûs, T., Humair, P.F., Grunau, C., Delaye, C., Renaud, F., 2004. Non-Mendelian
528 transmission of alleles at microsatellite loci: an example in *Ixodes ricinus*, the vector of
529 Lyme disease. *Int. J. Parasitol.* 34, 943-950.

530 De Meeûs, T., McCoy, K.D., Prugnolle, F., Chevillon, C., Durand, P., Hurtrez-Boussès, S.,
531 Renaud, F., 2007. Population genetics and molecular epidemiology or how to "débusquer
532 la bête". *Infect. Genet. Evol.* 7, 308-332.

533 Delaye, C., Aeschlimann, A., Renaud, F., Rosenthal, B., De Meeûs, T., 1998. Isolation and
534 characterization of microsatellite markers in the *Ixodes ricinus* complex (Acari : Ixodidae).
535 *Mol. Ecol.* 7, 360-361.

536 Delaye, C., Béati, L., Aeschlimann, A., Renaud, F., De Meeûs, T., 1997. Population
537 genetic structure of *Ixodes ricinus* in Switzerland from allozymic data: no evidence of
538 divergence between nearby sites. *Int. J. Parasitol.* 27, 769-773.

539 Fagerberg, A.J., Fulton, R.E., Black, W.C., 2001. Microsatellite loci are not abundant in all
540 arthropod genomes: analyses in the hard tick, *Ixodes scapularis* and the yellow fever
541 mosquito, *Aedes aegypti*. *Insect Mol. Biol.* 10, 225-236.

542 Fox, J., 2005. The R commander: a basic statistics graphical user interface to R. *J. Stat.*
543 *Software* 14, 1–42.

544 Fox, J., 2007. Extending the R commander by "plug in" packages. *R News* 7, 46–52.

545 Goudet, J., 1995. FSTAT (Version 1.2): A computer program to calculate F-statistics. *J.*
546 *Hered.* 86, 485-486.

547 Goudet, J., 2003. Fstat (ver. 2.9.4), a program to estimate and test population genetics
548 parameters. Available at <http://www.t-de-meeus.fr/Programs/Fstat294.zip>, Updated from
549 Goudet (1995).

550 Goudet, J., Raymond, M., De Meeûs, T., Rousset, F., 1996. Testing differentiation in
551 diploid populations. *Genetics* 144, 1933-1940.

552 Gulia-Nuss, M., Nuss, A.B., Meyer, J.M., Sonenshine, D.E., Roe, R.M., Waterhouse, R.M.,
553 Sattelle, D.B., de la Fuente, J., Ribeiro, J.M., Megy, K., Thimmapuram, J., Miller, J.R.,
554 Walenz, B.P., Koren, S., Hostetler, J.B., Thiagarajan, M., Joardar, V.S., Hannick, L.I.,
555 Bidwell, S., Hammond, M.P., Young, S., Zeng, Q.D., Abrudan, J.L., Almeida, F.C., Ayllon,
556 N., Bhide, K., Bissinger, B.W., Bonzon-Kulichenko, E., Buckingham, S.D., Caffrey, D.R.,
557 Caimano, M.J., Croset, V., Driscoll, T., Gilbert, D., Gillespie, J.J., Giraldo-Calderon, G.I.,
558 Grabowski, J.M., Jiang, D., Khalil, S.M.S., Kim, D., Kocan, K.M., Koci, J., Kuhn, R.J.,
559 Kurtti, T.J., Lees, K., Lang, E.G., Kennedy, R.C., Kwon, H., Perera, R., Qi, Y.M., Radolf,
560 J.D., Sakamoto, J.M., Sanchez-Gracia, A., Severo, M.S., Silverman, N., Simo, L., Tojo, M.,
561 Tornador, C., Van Zee, J.P., Vazquez, J., Vieira, F.G., Villar, M., Wespiser, A.R., Yang,
562 Y.L., Zhu, J.W., Arensburger, P., Pietrantonio, P.V., Barker, S.C., Shao, R.F., Zdobnov,
563 E.M., Hauser, F., Grimmelikhuijzen, C.J.P., Park, Y., Rozas, J., Benton, R., Pedra, J.H.F.,
564 Nelson, D.R., Unger, M.F., Tubio, J.M.C., Tu, Z.J., Robertson, H.M., Shumway, M., Sutton,

565 G., Wortman, J.R., Lawson, D., Wikel, S.K., Nene, V.M., Fraser, C.M., Collins, F.H.,
566 Birren, B., Nelson, K.E., Caler, E., Hill, C.A., 2016. Genomic insights into the *Ixodes*
567 *scapularis* tick vector of Lyme disease. *Nat Commun* 7.

568 Hardy, G.H., 1908. Mendelian proportions in a mixed population. *Science* 28, 49-50.

569 Hedrick, P.W., 1999. Perspective: Highly variable loci and their interpretation in evolution
570 and conservation. *Evolution* 53, 313-318.

571 Kaboré, J., MacLeod, A., Jamonneau, V., Ilboudo, H., Duffy, C., Camara, M., Camara, O.,
572 Belem, A.M.G., Bucheton, B., De Meeûs, T., 2011. Population genetic structure of Guinea
573 *Trypanosoma brucei gambiense* isolates according to host factors. *Infect. Genet. Evol.* 11,
574 1129-1135.

575 Keirans, J.E., Needham, G.R., Oliver, J.H., 1999. The *Ixodes (Ixodes) ricinus* complex
576 worldwide. Diagnosis of the species in the complex, hosts and distribution, in: Needham,
577 G.R., R., M., Horn, D.J., Welbourn, W.C. (Eds.), *Acarology IX: Symposia*. Ohio Biological
578 Survey, Columbus, Ohio, pp. 341-347.

579 Kempf, F., De Meeûs, T., Vaumourin, E., Noel, V., Taragel'ová, V., Plantard, O., Heylen,
580 D.J.A., Eraud, C., Chevillon, C., McCoy, K.D., 2011. Host races in *Ixodes ricinus*, the
581 European vector of Lyme borreliosis. *Infect. Genet. Evol.* 11, 2043-2048.

582 Kempf, F., McCoy, K.D., De Meeûs, T., 2010. Wahlund effects and sex-biased dispersal in
583 *Ixodes ricinus*, the European vector of Lyme borreliosis: new tools for old data. *Infect.*
584 *Genet. Evol.* 10, 989-997.

585 Manangwa, O., De Meeûs, T., Grébaud, P., Segard, A., Byamungu, M., Ravel, S., 2019.
586 Detecting Wahlund effects together with amplification problems : cryptic species, null
587 alleles and short allele dominance in *Glossina pallidipes* populations from Tanzania. *Mol.*
588 *Ecol. Res.* 19, 757-772.

589 Nei, M., Chesser, R.K., 1983. Estimation of fixation indices and gene diversities. *Ann.*
590 *Hum. Genet.* 47, 253-259.

591 Norris, D.E., Klompen, J.S.H., Keiransand, J.E., Black, I.W.C., 1996. Population genetics
592 of *Ixodes scapularis* (Acari: Ixodidae) based on mitochondrial 16S and 12S genes. *J.*
593 *Med. Entomol.* 33, 78-89.

594 Qiu, W.G., Dykhuizen, D.E., Acosta, M.S., Luft, B.J., 2002. Geographic uniformity of the
595 Lyme disease spirochete (*Borrelia burgdorferi*) and its shared history with tick vector
596 (*Ixodes scapularis*) in the Northeastern United States. *Genetics* 160, 833-849.

597 R-Core-Team, 2018. R: A Language and Environment for Statistical Computing, Version
598 3.5.0 (2018-04-23) ed. R Foundation for Statistical Computing, Vienna, Austria,
599 <http://www.R-project.org>.

600 Rousset, F., 1996. Equilibrium values of measures of population subdivision for stepwise
601 mutation processes. *Genetics* 142, 1357-1362.

602 Rulison, E.L., Kuczaj, I., Pang, G., Hickling, G.J., Tsao, J.I., Ginsberg, H.S., 2013.
603 Flagging versus dragging as sampling methods for nymphal *Ixodes scapularis* (Acari:
604 Ixodidae). *J. Vector Ecol.* 38, 163-167.

605 Sakamoto, J.M., Goddard, J., Rasgon, J.L., 2014. Population and demographic structure
606 of *Ixodes scapularis* Say in the eastern United States. *PLoS One* 9, e101389.

607 Séré, M., Thévenon, S., Belem, A.M.G., De Meeûs, T., 2017a. Comparison of different
608 genetic distances to test isolation by distance between populations. *Heredity* 119, 55-63.

609 Séré, M., Thévenon, S., Belem, A.M.G., De Meeûs, T., 2017b. Comparison of different
610 genetic distances to test isolation by distance between populations. *Heredity* 119, 55-63.

611 Van Oosterhout, C., Hutchinson, W.F., Wills, D.P.M., Shipley, P., 2004. MICRO-
612 CHECKER: software for identifying and correcting genotyping errors in microsatellite data.
613 *Mol Ecol Notes* 4, 535-538.

614 Wang, C., Schroeder, K.B., Rosenberg, N.A., 2012. A maximum-likelihood method to
615 correct for allelic dropout in microsatellite data with no replicate genotypes. *Genetics* 192,
616 651-669.

617 Waples, R.S., 2006. A bias correction for estimates of effective population size based on
618 linkage disequilibrium at unlinked gene loci. *Conserv Genet* 7, 167-184.

619 Waples, R.S., 2015. Testing for Hardy-Weinberg proportions: have we lost the plot? *J.*
620 *Hered.* 106, 1-19.

621 Waples, R.S., 2018. Null alleles and $F_{IS} \times F_{ST}$ correlations. *J. Hered.* In press.

622 Waples, R.S., Do, C., 2010. Linkage disequilibrium estimates of contemporary N_e using
623 highly variable genetic markers: a largely untapped resource for applied conservation and
624 evolution. *Evol. Appl.* 3, 244-262.

625 Wattier, R., Engel, C.R., Saumitou-Laprade, P., Valero, M., 1998. Short allele dominance
626 as a source of heterozygote deficiency at microsatellite loci: experimental evidence at the
627 dinucleotide locus Gv1CT in *Gracilaria gracilis* (Rhodophyta). *Mol. Ecol.* 7, 1569-1573.

628 Weinberg, W., 1908. Über den Nachweis der Verebung beim Menschen. *Jahresheft des*
629 *Vereins für Vaterländische Naturkunde in Württemberg* 64, 368-382.

630 Weir, B.S., Cockerham, C.C., 1984. Estimating F-statistics for the analysis of population
631 structure. *Evolution* 38, 1358-1370.

632 Wright, S., 1965. The interpretation of population structure by F-statistics with special
633 regard to system of mating. *Evolution* 19, 395-420.

634 Yuval, B., Spielman, A., 1990. Duration and Regulation of the Developmental Cycle of
635 *Ixodes dammini* (Acari: Ixodidae). J. Med. Entomol. 27, 196-201.
636
637