

Your manuscript on altitudinal divergence in snapdragon plants treats an interesting and timely question mostly with appropriate methods. However, I do have several difficulties to evaluate the quality of this research because important information appears to be missing from the manuscript. This concerns: I), most importantly, whether there were genetic or phenotypic differences between two subspecies that were apparently pooled here, according to Material and Methods and II) the experimental design such as spatial arrangement in the garden (how plants were arranged in how many containers) that should be taken account of as a random effect in the analyses, as well as crossing design and number of individuals per family used (see comments below).

Regarding the subspecies, an explicit analysis would be most helpful (phenotypic and genetic differentiation, for example PCoA or structure analysis, potentially presented in the supplementary material). Pooling of the subspecies would probably only be appropriate if there are no considerable phenotypic differences or genetic differentiation between them.  $F_{st}$  in these analyses is used to reflect putatively neutral genetic differentiation -- this might be difficult assumption if the dataset includes subspecies that are partially reproductively isolated from each other. The description of the subspecies in the method section as well as the citations do fuel this concern. Differences in habitat and flower color between subspecies are described —even pollinator-mediated reproductive isolation could be present in this system. Very careful and convincing analyses and explanations would be needed to motivate the pooling. And —if there are no differences between subspecies, would that not suggest that there might just be a flower color polymorphism?

It is important to note that, without this potentially troublesome pooling of subspecies, the number of populations would probably be too low for the present analysis and this is why this issue is so critical.

Provided that pooling can be justified, and the design clarified, this manuscript could become a good one, however, further improvements in the analysis and presentation are also needed. The main changes that I believe are necessary are:

1) Focus on biology. The paper would be more interesting if focus was shifted from methods and general interpretations to the biology of altitudinal adaptation. This would be facilitated by presenting the trait-altitude correlations first. These results are interesting for all traits, currently they are only presented for 3 out of 13 traits. From that you could move on to more complex analyses to identify the underlying causes of such trait clines. In the current version, there is overall rather little on HOW the traits change along altitude, for example, there are no details on leaf traits, even though there is a lot of literature of variation in SLA (see citations in literature mentioned below). It would also be very good if explicit expectations for all traits or for trait groups, based on the literature, could be added in the introduction. I would also appreciate an introduction ending in specific questions.

2) In the methods part, you write that you used the between family variance component to estimate the within population additive genetic variance for a trait, as is commonly done. However, Table S1 states, for reproductive traits, the total number of families is 372 and the number of individuals is 380, such that you cannot have had more than one individual per family in most cases, unless you used a subset of families with more individuals (I could not find information on the number of individuals per family). With 1 individual per family it should be very difficult or impossible to estimate the between family variance component. In Table S3 you do not give the within family variance component so I wonder whether what you state as the within population variance component really is the between family variance component. This issue is in need of clarification. It would also be good if you could explain why you have fewer measurements for reproductive traits than for vegetative traits.

3) Methods, QST - FST. I found the description of the QST - FST comparisons, using expected distributions, unclear and in part contradictory (see comments in the text). In particular, it is not clear to me how you arrived at the confidence intervals for QST on Figure 2. It would fit better if you presented the distribution of expected values for QST together with the observed values, this is the method you stated in the methods section. Alternatively, you could change the methods section.

4) Analysis of pairwise data: the Methods section describes the commonly used and suitable analyses for correlation tests among matrices (pairwise data), Mantel tests and partial Mantel tests, controlling for the effect of neutral genetic divergence. In the results however, a regression analysis is presented (Fig. 3) that is not suitable for this type of non-independent data points. The text states that the Mantel test was used to assess significance of the regression — this is not possible. The actual data points are

also not displayed, this is a bit worrisome due to the subspecies issue. My suggestion is to re-work this analysis according to the methods described and to display the data points on Fig. 3.

5) (Local) adaptation, interpretation. The interpretation of what constitutes local adaptation is highly variable throughout the manuscript. At some places, the common cautious interpretation of QST - FST analyses, that they can provide hypotheses on traits that may have adaptively diverged due to selection (see for example review by Leinonen et al. 2013, Nature Reviews Genetics) is used. At quite some other places, however (abstract, introduction, discussion, marked in detailed comments), it appears that local adaptation is inferred whenever trait divergence cannot be attributed to either the altitudinal gradient or neutral divergence. I cannot be sure that the text is really meant that way, of course, but I testify here that it reads that way. This interpretation would be incorrect (genetic drift and many other processes could be responsible for trait divergence instead). Any text that can potentially be misunderstood is probably better changed.

6) Interpretation of the results on germination. The corresponding paragraph in the discussion unfortunately appears contradictory and the two types of results, overall QST - FST and pairwise QST - FST have not been compared (see comments in the text). Moreover, the text also appears as if it was not clear to the authors that a lack of quantitative genetic differentiation in traits does not exclude phenotypic plasticity (i.e., this is not really an alternative interpretation). It seems that this part needs adjustment.

7) Interpretation of genetic divergence. You invoke reproductive isolation, when it appears that isolation by distance would be a sufficient interpretation, unless you refer to the subspecies differences here (see comments above and in the text). However, this cannot be evaluated from the data currently presented, as details on genetic differentiation are missing (see above). I suggest that this interpretation is re-worked once a more detailed analysis is presented.

8) Literature. The manuscript would much benefit by adding more literature on the analysis of altitudinal gradients and transplant experiments, for example: Luo, Y., A. Widmer, and S. Karrenberg. 2015. The roles of genetic drift and natural selection in quantitative trait divergence along an altitudinal gradient in *Arabidopsis thaliana*. *Heredity* 114:220.  
Halbritter, A. H., S. Fior, I. Keller, R. Billeter, P. Edwards, R. Holderegger, S. Karrenberg, A. R. Pluess, A. Widmer, and J. M. Alexander. 2018. Trait differentiation and adaptation of plants along elevation gradients. *J. Evol. Biol.*

[Disclaimer: these are papers I am an author on-- but they really are related here... as are many of the citations in them].

9) Methods are not really new. Throughout the intro and the discussion, the paper reads as if this mainly is a method advancement. This appears somewhat exaggerated, as similar methods have been used before and since a longer time (>5 yrs). I feel that the work of others should be acknowledged much more here, only few citations on such studies that are able to identify selective agents are given (for example, Hangartner et al).

10) Shortening. I find that the text could likely benefit from shortening, in particular in the introduction and discussion, probably to at least 2/3 of its present length. I have indicated several parts that appear repetitive in the detailed comments.

11) More comments. I have made comments similar to the above comments as well as many further detailed comments directly on the text using pdf commenting tools in Adobe Acrobat. I am not repeating these detailed comments here. Such a list of details would be very difficult to write and read, because the text did not include any line numbers to refer to.

I hope that you will find my comments clear and helpful for further revisions of your work.

With best wishes,

Sophie Karrenberg

## **Evolutionary inference from $Q_{ST}$ - $F_{ST}$ comparisons: disentangling local adaptation from altitudinal gradient selection in snapdragon plants**

Sara Marin<sup>1</sup>, Juliette Archambeau<sup>1,2</sup>, Vincent Bonhomme<sup>3</sup>, Mylène Lascoste<sup>1</sup> and Benoit Pujol<sup>1,4\*</sup>

<sup>1</sup> Laboratoire Évolution & Diversité Biologique (EDB UMR 5174), Université Fédérale de Toulouse Midi-Pyrénées, CNRS, IRD, UPS, Toulouse, France

<sup>2</sup> BIOGECO, INRA, University of Bordeaux, Pessac, France

<sup>3</sup> Institut des Sciences de l'Évolution (ISEM), équipe "Dynamique de la biodiversité, anthropo-écologie", UMR 5554, Université de Montpellier, CNRS, IRD, EPHE, Place Eugène Bataillon, Cc 065, 34095 Montpellier cedex 05, France

<sup>4</sup> EPHE, PSL Research University, UPVD, CNRS, USR 3278 CRIOBE, F-66360 Perpignan, France.

\* Author for correspondence: [benoit.pujol@criobe.pf](mailto:benoit.pujol@criobe.pf)

## ABSTRACT

Phenotypic differentiation among natural populations can be explained by natural selection or by neutral processes such as drift. There are many examples in the literature where comparing the effects of these processes on multiple populations has allowed the detection of local adaptation. However, these studies rarely identify the agents of selection. Whether population adaptive divergence is caused by local features of the environment, or by the environmental demand emerging at a more global scale, for example along altitudinal gradients, is a question that remains poorly investigated. Here, we measured neutral genetic ( $F_{ST}$ ) and quantitative genetic ( $Q_{ST}$ ) differentiation among 13 populations of snapdragon plants (*Antirrhinum majus*) in a common garden experiment. We found low but significant genetic differentiation at putatively neutral markers, which supports the hypothesis of either ongoing pervasive homogenisation via gene flow between diverged populations or reproductive isolation between disconnected populations. Our results also support the hypothesis of local adaptation involving phenological, morphological, reproductive and functional traits. They also showed that phenotypic differentiation increased with altitude for traits reflecting the reproduction and the phenology of plants, thereby confirming the role of such traits in their adaptation to environmental differences associated with altitude. Our approach allowed us to identify candidate traits for the adaptation to climate change in snapdragon plants. Our findings imply that environmental conditions changing with altitude, such as the climatic envelope, influenced the adaptation of multiple populations of snapdragon plants on the shape of their adaptation to local environmental features. They also have implications for the study of adaptive evolution in structured populations because they highlight the need to disentangle the adaptation of plant populations to climate envelopes and altitude from the confounding effects of selective pressures acting specifically at the local scale of a population.

## KEYWORDS

Local adaptation, altitudinal gradient, climate change, quantitative genetics, *Antirrhinum majus*

## INTRODUCTION

Local adaptation - the evolutionary response to local selection that makes populations fitter in their own local habitat than in any other population's local habitat - is widespread in both plant and animal species (Kawecki & Ebert, 2004; Tuomas Leinonen, McCairns, O'Hara, & Merila, 2013). One very popular indirect approach to assess whether local adaptation has shaped trait quantitative genetic variation is the  $Q_{ST}$ - $F_{ST}$  comparison (McKay & Latta, 2002; Merila & Crnokrak, 2001; Spitze, 1993). It consists in the comparison of population genetic differentiation (e.g.,  $D$  and  $F_{ST}$ ) estimated for putatively neutral molecular markers (Edelaar & Björklund, 2011; Rousset, 1997) with population quantitative genetic differentiation ( $Q_{ST}$ ) estimated for phenotypic traits of interest (Whitlock, 2008). The objective is to test whether trait quantitative genetic differentiation among populations is more likely the result of divergent selection or neutral evolutionary divergence (e.g., as a result of drift). This approach has three main advantages. First, the necessary data is relatively easy to acquire (phenotypic measures and genotypes). Second, population genetic differentiation statistics are relatively easy to estimate. Third, conclusions are mostly derived straightforwardly from results:  $Q_{ST} > F_{ST}$  is taken as evidence for local adaptation,  $Q_{ST} < F_{ST}$  is taken as evidence for stabilizing selection, and  $Q_{ST} = F_{ST}$  means that selection is not required to explain population differentiation because neutral evolution might be responsible for the divergence. There has been some debate around the accuracy of  $Q_{ST}$ - $F_{ST}$  approaches. As a result, the advantages and limits of this method are relatively well known, and a variety of possible methodological adjustments to the estimation of differentiation are available (De Villemereuil & Gaggiotti, 2015; Edelaar, Burraco, & Gomez-Mestre, 2011; Ovaskainen, Karhunen, Zheng, Arias, & Merilä, 2011; Whitlock, 2008; Whitlock & Gilbert, 2012). Although some of these adjustments were later discarded (Edelaar et al., 2011; Whitlock, 2011), more than thirty years of extensive use have resulted in wide acknowledgement of the  $Q_{ST}$ - $F_{ST}$  approach as a method testing for the hypothesis that local adaptation might explain observed patterns of differentiation. The influence of environmental effects on phenotypic variation that might result in biasing  $Q_{ST}$ - $F_{ST}$  comparisons conducted directly in wild populations has also been debated (Pujol, Wilson, Ross, & Pannell, 2008). Yet, the inclusion of ecological factors in this type of analysis remains rare (Hangartner, Laurila, & Räsänen, 2011; Karhunen, Ovaskainen, Herczeg, & Merilä, 2014). Consequently, the identification of local habitat selective pressures that might drive local adaptation is often missing from  $Q_{ST}$ - $F_{ST}$  approaches.

The general absence of identified ecological drivers for local adaptation in  $Q_{ST}$ - $F_{ST}$  approaches is not surprising. This is likely because it is a correlative approach, which limits conclusions about causality, and because it is impossible to record and include in the analysis every ecological variable differing between populations. Consequently, confounding effects from unmeasured variables can never be ruled out. It is nevertheless possible to investigate the link between changes in population divergence and changes in ecological variables at the scale of multiple populations. This does not allow identifying the local features of the environment of every population that drive local adaptation *per se*. It can however shed light on traits that are involved with adaptation to a particular ecological background at a larger scale. In fact, the exploration of ecologically explicit adaptive scenarios based on  $Q_{ST}$ - $F_{ST}$  approaches that are not particularly “local” has broader implications in the actual context of environmental change (e.g., climate change, pollution) because they affect multiple populations simultaneously. For example, identifying candidate traits that are involved with adaptation to different climates has obvious implications for climate change remediation (De Villemereuil, Mousterde, Gaggiotti, & Till-Bottraud). Other approaches targeting the molecular variation of adaptive genes also allow us to investigate the same question (Ahuja, de Vos, Bones, & Hall, 2010; Hoffmann & Sgro, 2011), but most of them cover less populations and are more demanding in time and resources. Breaking down this accessibility barrier by using less demanding approaches has implications for the generalisation of findings. It is therefore only logical that the general challenge of  $Q_{ST}$ - $F_{ST}$  approaches is shifting progressively from detecting the signature of local adaptation to identifying the ecological drivers of selection (Hangartner et al., 2011; Whitlock, 2008).

There is a growing number of studies investigating whether population quantitative genetic divergence is shaped by environmental constraints (Gomez-Mestre, Tejedo, & Ashley, 2004; Palo et al., 2003; Raeymaekers, Van Houdt, Larmuseau, Geldof, & Volckaert, 2007). For example, Hangartner et al. (2011) showed that ~~population~~ adaptive divergence in metamorphic size was associated with pond acidity in moor frogs. They also identified that other traits such as larval period and growth rate were involved with adaptation to latitude and predation. This was possible because they investigated whether the increase of environmental differences between populations was associated with an increase in  $Q_{ST}$  for various traits. This type of approach can ultimately inform us about the vulnerability of populations to climate change because comparing their adaptive divergence along altitudinal gradients can serve as a first-order approximation of the possible effects of ongoing climate change (Byars, Papst, & Hoffmann, 2007; Gonzalo-Turpin & Hazard, 2009). There is evidence for the adaptive

signature of past climate change in multiple populations growing number of studies on this topic (De Villemereuil et al.; Luquet, Léna, Miaud, & Plénet, 2014; McKay et al., 2001; Muir, Biek, Thoma, & Mable, 2014). For example, Muir et al. (2014) showed that local adaptation had shaped the five traits they had measured in common frog populations by using a common garden experiment, and that two of these traits, i.e., larval period and growth rate, played a role in their adaptation to altitude. In plants, reciprocal transplants between altitudes that are used to compare plant fitness directly in their native habitat and in the foreign habitat have been mostly preferred to  $Q_{ST}$ - $F_{ST}$  indirect approaches that are conducted in common gardens (Angert, Schemske, & Geber, 2005; Etterson, 2004; Kim & Donohue, 2013). Using an indirect approach in controlled conditions can however present an advantage when the conditions for the reciprocal transplant cannot be met, e.g., when gene flow between populations on multiple sites that would result in mixing up the genomic background of all these populations cannot be avoided. In such cases,  $Q_{ST}$ - $F_{ST}$  comparisons along altitudinal gradients might present a rare opportunity for exploring hypotheses about candidate traits for the adaptation of plants to climatic differences.

Functional ecology studies predict that eco-physiological and reproductive traits play a crucial role in the adaptation of plants to the later and shorter availability of favourable environmental conditions that characterise seasons at higher altitudes (Körner, 1999). Selection should therefore favour smaller plants, germinating later but reproducing earlier, with faster growth at higher altitude. By reciprocity, plants from lower altitude that are confronted with warmer climates should present opposite adaptations. However, these hypotheses have rarely been tested using empirical data from multiple natural populations across altitudinal gradients. Here we present the results from a common garden experiment where we evaluated the adaptive significance of multiple phenological, morphological and functional traits in populations of snapdragon plants (*Antirrhinum majus* L.). We estimated neutral genetic differentiation ( $F_{ST}$ ), trait heritability ( $h^2$ ), and quantitative genetic differentiation ( $Q_{ST}$ ) among 13 *A. majus* populations. We estimated the genetic differentiation between populations at putatively neutral microsatellite markers (Debout, Lhuillier, Malé, Pujol, & Thébaud, 2012; Pujol et al., 2017) to test for the hypothesis that gene flow was limited between populations, which sets the stage for local adaptation. We then tested for the hypothesis of local adaptation by comparing  $Q_{ST}$  and  $F_{ST}$ . ally, we investigated whether quantitative genetic differentiation increased with altitude, with the hypothesis that broader environmental changes associated with altitude, such as the climatic envelope, can explain more population adaptive trait differentiation than local


population differences. Our approach ultimately participates to the collective evaluation of the  $Q_{ST}$ - $F_{ST}$  approach, which is generally used for the study of local adaptation in structured populations as a tool to identify the adaptive response of local populations to altitudinal gradients. 

## MATERIAL AND METHODS

### Study system

*Antirrhinum majus* L. (Plantaginaceae) is a hermaphroditic, self-incompatible, short-lived perennial, which produces annual inflorescences with zygomorphic flowers. These flowers are arranged in raceme and produced at the end of the stem, thereby preventing further vegetative elongation (terminal flowering). This plant is characterized by a patchy distribution in southern Europe centred over the Pyrenees Mountains. Two subspecies, that are interfertile, occupy this geographic area: *A. m. ssp. pseudomajus*, characterized by magenta flowers, and *A. m. ssp. striatum*, characterized by yellow flowers. The geographic range of *A. m. striatum* is surrounded by the range of *A. m. pseudomajus* (Khimoun et al., 2011). Within the species geographic range, the two subspecies are distributed parapatrically and come into contact at the margins of their ranges (Andalo et al., 2010). The transition between subspecies in the contact zones occurs over a very short distance (<1km) (Whibley et al., 2006). In the west part and the east part of the contact perimeter, there is evidence for introgressive hybridization between *A. m. ssp. pseudomajus* and *A. m. ssp. striatum* (Khimoun et al., 2011). The clear separation between the distribution of *A. m. ssp. pseudomajus* and *A. m. ssp. striatum* is not explained by habitat differences, as illustrated by the substantial overlap of environmental conditions between the two species (Khimoun et al., 2013). In fact, the range of environmental conditions characterizing the ecological niche of *A. m. ssp. pseudomajus* is almost twice as large as that of *A. m. ssp. striatum* (Khimoun et al., 2013). Both subspecies occur from sea level to an altitude of 1900 m (Andalo et al., 2010), on limestone or siliceous substrates and in habitats with contrasted moisture regimes (rainfall 500-1000 mm per year), where they form restricted patches mostly in rocky outcrops and screes. *A. majus* thrives in disturbed habitats, and is especially common along roadside and railway embankments. Thirteen wild populations were sampled in 2011 across the geographic range (between north-eastern Spain and south-western France), with four populations of *A. m. ssp. striatum*, six populations of *A. m. ssp. pseudomajus*, and three populations (one *A. m. ssp. striatum*, and two *A. m. ssp. pseudomajus*) from the contact zones (Figure 1; Table S1, Supporting Information). For each subspecies and in the contact

zone, we sampled populations from low and high altitude habitats. This is because populations sampled along elevation gradients are likely to be confronted to contrasted environmental conditions (Figure S1, Supporting Information). Populations were sampled in different valleys or on different summits in order to avoid spatial autocorrelation in the data, and to avoid shared phylogeographic history between populations from similar altitudes.



**FIGURE 1** Map of *Antirrhinum majus* populations that were sampled across the geographic range of the species in Southern France. Dots represent *A. m. ssp. pseudomajus* populations, squares represent *A. m. ssp. striatum* populations and triangles represent populations from the contact zone. Grey lines represent elevation isoclines every 250 m.

### Common garden experiment

In each wild population, mature plants of *A. majus* were randomly sampled for seeds across their entire spatial distribution in October 2011 (Table S1, Supporting Information). We avoided maternal environmental effects in our common garden experiment by producing a seed bank from plants grown in similar conditions in a greenhouse at the CNRS Station of Experimental Ecology in Mounss. Seeds sampled in the wild were sown in spring 2012 in individual pots (9 × 9 × 10 cm) filled with universal compost. Plants germinated and grew with no nutrient addition under an average temperature from 15 to 28°C and weekly watering.

Mature plants were hand-pollinated during summer 2012. Crosses were conducted within populations where mates were assigned randomly. Seeds were stored in the seed bank (room temperature, dark, dry conditions). We sowed seeds from the seed bank in spring 2014 in a common garden at ENSFEA (Toulouse, France). Plants were grown outdoor in individual pots (9 × 9 × 10 cm) that were positioned in plastic containers (600 × 400 × 120 mm). Their growth was monitored during the summer 2014. Plants grew in pots filled with universal compost, with no nutrient addition, under outdoor climatic conditions (average month temperatures ranging from 20.6 to 21.5°C and cumulative monthly rainfall ranging from 28.3 to 73.4 mm). The bottom of each container was covered with an irrigation sheet (400 g.m<sup>-2</sup>) that allowed the regulation of compost moisture. Plants were supplied with water in case of prolonged drought. Damage caused by herbivorous insects were contained by using a wintering veil. This veil also limited pollination. The spatial location of each plant in the experimental setting was distributed randomly to avoid spatial effects on plants caused by uncontrolled environmental conditions.

### **Molecular and phenotypic data**

In terms of molecular markers, we genotyped the 13 different populations ( $N = 637$  individuals) for 23 putatively neutral microsatellite markers that were developed for population genetic studies (Debout et al., 2012; Pujol et al., 2017). In terms of phenotypic traits, we measured multiple phenological, morphological, functional and reproductive characters on each individual. Phenological traits included germination date, flowering date, and time to flowering (the number of days separating the germination of a plant and the emergence of its first flower). At the time of first flowering, we measured vegetative traits: main stem basal diameter (Diameter), main stem vegetative length (Vegetative Length), number of vegetative nodes on the main stem (Nodes), total number of leaves on the plant (Leaves), and number of branches on the plant (Branches). We also measured these vegetative traits for plants that did not flower but stopped growing at the end of experiment. As a result, vegetative traits were characterised for all plants. At the end of the experiment, we also measured reproductive traits on flowering plants ( $N = 380$ ): stem total length (Total Length), inflorescence length, number of flowers (Flowers). Additionally, we measured a functional trait when possible: specific leaf area (SLA). SLA was calculated as the ratio between the cumulated area of five mature but non senescent fresh leaves and their oven-dried mass (Pérez-Harguindeguy et al., 2016; Pujol, Salager, Beltran, Bousquet, & McKey, 2008). Leaf area was measured by using the R package Momocs v. 1.2.9 (Bonhomme, Picq, Gaucherel, & Claude, 2014). We also measured a developmental trait, average Internode Length for the total stem vegetative length.

## **$Q_{ST}$ - $F_{ST}$ comparison to evaluate local adaptation and altitudinal selection**

To explore **the effect of local adaptation on traits**, we tested for the difference between the  $Q_{ST}$  observed value (global population quantitative genetic differentiation for a trait) and its expected value under the hypothesis of neutral evolution by using the approach of Whitlock and Guillaume (2009). As a result, we did not compare  $Q_{ST}$  with average  $F_{ST}$  but with the  $Q_{ST}$  **expected** distribution under a scenario of neutral evolution. This distribution of neutral  $Q_{ST}$  was obtained by using simulations parametrized on the basis of observed mean  $F_{ST}$  (neutral genetic differentiation at molecular markers) and predicted from the  $\chi^2$  distribution. **The expected neutral  $Q_{ST}$  distribution for each trait was generated by bootstrapping ( $N = 10^5$ ).** We then inferred  $P$  values associated with our point estimates of  $Q_{ST}$ - $F_{ST}$  differences. **Finally, we used** **the modification by Lind et al. (2010) of the approach of Whitlock and Guillaume (2009) to estimate variance components.** The choice of this method was motivated by the fact that it suits perfectly data like ours (population count around ~~ten~~ around ten, and availability of molecular markers such as microsatellites rather than genomic markers). All procedures were implemented in the **R environment v. 3.5.0 (R-Core-Team, 2018).** **then** evaluated the correlation between matrices of pairwise population altitudinal and  $F_{ST}$  distances, and between pairwise population altitudinal and  $Q_{ST}$  distances. The significance ( $\alpha=0.05$ ) of these correlations was tested by using Mantel partial correlation tests (Mantel, 1967). Some drawbacks have been reported for this method, in particular in the presence of spatial autocorrelation. However, we limited the possibility for data spatial autocorrelation in altitude at the scale of populations by using a specific sampling scheme (see above). The option of using a Bayesian approach to overcome this issue was not available to us as it would have required using around 1000 molecular markers (Bradburd, Ralph, & Coop, 2013). We tested for Mantel partial correlations by using 9999 permutations (*mantel.rtest* command in the R package *ade4* (Chessel, Dufour, & Thioulouse, 2004)).

## **Calculation of $F_{ST}$ , $h^2$ , and $Q_{ST}$**

To compute  $F_{ST}$ , we used estimates from the study by Pujol et al. (2017) for both pairwise population  $F_{ST}$  estimates and global population overall estimates. Potential effects on population genetic differentiation of geographic coordinates (Latitude and Longitude), pairwise geographic distance, subspecies and altitude were controlled for by using an AMOVA approach. Genetic diversity at each locus, which measure was required for the  $Q_{ST}$ - $F_{ST}$  comparative approach that we used, was estimated by using the GENETIX 4.05 program


(Belkhir, Borsa, Chikhi, Raufaste, & Bonhomme, 2004). Narrow-sense heritabilities ( $h^2$ ) were estimated for each phenotypic trait. They were calculated as  $h^2 = V_A / V_P$ , with  $V_P$  the population phenotypic variance and  $V_A$  the additive genetic variance.  $V_A$  was estimated as  $2V_W$ , with  $V_W$  being the within-population genetic variance.  $V_W$  was estimated by the among-family variance. We multiplied  $V_W$  by two in the calculation of  $h^2$  because we used a full-sib crossing design (Roff, 1997). No environmental source of phenotypic variance due to the ecological conditions of the location of origin of populations could in theory bias  $Q_{ST}$  estimates because data was obtained from a common garden experiment (Benoit Pujol et al., 2008). Furthermore, at the scale of the common garden, plant locations were assigned randomly.  $Q_{ST}$  estimates were calculated by using the following formula (Spitze, 1993):  $Q_{ST} = V_B / (V_B + 2V_W)$  with  $V_B$  being the trait genetic variance among populations. All variance components were estimated by using a Linear Mixed Model approach implemented in the R package lme4 v. 1.1.17 (Bates, 2005). When a variance component was not significant, it was considered as null in further calculations. In these models, population and family were included as random factors. We verified the normality of the distribution of residuals (Shapiro test) and their homoscedasticity (Breusch-Pagan test). When necessary (as for pairwise  $Q_{ST}$  calculation), data was linearized by using a square root transformation.

## RESULTS

### Local adaptation

Average phenotypic values and their standard errors were compared among the 13 experimental populations for the 13 analysed traits (Table S2, Supporting Information). For phenological traits, average germination dates of populations spanned across seven days (ranging from the 194<sup>th</sup> to the 201<sup>th</sup> calendar day of 2014) while the average flowering date spanned across 15 days (ranging from the 236<sup>th</sup> to 251<sup>th</sup> calendar day of 2014). Among populations, the time to flowering ranged from 43 to 59 days. For vegetative traits, amongst populations, Diameter ranged from 3.37 to 4.14 mm, Vegetative Length ranged from 19.95 to 35.74 cm, Nodes ranged from 10 to 17, Leaves ranged from 95 to 152, and Branches ranged from 12 to 20. Total Length, which included the length of the inflorescence, ranged from 25.18 to 54.49 cm. For reproductive traits, inflorescence length ranged from 6.86 to 20 cm and Flowers ranged from 6 to 19. Finally, we measured an average SLA ranging from 161.24 to 194.48 cm<sup>2</sup>.g<sup>-1</sup> and an Internode Length ranging from 1.75 to 2.43 cm.

Although not directly confronted to trait  $Q_{ST}$  values, we have represented on Figure 2 the low but significant average  $F_{ST}$  among populations ( $F_{ST} = 0.107$ ,  $P < 0.001$ ), which ranged from 5 to 20% in order to illustrate population neutral genetic differentiation (for more details on population pairwise neutral genetic differentiation, see Pujol et al., 2017). We found that most trait  $Q_{ST}$  values were not in the tail of the expected probability distribution of neutral traits, which means that the hypothesis of neutrality was rejected for these quantitative traits. Trait  $Q_{ST}$  values are represented on Figure 2. These values were generally significantly higher than the simulated distribution of their expected neutral values (Figure 2, Table S3, Supporting Information). However, for germination date,  $Q_{ST}$  was in fact lower than could be expected under neutrality. For time to flowering,  $Q_{ST}$  was in the tail of the expected probability distribution of neutral traits. Heritability estimates were generally comprised between 0.07 and 0.23. However, germination date, time to flowering, Nodes and Internode Length, had a heritability between 0.36 and 0.47 (Table S3, Supporting Information). Caution must be taken when considering  $h^2$  values because its estimation can be biased by the estimation of  $V_A$ . Indeed,  $h^2$  was calculated on the basis of all the families, without taking into account the differences of  $h^2$  between different populations. The family variance component was not significant and thereby considered null ( $V_W = 0$ ) for the Leaves, making simulations by the approach of Whitlock and Guillaume (2009) impossible.




**FIGURE 2** Values for observed  $Q_{ST}$  (represented by dots with its 95% confident interval indicated by black lines).  ~~$P$ -values~~ of statistical differences between observed values and the expected distribution of neutral traits are indicated by \* for significant differences, or *ns* for non-significant differences. Average population  $F_{ST}$  is represented by the dashed line ( $P < 0.001$ ).

### Increased quantitative genetic differentiation with altitude

The larger the difference in altitude was for a pair of populations, the higher was their neutral genetic differentiation, as estimated by  $F_{ST}$ . This altitudinal effect was weak but significant (Figure 3). The relationship with altitude was also significant for quantitative genetic differentiation as measured by  $Q_{ST}$ . It was only significant for three traits out of the 13 traits



that we measured (germination date, inflorescence length, Flowers). The slope of the altitudinal increase in differentiation was one order of magnitude higher for  $Q_{ST}$  than  $F_{ST}$  (Figure 3). Indeed, we found that per unit elevation of 500 meters,  $Q_{ST}$  of germination date increased by 11%,  $Q_{ST}$  of inflorescence length increased by 12.5%, and  $Q_{ST}$  of Flowers increased by 13.5%, while  $F_{ST}$  increased by 1.5%.




**FIGURE 3** Linear regression (coloured lines) of population pairwise quantitative trait differentiation ( $Q_{ST}$ ) on population pairwise differences in altitude (m) for Germination date, Inflorescence length and Flowers. Linear regression (grey line) of the population pairwise neutral genetic differentiation ( $F_{ST}$ ) on population pairwise differences in altitude (m). Regression equations and levels of significance ( $P$ ), obtained by using Model tests are presented in the figure. 95% confidence intervals are indicated by shaded areas.



### Changes in phenotypic values with altitude

Germination occurred significantly later in experimental populations that originated from higher altitudes. Germination date increased by 0.59% (1.15 days) per unit elevation of

500 meters (Figure 4a). Plants from populations originating from higher altitudes showed a significant reduction in the size of their inflorescence and in the number of flowers. Inflorescence length decreased by 13.14%, and Flowers decreased by 12.43% per unit elevation of 500 meters (Figure 4b, c).



**FIGURE 4** Variation of mean ( $\pm$ SE) per populations for Germination date (a), Inflorescence length (b) and Flowers (c) with altitude. Associated regression (solid line) equations and level of significance ( $P$ ) are presented for each phenotypic trait. Dots represent *A. m. ssp. pseudomajus* populations, squares represent *A. m. ssp. striatum* populations and triangles represent populations from the contact zone.

## DISCUSSION

Our results showed that neutral evolution and local adaptation could not explain on their own the extent to which *A. majus* populations from the highest altitudes had diverged from lower populations. Phenotypic differentiation increased significantly with altitude for the germination date, inflorescence size and number of flowers, with more than a two-fold increase at the highest altitudes. Our results also provide support for the hypothesis that neutral evolution had the potential to shape the diversity of *A. majus* populations over their geographic range. However, this potential could only account for around 10% of population phenotypic differentiation. On the top of their neutral divergence, it was necessary to invoke local adaptation for explaining how *A. majus* populations diverged from each other at multiple phenological, morphological, reproductive and functional traits. Our findings comfort the emerging idea that  $Q_{ST}$ - $F_{ST}$  comparisons can be used to go further than detecting the local adaptation of populations to the local demand of their surrounding - often-unidentified - environmental conditions (Edelaar et al., 2011; Ovaskainen et al., 2011; Whitlock, 2008; Whitlock & Gilbert, 2012). They highlight how some traits involved with adaptive evolution at the population level can be used to identify the ecological pressures underlying natural selection. They revealed that some traits, but not all the traits involved with *A. majus* local adaptation, are under natural selection by the ecological conditions changing with altitude that are tightly linked to the climatic envelope of populations.

### Support for the local adaptation scenario

Our results showed that quantitative genetic divergence was higher amongst *A. majus* populations for ten of the thirteen traits under study than what could be explained by neutral evolutionary divergence. They imply that local adaptation has shaped the phenotypic diversity of *A. majus* populations across their geographic range. Local adaptation is detected in the majority of  $Q_{ST}$ - $F_{ST}$  comparisons (T. Leinonen, O'Hara, Cano, & Merila, 2008). Our finding is robust against a range of neutral evolution scenarios for these traits that were extrapolated from the distribution of  $F_{ST}$  values based on multiple putatively neutral markers (Whitlock, 2008). Furthermore, our approach excludes the possibility that plasticity rather than divergent selection might have generated this signal of phenotypic divergence because we used a common garden experiment, and included trait heritability estimates in  $Q_{ST}$  calculations (Benoit Pujol et al., 2008; Spitze, 1993). Ten of the thirteen traits under study harboured the signature of divergent selection but two traits (the time to flowering and the number of the leaves) did not show departure from the baseline scenario of neutral evolution. One particular trait (germination date) was in fact more similar among populations than expected under neutrality. A scenario of stabilising selection is classically extrapolated in the case of similar results (Lamy, Plomion,

Kremer, & Delzon, 2012) but we argue that trait homogenisation caused during the experiment by phenotypic plasticity might be another plausible explanation.

The most likely evolutionary scenario applying to *A. majus* populations requires invoking a history of local adaptation in a complex background of gene flow and reproductive isolation. Neutral genetic divergence was weak but significant amongst populations. This indicates that mechanisms such as genetic drift have likely shaped differentially the genetic background of *A. majus* populations at the scale of their global geographic range. This can be interpreted as the genetic signature of reproductive isolation which restricted at least partly gene exchanges (Pujol et al., 2017). The mountain landscape in the Pyrenees was found to have likely reinforced reproductive isolation between *A. majus* populations (Pujol et al., 2017). However at the scale of *A. majus* neighbour populations from a contact zone, no genome wide barrier to gene flow was found (Ringbauer, Kolesnikov, Field, & Barton, 2018). At such local scale, gene flow is very likely rapidly homogenising population neutral genetic variation. Our findings thereby imply that Pyrenees mountains, which constitute a heterogeneous landscape that promotes complex patterns of connectivity amongst plant populations (Alberto et al., 2010), were prone to generate local adaptation in *A. majus*.

### **The ecological significance of local adaptation in *A. majus***

In the absence of environmental measures included in the  $Q_{ST}$ - $F_{ST}$  analysis, it is impossible to identify the potential environmental agents of local selection that shape the quantitative genetic variation of traits. The functions behind the traits that have diverged during local adaptation can nevertheless be used to discuss plausible evolutionary scenarios of natural selection. Our results imply the adaptive divergence of flowering traits (flowering date, number of flowers, inflorescence size) amongst *A. majus* populations. These traits are directly linked to the reproductive success of populations and thereby often under natural selection (Meagher, 1994; Ollerton, Winfree, & Tarrant, 2011; Van Kleunen, 2007). They have great ecological significance in wild populations. Environmental constraints on the amount of resources available to plants are known to constrain flowering (Meagher & Delph, 2001). Biotic interactions between plants and pollinators are also known to modulate selection on flowering traits (Elzinga et al., 2007; Schemske & Bradshaw, 1999). Climate and seasonality differences amongst populations can also impose selection pressures on the flowering phenology (Franks, Sim, & Weis, 2007). Furthermore, populations flowering at different periods are less likely to exchange genes. Local adaptation shaping the flowering phenology can therefore have a



feedback effect on population evolutionary divergence because it can reinforce this divergence by enhancing reproductive isolation. One plausible scenario of local adaptation is that a combination of these environmental pressures acts in fact as a complex agent of divergent selection amongst *A. majus* populations.

Our results also imply that local adaptation has shaped the vegetative architecture of plants that is specific to each *A. majus* population. They show that the quantitative genetic variation of several phenotypic traits characterising the vegetative growth and development of plants (stem diameter and length, number of nodes and internode distance, number of branches) has diverged among populations as a result of local adaptation. It is difficult to identify the environmental pressures underlying this finding because several environmental parameters (vegetation cover, wind, disturbance, temperature, water availability, etc.) can affect these traits. Every combination of them might have differed and be at the origin of divergent selection pressures between *A. majus* populations. Divergence in the genetic variation underlying the shape and size of plants was already found at the level of *Antirrhinum* species but its adaptive significance was not tested for (Langlade et al., 2005). The fact that a physiological trait, the Specific Leaf Area, is also involved with the local adaptation of *A. majus* populations suggests that the resource use ecological strategy of plants differs amongst populations. This is because population differences in this trait often reflect evolutionary divergence in the relative investment of resources in rapid growth (B. Pujol et al., 2008; Wilson, Thompson, & Hodgson, 1999).

Results like ours on the germination date of *A. majus* populations - lower quantitative genetic differentiation than expected under a scenario of neutral evolution - are widely acknowledged to imply stabilizing selection amongst populations (Lamy et al., 2012). Stabilizing selection of the germination date is a surprising finding because the timing of germination is often found to have evolved in response to divergent selection between habitats and populations (Donohue et al., 2005; Kalisz, 1986; Pujol et al., 2002; Weinig, 2000). The absence of fitness trade-offs between local environments might participate to explain this lack of divergence (Silvertown, 1981). The classical hypothesis suggested by this result is that a set of selective agents, which would be constant amongst populations, might act on this trait and favour a similar timing of germination across the species geographic and ecological range. Here we argue that an alternative but not mutually exclusive biological explanation can be invoked. There is evidence for the genetic basis of the germination timing in *A. majus* (Foley & Fennimore, 1998). Our

alternative explanation nevertheless implies that the germination timing of *A. majus* populations was shaped by phenotypic plasticity in our experiment. For instance, we observed different abundance periods amongst *A. majus* populations in the wild. It is likely that phenotypic differences between wild populations that are not found in common garden conditions reflect plastic rather than genetic differences (Benoit Pujol et al., 2008). Here, we argue that the phenotypic similarity of populations in a common garden experiment might reflect their similar adjustment by means of phenotypic plasticity to the similar environment.

### **Adaptive evolution of *A. majus* populations along the altitudinal gradient**

Our results imply that the quantitative genetic basis of three of the thirteen traits under study (number of flowers, inflorescence length and germination date) was shaped by divergent selection between populations from different altitudes. Neutral genetic differentiation increased with population altitudinal differences, which means that smaller populations, less genetically diverse, and/or more reproductively isolated might be found higher on mountains (Pujol et al., 2017). This increase was however very small, so that neutral evolution is unlikely to explain the increase in quantitative genetic divergence observed for these three traits when population altitudinal differences increased. This latter increase was indeed one order of magnitude stronger than the baseline level set by neutral divergence (Figure 3). The hypothesis that the altitudinal gradient in quantitative genetic differentiation that we detected for three traits could be explained by neutral evolution therefore received poor support from our data. Contrasted environmental conditions characterise different altitudes in the geographic range of *A. majus*. In particular, we emphasize here the clear differences in rainfall, temperature and atmospheric pressure (Figure S2, Supporting Information). Other non-measured or unidentified ecological aspects (e.g., pollinator abundance, neighbouring plant community, vegetation cover) are also likely to change with altitude. These environmental factors are therefore good candidate selective agents for the observed adaptive evolutionary divergence of *A. majus* populations with altitude.

Most studies on plant adaptation to altitude report the selection of shorter plants that are faster growers at higher altitudes (Körner, 1999). There is also evidence for adaptive changes in flowering phenology and leaf traits to higher altitude environments (Körner, 1999). This is because there is generally a shorter growth season at higher altitudes, and thereby selection for fast developing plants that reproduce before the season deteriorates (Körner, 1999). Our results did not support a similar scenario of selection at play in *A. majus*. In contrast with the usual

findings, our results showed that the divergent selection of flower production and germination timing between different altitudes predominates over flowering phenology, vegetative growth, and leaf traits in *A. majus* populations. *A. majus* plants that germinate later were selected for at higher altitudes. This is likely because it allows plants to track the later arrival of suitable climatic conditions for growth at higher altitudes (Körner, 1999). *A. majus* plants with smaller inflorescences and a reduced number of flowers were also selected for at higher altitudes. This can be the by-product of selection for the reduction of the whole plant size (Gonzalo-Turpin & Hazard, 2009). Alternatively, trade-offs in the resource allocation towards vegetative reproduction might be favoured at higher altitudes at the expense of sexual reproduction (Pluess & Stöcklin, 2005). The latter hypothesis is less likely in *A. majus* because we did not detect an increase in vegetative growth, and its vegetative reproduction in the wild is not documented. Our finding implies that environmental conditions becoming more stressful at higher altitudes (Körner, 2007) selected for reduced flower production.

Detecting the quantitative genetic signature of divergent selection imposed by environments that change with altitude has implications for the research on plant adaptation to climate change (De Vilmereuil et al.). Our results provide evidence for the adaptation of *A. majus* populations to altitude in the Pyrenees, which might imply that *A. majus* successfully evolved adaptations to climate differences. The range of climate conditions in these mountains is already changing and set to change even more in the near future as a result of climate change. Conditions at high altitudes are becoming more similar to conditions from lower altitudes (Urli et al., 2014). Although signatures of past adaptive evolution have sometimes been used to evaluate the ability of populations to adapt to climate change on the long run, we argue that past adaptation does not imply that populations still have the ability to adapt. These signatures are however useful to identify potential adaptive traits, as for example flower production and germination date in *A. majus*, which might play a key role in the adaptation to climatic differences. The presence of heritable variation for adaptive traits is a more direct predictor of the ability of populations to respond to selection in the absence of specific constraints (Charmantier, Garant, & Kruuk, 2014; Kruuk, Slate, & Wilson, 2008; Pujol et al., 2018). In our approach, we quantified significant heritable variation in *A. majus* for traits potentially involved with adaptation to altitude. Under the hypothesis that no mechanisms will impede the response to selection in the wild (Pujol et al., 2018), our results suggest that *A. majus* populations might have some potential to adapt to climate change.

## Conclusion

Our findings corroborate the utility of  $Q_{ST}$ - $F_{ST}$  approaches conducted in common garden experiments to explore whether adaptive evolutionary divergence is required to explain trait quantitative genetic differentiation amongst populations. It is widely acknowledged that such an approach can be used to identify traits that might be involved with local adaptation across the geographic range of a species. Our findings also ~~participate to~~ confirm the utility of the emerging use of  $Q_{ST}$ - $F_{ST}$  comparisons as a tool identifying candidate traits involved with the adaptation to selective agents acting on multiple populations rather than unknown local features of the environment. Here, we disentangled *A. majus* local adaptation from its response to selection along altitudinal gradients. This is a pertinent tool to explore how selection by ecological conditions that are currently changing, including climate envelopes, have already shaped the quantitative genetic variation of adaptive traits in the past. Combined with population estimates of the evolutionary potential to respond to selection, such approach might prove useful for building potential evolutionary scenarios of future adaptive evolution.

## Acknowledgments

We thank Jessica Cote for helpful discussions on the methods and David Field for helpful comments on the manuscript. This project has received funding from the European Research Council (ERC) under the European Union's horizon 2020 research and innovation program (grant agreement No ERC-CoG-2015-681484-ANGI) awarded to BP. This work was supported by funding from the French "Agence Nationale de la Recherche" (ANR-13-JSV7-0002 "CAPA") to BP. This project was also supported by the ANR funded French Laboratory of Excellence projects "LABEX TULIP" and "LABEX CEBA" (ANR-10-LABX-41, ANR-10-LABX-25-01).

## Reference List

- Ahuja, I., de Vos, R. C. H., Bones, A. M., & Hall, R. D. (2010). Plant molecular stress responses face climate change. *Trends in Plant Science*, 15(12), 664-674. doi:<https://doi.org/10.1016/j.tplants.2010.08.002>
- Alberto, F., Niort, J., Derory, J., Lepais, O., Vitalis, R., Galop, D., & Kremer, A. (2010). Population differentiation of sessile oak at the altitudinal front of migration in the French Pyrenees. *Molecular Ecology*, 19(13), 2626-2639. doi:doi:10.1111/j.1365-294X.2010.04631.x

- Andalo, C., Cruzan, M., Cazettes, C., Pujol, B., Burrus, M., & Thébaud, C. (2010). Post-pollination barriers do not explain the persistence of two distinct *Antirrhinum* subspecies with parapatric distribution. *Plant Systematics and Evolution*, 286(3), 223-234. doi:10.1007/s00606-010-0303-4
- Angert, A. L., Schamske, D. W., & Geber, M. (2005). The evolution of species' distributions: reciprocal transplants across the elevation ranges of *Mimulus cardinalis* and *M. lewisii*. *Evolution*, 59(8), 1671-1684. doi:10.1554/05-107.1
- Bates, D. (2005). Fitting linear mixed models in R using the lme4 package. *R News*, 5, 27-30.
- Belkhir, K., Borsa, P., Chikhi, L., Raufaste, N., & Bonhomme, F. (2004). *Genetix version 4.05 Logiciel sous Windows pour la génétique des populations*. Montpellier Laboratoire Génome, Populations, Interactions. CNRS UMR 5000, Université de Montpellier II.
- Bonhomme, V., Picq, S., Gaucherel, C., & Claude, J. (2014). Momocs: Outline Analysis Using R. *Journal of Statistical Software*, 56, i13.
- Bradburd, G. S., Ralph, P. L., & Coop, G. M. (2013). Disentangling the effects of geographic and ecological isolation on genetic differentiation. *Evolution*, 67(11), 3258-3273. doi:doi:10.1111/evo.12193
- Byars, S. G., Papst, W., & Hoffmann, A. A. (2007). Local adaptation and cogradient selection in the alpine plant, *Poa hiemata*, along a narrow altitudinal gradient. *Evolution*, 61(12), 2925-2941. doi:doi:10.1111/j.1558-5646.2007.00248.x
- Charmantier, A., Garant, D., & Kruuk, L. E. B. (2014). *Quantitative Genetics in the Wild*. Oxford: Oxford University Press
- Chessel, D., Dufour, A. B., & Thioulouse, J. (2004). The ade4 Package – I: One-Table Methods. *R News*, 4(1), 5–10.
- De Villemereuil, P., & Gaggiotti, O. E. (2015). A new FST-based method to uncover local adaptation using environmental variables. *Methods in Ecology and Evolution*, 6(11), 1248-1258. doi:doi:10.1111/2041-210X.12418
- De Villemereuil, P., Mouterde, M., Gaggiotti, O. E., & Till-Bottraud, I. Patterns of phenotypic plasticity and local adaptation in the wide elevation range of the alpine plant *Arabis alpina*. *Journal of Ecology*, 0(0). doi:doi:10.1111/1365-2745.12955
- Debout, G., Lhuillier, E., Malé, P.-J., Pujol, B., & Thébaud, C. (2012). Development and characterization of 24 polymorphic microsatellite loci in two *Antirrhinum majus* subspecies (Plantaginaceae) using pyrosequencing technology. *Conservation Genetics Resources*, 4(1), 75-79. doi:10.1007/s12686-011-9478-4

- Donohue, K., Dorn, L., Griffith, C., Kim, E., Aguilera, A., Polisetty, C. R., . . . Galloway, L. (2005). The evolutionary ecology of seed germination of *Arabidopsis thaliana*: variable natural selection on germination timing. *Evolution*, *59*(4), 758-770. doi:10.1554/04-418
- Edelaar, P. I. M., & Björklund, M. (2011). If  $F_{ST}$  does not measure neutral genetic differentiation, then comparing it with  $Q_{ST}$  is misleading. Or is it? *Molecular Ecology*, *20*(9), 1805-1812. doi:10.1111/j.1365-294X.2011.05051.x
- Edelaar, P. I. M., Burraco, P., & Gomez-Mestre, I. (2011). Comparisons between  $Q_{ST}$  and  $F_{ST}$ —how wrong have we been? *Molecular Ecology*, *20*, 4830-4839. doi:10.1111/j.1365-294X.2011.05333.x
- Elzinga, J. A., Atlan, A., Biere, A., Gigord, L., Weis, A. E., & Bernasconi, G. (2007). Time after time: flowering phenology and biotic interactions. *Trends in ecology & evolution*, *22*(8), 432-439. doi:<https://doi.org/10.1016/j.tree.2007.05.006>
- Etterson, J. R. (2004). Evolutionary potential of *Chamaecrista fasciculata* in relation to climate change. II. Genetic architecture of three populations reciprocally planted along an environmental gradient in the great plains. *Evolution*, *58*(7), 1459-1471.
- Foley, M. E., & Fennimore, S. A. (1998). Genetic basis for seed dormancy. *Seed Science Research*, *8*(2), 173-182. doi:10.1017/S0960258500004086
- Franks, S. J., Sim, S., & Weis, A. E. (2007). Rapid evolution of flowering time by an annual plant in response to a climate fluctuation. *Proceedings of the National Academy of Sciences*, *104*(4), 1278-1282. doi:10.1073/pnas.0608379104
- Gomez-Mestre, I., Tejedo, M., & Ashley, M. (2004). Contrasting patterns of quantitative and neutral genetic variation in locally adapted populations of the Natterjack toad, *Bufo calamita*. *Evolution*, *58*(10), 2343-2352. doi:10.1554/03-671
- Gonzalo-Turpin, H., & Hazard, L. (2009). Local adaptation occurs along altitudinal gradient despite the existence of gene flow in the alpine plant species *Festuca eskia*. *Journal of Ecology*, *97*(4), 742-751. doi:10.1111/j.1365-2745.2009.01509.x
- Hangartner, S., Laurila, A., & Räsänen, K. (2011). Adaptive divergence in moor frog (*Rana arvalis*) populations along an acidification gradient: Inferences from  $Q_{ST}$ – $F_{ST}$  correlations. *Evolution*, *66*(3), 867–881. doi:10.1111/j.1558-5646.2011.01472.x
- Hoffmann, A. A., & Sgro, C. M. (2011). Climate change and evolutionary adaptation. *Nature*, *470*(7335), 479-485.
- Kalisz, S. (1986). Variable selection on the timing of germination in *Colinsia verna* (Scrophulariaceae). *Evolution*, *40*(3), 479-491. doi:doi:10.1111/j.1558-5646.1986.tb00501.x

- Karhunen, M., Ovaskainen, O., Herczeg, G., & Merilä, J. (2014). Bringing habitat information into statistical tests of local adaptation in quantitative traits: A case study of nine-spined sticklebacks. *Evolution*, *68*(2), 559-568. doi:doi:10.1111/evo.12268
- Kawecki, T. J., & Ebert, D. (2004). Conceptual issues in local adaptation. *Ecology Letters*, *7*(12), 1225-1241.
- Khimoun, A., Burrus, M., Andalo, C., Liu, Z. L., Vicédo-Cazettes, C., Thébaud, C., & Pujol, B. (2011). Locally asymmetric introgressions between subspecies suggest circular range expansion at the *Antirrhinum majus* global scale. *Journal of Evolutionary Biology*, *24*(7), 1433-1441. doi:10.1111/j.1420-9101.2011.02276.x
- Khimoun, A., Cornuault, J., Burrus, M., Pujol, B., Thebaud, C., & Andalo, C. (2013). Ecology predicts parapatric distributions in two closely related *Antirrhinum majus* subspecies. *Evolutionary Ecology*, *27*, 51-64. doi:10.1007/s10682-012-9574-2
- Kim, E., & Donohue, K. (2013). Local adaptation and plasticity of *Erysimum capitatum* to altitude: its implications for responses to climate change. *Journal of Ecology*, *101*(3), 796-805. doi:doi:10.1111/1365-2745.12077
- Körner, C. (1999). *Alpine Plant Life: Functional Plant Ecology of High Mountain Ecosystems.*: Springer-Verlag, Berlin, Germany.
- Körner, C. (2007). The use of 'altitude' in ecological research. *Trends in ecology & evolution*, *22*(11), 569-574. doi:<https://doi.org/10.1016/j.tree.2007.09.006>
- Kruuk, L. E. B., Slate, J., & Wilson, A. J. (2008). New answers for old questions: The evolutionary quantitative genetics of wild animal populations. *Annual Review of Ecology, Evolution, and Systematics*, *39*(1), 525-548.
- Lamy, J.-B., Plomion, C., Kremer, A., & Delzon, S. (2012).  $QST < FST$  As a signature of canalization. *Molecular Ecology*, *21*(23), 5646-5655. doi:10.1111/mec.12017
- Langlade, N. B., Feng, X., Dransfield, T., Copsey, L., Hanna, A. I., Thebaud, C., . . . Coen, E. (2005). Evolution through genetically controlled allometry space. *Proceedings of the National Academy of Sciences*, *102*(29), 10221-10226.
- Leinonen, T., McCairns, R. J. S., O'Hara, R. B., & Merila, J. (2013).  $QST-FST$  comparisons: evolutionary and ecological insights from genomic heterogeneity. *Nat Rev Genet*, *14*(3), 179-190.
- Leinonen, T., O'Hara, R. B., Cano, J. M., & Merila, J. (2008). Comparative studies of quantitative trait and neutral marker divergence: a meta-analysis. *Journal of Evolutionary Biology*, *21*(1), 1-17.

- Lind, M. I., Ingvarsson, P. K., Johansson, H., Hall, D., & Johansson, F. (2010). Gene flow and selection on phenotypic plasticity in an island system of *Rana temporaria*. *Evolution*, 65, 684-697. doi:10.1111/j.1558-5646.2010.01122.x
- Luquet, E., Léna, J. P., Miaud, C., & Plénet, S. (2014). Phenotypic divergence of the common toad (*Bufo bufo*) along an altitudinal gradient: evidence for local adaptation. *Heredity*, 114, 69. doi:10.1038/hdy.2014.71
- Mantel, N. (1967). The detection of disease clustering and a generalized regression approach. *Cancer Research*, 27(2 Part 1), 209-220.
- McKay, J. K., Bishop, J. G., Lin, J.-Z., Richards, J. H., Sala, A., & Mitchell-Olds, T. (2001). Local adaptation across a climatic gradient despite small effective population size in the rare sapphire rockcress. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 268(1477), 1715-1721. doi:10.1098/rspb.2001.1715
- McKay, J. K., & Latta, R. G. (2002). Adaptive population divergence: markers, QTL and traits. *Trends in ecology & evolution*, 17(6), 285-291.
- Meagher, T. R. (1994). The quantitative genetics of sexual dimorphism in *Silene Latifolia* (Caryophyllaceae) .2. Response to sex-specific selection. *Evolution*, 48(4), 939-951.
- Meagher, T. R., & Delph, L. F. (2001). Individual flower demography, floral phenology and floral display size in *Silene latifolia*. *Evolutionary Ecology Research*, 3(7), 845-860.
- Merila, J., & Crnokrak, P. (2001). Comparison of genetic differentiation at marker loci and quantitative traits. *Journal of Evolutionary Biology*, 14(6), 892-903.
- Muir, A. P., Biek, R., Thoma, s. R., & Mable, B. K. (2014). Local adaptation with high gene flow: temperature parameters drive adaptation to altitude in the common frog (*Rana temporaria*). *Molecular Ecology*, 23(3), 561-574. doi:doi:10.1111/mec.12624
- Ollerton, J., Winfree, R., & Tarrant, S. (2011). How many flowering plants are pollinated by animals? *Oikos*, 120(3), 321-326. doi:10.1111/j.1600-0706.2010.18644.x
- Ovaskainen, O., Karhunen, M., Zheng, C., Arias, J. M. C., & Merilä, J. (2011). A new method to uncover signatures of divergent and stabilizing selection in quantitative traits. *Genetics*, 189(2), 621-632. doi:10.1534/genetics.111.129387
- Palo, J. U., O'Hara, R. B., Laugen, A. T., Laurila, A., Primmer, C. R., & Merilä, J. (2003). Latitudinal divergence of common frog (*Rana temporaria*) life history traits by natural selection: evidence from a comparison of molecular and quantitative genetic data. *Molecular Ecology*, 12(7), 1963-1978. doi:doi:10.1046/j.1365-294X.2003.01865.x
- Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., . . . Cornelissen, J. H. C. (2016). *Corrigendum to: New handbook for standardised*


- measurement of plant functional traits worldwide. *Australian Journal of Botany*, 64(8), 715-716. doi:[https://doi.org/10.1071/BT12225\\_CO](https://doi.org/10.1071/BT12225_CO)
- Pluess, A. R., & Stöcklin, J. (2005). The importance of population origin and environment on clonal and sexual reproduction in the alpine plant *Geum reptans*. *Functional Ecology*, 19(2), 228-237. doi:doi:10.1111/j.0269-8463.2005.00951.x
- Pujol, B., Archambeau, J., Bontemps, A., Lascoste, M., Marin, S., & Meunier, A. (2017). Mountain landscape connectivity and subspecies appurtenance shape genetic differentiation in natural plant populations of the snapdragon (*Antirrhinum majus* L.). *Botany Letters*, 164(2), 111-119. doi:10.1080/23818107.2017.1310056
- Pujol, B., Blanchet, S., Charmantier, A., Danchin, E., Facon, B., Marrot, P., . . . Winney, I. (2018). The missing response to selection in the wild. *Trends in ecology & evolution*, 33(5), 337-346. doi:10.1016/j.tree.2018.02.007
- Pujol, B., Gigot, G., Laurent, G., Pinheiro-Kluppel, M., Elias, M., Hossaert-McKey, M., & McKey, D. (2002). Germination ecology of cassava (*Manihot esculenta* Crantz, Euphorbiaceae) in traditional agroecosystems: Seed and seedling biology of a vegetatively propagated domesticated plant. *Economic Botany*, 56(4), 366-379.
- Pujol, B., Salager, J. L., Beltran, M., Bousquet, S., & McKey, D. (2008). Photosynthesis and leaf structure in domesticated cassava (Euphorbiaceae) and a close wild relative: have leaf photosynthetic parameters evolved under domestication? *Biotropica*, 40(3), 305-312.
- Pujol, B., Wilson, A. J., Ross, R. I. C., & Pannell, J. R. (2008). Are QST-FST comparisons for natural populations meaningful? *Molecular Ecology*, 17(22), 4782-4785.
- R-Core-Team. (2018). *R: A Language and Environment for Statistical Computing*. : R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- Raeymaekers, J. A. M., Van Houdt, J. K. J., Larmuseau, M. H. D., Geldof, S., & Volckaert, F. A. M. (2007). Divergent selection as revealed by  $P_{ST}$  and QTL-based  $F_{ST}$  in three-spined stickleback (*Gasterosteus aculeatus*) populations along a coastal-inland gradient. *Molecular Ecology*, 16(4), 891-905.
- Ringbauer, H., Kolesnikov, A., Field, D. L., & Barton, N. H. (2018). Estimating barriers to gene flow from distorted isolation by distance patterns. *Genetics*. doi:10.1534/genetics.117.300638
- Roff, D. A. (1997). *Evolutionary Quantitative Genetics*. New York: Chapman and Hall.
- Rousset, F. (1997). Genetic differentiation and estimation of gene flow from F-Statistics under isolation by distance. *Genetics*, 145(4), 1219-1228.

- Schemske, D. W., & Bradshaw, H. D. (1999). Pollinator preference and the evolution of floral traits in monkeyflowers (*Mimulus*). *Proceedings of the National Academy of Sciences*, 96(21), 11910-11915. doi:10.1073/pnas.96.21.11910
- Silvertown, J. W. (1981). Seed size, life span, and germination date as coadapted features of plant life history. *The American Naturalist*, 118(6), 860-864. doi:10.1086/283876
- Spitze, K. (1993). Population-Structure in *Daphnia obtusa* - Quantitative genetic and allozymic variation. *Genetics*, 135(2), 367-374.
- Urli, M., Delzon, S., Eyermann, A., Couallier, V., García-Valdés, R., Zavala, M. A., & Porté, A. J. (2014). Inferring shifts in tree species distribution using asymmetric distribution curves: a case study in the Iberian mountains. *Journal of Vegetation Science*, 25(1), 147-159. doi:doi:10.1111/jvs.12079
- Van Kleunen, M. (2007). Adaptive genetic differentiation in life-history traits between populations of *Mimulus guttatus* with annual and perennial life-cycles. *Evolutionary Ecology*, 21, 185-199.
- Weinig, C. (2000). Differing selection in alternative competitive environments: shade-avoidance responses and germination timing. *Evolution*, 54(1), 124-136. doi:10.1554/0014-3820(2000)054[0124:DSIACE]2.0.CO;2
- Whibley, A. C., Langlade, N. B., Andalo, C., Hanna, A. I., Bangham, A., Thebaud, C., & Coen, E. (2006). Evolutionary paths underlying flower color variation in *Antirrhinum*. *Science*, 313(5789), 963-966. doi:10.1126/science.1129161
- Whitlock, M. C. (2008). Evolutionary inference from QST. *Molecular Ecology*, 17(8), 1885-1896. doi:doi:10.1111/j.1365-294X.2008.03712.x
- Whitlock, M. C. (2011).  $G'_{ST}$  and  $D$  do not replace  $F_{ST}$ . *Molecular Ecology*, 20(6), 1083-1091. doi:10.1111/j.1365-294X.2010.04996.x
- Whitlock, M. C., & Gilbert, K. J. (2012). QST in a hierarchically structured population. *Molecular Ecology Resources*, 12(3), 481-483. doi:doi:10.1111/j.1755-0998.2012.03122.x
- Whitlock, M. C., & Guillaume, F. (2009). Testing for spatially divergent selection: Comparing QST to FST. *Genetics*, 183(3), 1055-1063. doi:10.1534/genetics.108.099812
- Wilson, P. J., Thompson, K. E. N., & Hodgson, J. G. (1999). Specific leaf area and leaf dry matter content as alternative predictors of plant strategies. *New Phytologist*, 143(1), 155-162. doi:undefined


## Supplementary information

**TABLE S1** Description of the parental plant populations and characteristics of study system: number of families (*Nfam*), vegetative sampling size (*Nv*), reproductive sampling size (*Nr*).


Acronym	Latitude	Longitude	Location	Altitude (m)	Subspecies	Description	<i>Nfam</i>	<i>Nv</i>	<i>Nr</i>
BAG	43.10	2.98	Bages	6	<i>pseudomajus</i>	Dunes on seaside (rocky / herbaceous)	40	67	45
BAN	42.49	3.12	Banyuls-sur-Mer	61	<i>pseudomajus</i>	Roadside bank (rocky)	32	54	33
THU	42.64	2.72	Thuir	130	contact	Roadside bank (herbaceous)	34	60	40
LAG	43.09	2.58	Lagrasse	149	contact	Roadside bank (rocky / herbaceous)	32	55	44
BES	42.21	2.67	Besalú	195	<i>pseudomajus</i>	Stone walls in village	39	69	45
LUC	42.97	2.26	Luc-sur-Aude	227	<i>striatum</i>	Roadside bank and river-side bank (rocky)	19	29	18
RIP	42.21	2.20	Ripoll	709	<i>pseudomajus</i>	Roadside bank (herbaceous)	9	16	11
LYS	42.83	2.20	'Pierre-Lys' gorge	713	<i>striatum</i>	Roadside bank (rocky / herbaceous)	32	53	29
CAL	42.10	1.83	Berga	838	<i>pseudomajus</i>	Roadside bank (herbaceous)	42	69	28
PAR	42.31	2.20	Pardines	1118	<i>pseudomajus</i>	Roadside bank (herbaceous)	32	58	36
SAL	42.23	1.74	Saldes	1126	contact	Banks in pasture (herbaceous)	30	55	26
MIJ	42.73	2.04	Mijanès	1347	<i>striatum</i>	Roadside bank (herbaceous)	10	18	10
MON	42.51	2.12	Mont-Louis citadelle	1564	<i>striatum</i>	Stone walls on fortifications	21	34	15
<i>All populations</i>							<i>372</i>	<i>637</i>	<i>380</i>



**FIGURE S1** Spring and summer average temperatures and rainfall in the 13 populations. Population average temperature (a) and average rainfall (b) for the spring (triangles) and summer (dots) seasons as a function of altitude. Bioclimatic data was extracted from the *WorldClim* database ([www.worldclim.org](http://www.worldclim.org)).



**FIGURE S1bis** Annual average temperatures and rainfall in the 13 populations. Population average temperature (a) and average rainfall (b) as a function of altitude. Bioclimatic data was extracted from the *WorldClim* database ([www.worldclim.org](http://www.worldclim.org)).



**FIGURE S2** Linear regression lines of population pairwise average temperature (orange) and average rainfall (black) for a year with population pairwise differences in altitude (m). Regression equations and level of significance ( $P$ ), obtained by Mantel test, are presented in figure and its 95% confidence interval is indicated by the shaded area. Bioclimatic data was extracted from the *WorldClim* database ([www.worldclim.org](http://www.worldclim.org)).

1 **TABLE S2** Summary of quantitative traits according to the sampling site. The table exhibits the average of phenotypic values of the common-garden  
 2 experiment for all combined populations and different populations. All mean values are accompanied by their standard error.

Phenotypic Traits	All populations	BAG	BAN	THU	LAG	BES	LUC	RIP	LYS	CAL	PAR	SAL	MIJ	MON
Germination date	197.45 ± 0.45	197.51 ± 1.38	196.31 ± 1.37	195.2 ± 1.32	197.16 ± 1.62	196.22 ± 1.18	194.43 ± 1.48	194.19 ± 2.55	200.06 ± 1.7	198.72 ± 1.74	198.84 ± 1.43	198.17 ± 1.48	200.44 ± 2.94	198.47 ± 1.83
Flowering date	241.73 ± 0.66	240.33 ± 1.87	245.73 ± 2.47	237.4 ± 1.24	243.52 ± 2.86	244.09 ± 1.81	237.94 ± 2.15	236.5 ± 1.34	239.25 ± 2.55	238.64 ± 1.3	243.83 ± 1.84	250.04 ± 3.12	240.9 ± 3.18	236.4 ± 1.05
Time to flowering	47.97 ± 0.48	46.56 ± 1.21	51 ± 2.26	45.5 ± 0.86	48.45 ± 1.96	49.40 ± 1.14	44.67 ± 1.18	48.5 ± 1.34	45.11 ± 1.28	45.96 ± 0.83	48.83 ± 1.66	56.36 ± 2.52	47 ± 1.11	43.8 ± 1.3
Diameter (mm)	3.73 ± 0.04	3.59 ± 0.10	4.14 ± 0.15	4.03 ± 0.12	3.59 ± 0.11	3.65 ± 0.13	3.42 ± 0.15	3.98 ± 0.19	3.37 ± 0.14	3.85 ± 0.11	3.71 ± 0.14	3.82 ± 0.13	3.38 ± 0.18	3.76 ± 0.16
Vegetative Length (cm)	26.83 ± 0.34	23.4 ± 0.79	35.74 ± 1.18	30.63 ± 0.71	24.32 ± 0.89	28.97 ± 1.18	21.66 ± 0.85	30.91 ± 2.1	21.68 ± 0.88	27.69 ± 1.02	28.89 ± 1.28	27.01 ± 1.01	22 ± 1.52	19.95 ± 0.73
Nodes	13.21 ± 0.16	12.46 ± 0.41	15.07 ± 0.52	13.33 ± 0.55	11.07 ± 0.37	14.67 ± 0.52	12.69 ± 0.50	16.38 ± 1.02	11.62 ± 0.43	13.7 ± 0.42	12.95 ± 0.41	15.89 ± 0.65	10.61 ± 0.65	10 ± 0.41
Leaves	124.72 ± 2.12	133.55 ± 6.35	134.28 ± 7.73	122.82 ± 6.5	113.82 ± 5.74	143.19 ± 8.6	123.41 ± 11.29	141 ± 12.56	107.42 ± 5.67	122.23 ± 4.89	109.95 ± 6.54	151.36 ± 7.47	95.72 ± 10.7	98.5 ± 5.82
Branches	16.77 ± 0.29	16.97 ± 0.75	19.94 ± 1.21	16.9 ± 0.73	15.02 ± 0.78	19.71 ± 1.19	16.24 ± 1.76	19.5 ± 1.8	14.11 ± 0.77	15.94 ± 0.77	15.84 ± 0.9	19.27 ± 0.88	13.22 ± 1.55	12.35 ± 0.67
Total Length (cm)	45.18 ± 0.68	43.74 ± 2.19	54.49 ± 2.02	52.29 ± 1.57	43.84 ± 1.72	45.94 ± 1.95	35.03 ± 1.65	48.49 ± 3.59	38.96 ± 2.22	49 ± 2.46	51.14 ± 1.83	41.28 ± 1.84	36.66 ± 1.99	25.18 ± 1.47
Inflorescence length (cm)	16.82 ± 0.51	20.00 ± 1.77	16.42 ± 1.68	21.92 ± 1.6	18.58 ± 1.47	15.92 ± 1.36	11.83 ± 1.57	15.15 ± 2.87	15.03 ± 1.95	17.52 ± 1.89	18.47 ± 1.51	12.79 ± 1.47	11.57 ± 2.12	6.86 ± 1.1
Flowers	15.95 ± 0.49	18.09 ± 1.70	15.48 ± 1.54	18.44 ± 1.34	16.75 ± 1.41	17.3 ± 1.43	14.78 ± 2.42	14.55 ± 2.86	12.55 ± 1.46	15.75 ± 1.72	18.77 ± 1.73	13.96 ± 1.46	10.9 ± 2.09	6.36 ± 1.1
Internode Length (cm)	2.09 ± 0.02	1.95 ± 0.07	2.43 ± 0.08	2.43 ± 0.07	2.26 ± 0.08	2.01 ± 0.07	1.76 ± 0.09	1.9 ± 0.08	1.92 ± 0.08	2.06 ± 0.07	2.24 ± 0.08	1.75 ± 0.06	2.11 ± 0.13	2.07 ± 0.09
SLA (cm <sup>2</sup> /g)	176.08 ± 1.96	173.29 ± 4.22	178.44 ± 5.16	179.55 ± 6.09	182.1 ± 10.37	194.48 ± 5.75	161.24 ± 5.94	165.9 ± 10.9	179.53 ± 8.04	171.29 ± 4.93	189.06 ± 6	162.62 ± 6.58	178.74 ± 14.3	144.58 ± 5.93

3

4 **TABLE S3** Differentiation for phenotypic traits among the 13 populations of *A. majus* grown in our common-garden experiment. Values for the within-  
 5 population genetic variance ( $V_w$ ), among-population genetic variance ( $V_b$ ), heritability ( $h^2$ ) and observed (Obs.)  $Q_{ST}$  are given.  $F_{ST}$  overall populations was  
 6 0.107 ( $P < 0.001$ ).  $P$ -values indicate statistical differences between observed and expected values of  $Q_{ST}-F_{ST}$  under a neutrality hypothesis. The 95% CI of  $Q_{ST}$   
 7 ( $Q_{ST} .CI$ ) and the  $P$  values of the  $Q_{ST}-F_{ST}$  contrast are based on  $10^5$  bootstrap replicates, following Whitlock & Guillaume (2009). Significant values are indicated  
 8 in bold. The degrees of freedom used in the bootstrapping procedures are 12 for the among-population component ( $V_b$ ) and are given in this table for the  
 9 within-population component (df  $V_w$ ).



Traits	$V_w$	df. $V_w$	$V_b$	$h^2$	$Q_{ST}$ -obs	$Q_{ST}$ -CI	$P$ -right	$P$ -left
Germination date	23.0964	356	0	0.36	0	NA	1	<b>0</b>
Flowering date	5.3673	259	9.828	0.07	0.48	0.247-0.644	<b>0</b>	1
Time to flowering	16.2058	259	7.4606	0.40	0.19	0.077-0.312	0.039	0.961
Diameter	0.0559	359	0.035	0.12	0.24	0.102-0.383	<b>0.003</b>	0.997
Vegetative Length	1.8973	359	19.9007	0.07	0.84	0.660-0.912	<b>0</b>	1
Nodes	3.1459	359	3.285	0.47	0.34	0.160-0.514	<b>0</b>	1
Leaves	0	359	217.3899	0	1	NA	NA	NA
Branches	1.7905	356	5.0482	0.08	0.59	0.332-0.736	<b>0</b>	1
Total Length	15.5028	266	57.5775	0.23	0.65	0.403-0.787	<b>0</b>	1
Inflorescence length	10.5535	263	10.7657	0.23	0.34	0.155-0.507	<b>0</b>	1
Flowers	6.8182	263	5.8013	0.16	0.30	0.133-0.462	<b>0</b>	1
Internode Length	0.0641	359	0.0428	0.42	0.25	0.107-0.397	<b>0</b>	1
SLA	146.6081	359	108.8237	0.13	0.27	0.118-0.422	<b>0</b>	1

10

11

12