

1 Distribution of iridescent colours in hummingbird communities results
2 from the interplay between selection for camouflage and communication

3 Hugo Gruson^{*a}, Marianne Elias^b, Juan L. Parra^c, Christine Andraud^d, Serge Berthier^e, Claire
4 Doutrelant^a, and Doris Gomez^{a,e}

5 ^aCEFE, Univ Montpellier, CNRS, Univ Paul Valéry Montpellier 3, EPHE, IRD, Montpellier, France

6 ^bISYEB, CNRS, MNHN, Sorbonne Université, EPHE, 45 rue Buffon CP50, Paris, France

7 ^cGrupo de Ecología y Evolución de Vertebrados, Instituto de Biología, Universidad de Antioquia,
8 Medellín, Colombia

9 ^dCRC, MNHN, Ministère de la Culture et de la Communication, CNRS, Paris, France

10 ^eINSP, Sorbonne Université, CNRS, Paris, France

11 **Abstract**

12 Identification errors between closely related, co-occurring, species may lead to misdirected social interactions such
13 as costly interbreeding or misdirected aggression. This selects for divergence in traits involved in species identification
14 among co-occurring species, resulting from character displacement. On the other hand, predation may select for crypsis,
15 potentially leading co-occurring species that share the same environment and predators to have a similar appearance.
16 However, few studies have explored how these antagonistic processes influence colour at the community level. Here,
17 we assess colour clustering and overdispersion in ~~multiple hummingbird communities~~ 189 hummingbird communities,
18 tallying 112 species, across Ecuador and ~~identify the processes suggest possible evolutionary mechanisms~~
19 controlling for species phylogenetic relatedness. In hummingbirds, most colours are iridescent structural colours, defined
20 as colours that change with the illumination or observation angle. Because small variations in the underlying structures
21 can have dramatic effects on the resulting colours and because iridescent structures can produce virtually any hue and
22 brightness, we expect iridescent colours to respond finely to selective pressures. Moreover, we predict that hue angular
23 dependence – a specific aspect of iridescent colours – may be used as an additional channel for species recognition.
24 In our hummingbird assemblages in Ecuador, we find support for colour overdispersion in ~~specific body ventral and~~
25 facial patches at the community level even after controlling for the phylogeny, especially on iridescence-related traits,
26 suggesting character displacement among co-occurring species. We also find colour clustering at the community level
27 on dorsal patches, suspected to be involved in camouflage, suggesting that the same cryptic colours are selected among
28 co-occurring species.

*hugo.gruson@normalesup.org

29 **Keywords:** Reproductive Character Displacement, Agonistic Character Displacement, Camouflage, Structural Colours,
30 Angle-Dependent Colouration, Community structure, Ecuador

31

32 Colour is a complex communication channel widespread among various taxa and involved in many ecological and
33 evolutionary processes [1]. It can be described by multiple variables, including hue (colour in its common sense, such as
34 red, green, blue, etc.) and brightness (average level of grey of a colour, i.e. whether the object is light or dark). Colours
35 can be produced by two non-mutually exclusive means: pigmentary colours are produced by the selective absorption of
36 incoming light by pigments, while structural colours are produced by the interaction of incoming light with nanostructures,
37 causing diffraction, interferences or scattering [2]. Among structural colours, iridescent colours are characterised by a shift
38 in hue with changes in illumination or observation angle [3]. Iridescent colours are found in many bird families such as
39 Anatidae (ducks) Phasianidae (fowls), Sturnidae (starlings), or Trochilidae (hummingbirds), and thought to be involved
40 in numerous adaptations [4]. But evolution of iridescent colours at the community level remains poorly understood. Yet,
41 ~~they may display evolutionary patterns that differ from~~ evolutionary patterns of iridescent colours, which remain poorly
42 studied and understood, may differ from that of non-iridescent colours. Indeed, as opposed to other types of colours,
43 iridescent colours can produce virtually any hue and are expected to respond more readily and finely to selection, because
44 large changes of hue can be achieved by small changes in the underlying structures [5]. They can also result in directional
45 colours only seen at specific angles, as well as highly reflective colours [6].

46 Because colours are involved in many different ecological processes, they are subject to multiple selection pressures,
47 often with opposite effects [7]. Colour may indeed increase or decrease detectability of an animal depending on the colour
48 contrast with its surroundings. In particular, colour can reduce predation risk via crypsis or aposematism or serve as a
49 means of species identification. In this case, two opposite evolutionary forces act on colours: (i) On the one hand, species
50 living in the same environment are likely experiencing similar selective pressures, such as predation. The environment
51 is characterised by ambient light and vegetation, which both influence greatly which colours are poorly detectable and
52 which colours are highly detectable [8, 9]. We thus expect co-occurring species to harbour the same, poorly detectable,
53 colours as this would decrease the risk of being detected by predators, thereby causing a clustering pattern in colouration
54 at the community level, all else being equal. This colour clustering can result from convergence between sympatric species
55 (evolutionary process), from environmental filtering (ecological process), i.e. species sorting locally according to the traits
56 they harbour, or a mixture of the two (detailed in table 1). (ii) On the other hand, sympatric closely-related species
57 are more likely to face problems of species recognition, eventually resulting in reproductive interference - a phenomenon
58 where an individual courts or mates with individuals of another species, producing no offspring or low fertility hybrids,
59 leading to costly interbreeding [10]. Species misidentification can also lead to misdirected aggression and costly fighting
60 when individuals compete over resources or territories. Hence, any feature that would enhance species recognition is
61 expected to be selected for. In this context, closely related species living in sympatry should be under strong selective
62 pressure to diverge in traits involved in communication, if divergence enhances species recognition. Divergence can result
63 from a process called character displacement (RCD for reproductive character displacement, ACD for agonistic character

64 displacement; evolutionary process) [11–13] or from species sorting (ecological process). For ACD, it is worth noting that
65 traits are expected to diverge only in case of moderate ecological competition, whereas they should converge in case of
66 high competition [13, 14]. Multiple empirical studies have shown character displacement for songs (e.g. Gerhardt [15] in
67 frogs and Grant and Grant [16] in birds), or olfactory signals [17]. However, fewer studies have looked at divergence in
68 colour patterns (but see Sætre et al. [18], Naisbit et al. [19], Lukhtanov et al. [20], Martin et al. [21], Doutrelant et al. [22],
69 and Hemingson et al. [23]). Almost all these studies were at the species level, and at best involved comparison between
70 closely related species. Many of them also did not use objective spectrometry measurements and instead relied on human
71 vision, which did not allow them to analyse colours as perceived by the intended receiver, in the case of this study: birds
72 [24–27] .

73 In birds, it has been ~~showed~~shown that colouration is under different selective pressures depending on the body patch
74 location: dorsal patches, which are exposed to aerial predators, are mainly involved in camouflage while ventral and facial
75 patches are mainly involved in communication [7, 28]. In this study, we test this hypothesis for iridescent colours at
76 the community level by looking at phenotypic structure in hummingbird local assemblages across different body parts.
77 Hummingbirds are an interesting study system to test this hypothesis as various published accounts of sexual displays and
78 aggressive encounters among hummingbirds have made clear that certain feather patches such as the crown and throat are
79 consistently used during these displays [29–32]. On the other hand, colours displayed on the dorsal side of hummingbirds
80 tend to resemble background colours and thus have been suggested to be cryptic [33]. Accordingly, we predict that co-
81 occurring hummingbird species should display similar hues on dorsal patches, leading to phenotypic clustering of hues
82 (i.e. co-occurring species are more similar than expected by chance, prediction 1) and different hues on ventral patches,
83 resulting in a phenotypic overdispersion pattern (i.e. co-occurring species are more dissimilar than expected by chance,
84 prediction 2). For brightness, we can formulate two alternative predictions: on the one hand, it might evolve in the same
85 way as hue, also because of reproductive character displacement and selection for camouflage, leading to the same outcome
86 as for hue (prediction 3, equivalent to predictions 1 and 2 but for brightness). On the other hand, because brightness level
87 positively correlates with signal conspicuousness, poorly detectable signals have similar brightness, and highly detectable
88 signals have similar brightness. Hence, we may instead expect that species co-occurring should converge for brightness on
89 all patches (prediction 3bis) if the same patches are involved in the same ecological process (communication or camouflage).

90 Compared to other types of colouration, iridescent colours might enable species recognition on another dimension in
91 the sensory space. Two species can have the same hue or brightness at a given angle but can differ at another angle,
92 via an additional variable we call "hue shift". Because hue shift cannot be seen at long distances, it may allow species
93 to diverge without interfering with camouflage against predators [4, 34]. Accordingly, we predict overdispersion for hue
94 shift not only on ventral patches, but also on dorsal patches (prediction 4). However, hue shift is often highly correlated
95 with hue due to the optics underlying iridescence (Dakin and Montgomerie [35] for example reported $R^2 \geq 0.95$ for the
96 correlation between hue and hue shift). We test this correlation with the data from this article and discuss how it may
97 impact our results.

98 At the community level, we predict that community colour volume (also known as functional richness $FRic$ in functional

99 ecology [36]) and brightness range increase with species richness more than expected in a random species assemblage (null
100 model) because co-occurring species would use different colours (hue or brightness) (prediction 5).

101 Here we test our five predictions by quantifying both iridescent and non-iridescent colours of 189 hummingbird assem-
102 blages in Ecuador that include 112 species and span a large variety of habitats, and by assessing the phenotypic structure
103 (clustering, random distribution, overdispersion of colours) and investigate the underlying processes by taking into account
104 species phylogenetic relatedness within these assemblages. Comparing the uncorrected and the phylogenetically-corrected
105 phenotypic structure of hummingbird communities will allow us to identify which mechanisms (character displacement,
106 species sorting with mutual exclusion of similar species, environmental filtering; as detailed in table 1) underlie the com-
107 munity structure of iridescent colours in hummingbirds.

108 **Materials and methods**

109 All scripts and data used to produce the results and figures from this article are available at [https://doi.org/10.5281/
110 zenodo.3355444](https://doi.org/10.5281/zenodo.3355444)

111 **Community data**

112 Hummingbirds are particularly suited as a study system to explore the possible effect of reproductive character displace-
113 ment on iridescent colours because (i) they display a large variety of hues [37] and all species harbour some iridescent
114 patches, many of which have a very strong angular dependence, rapidly shifting from e.g. pink to green or black [38,
115 39] (but note that many hummingbirds species also have non-iridescent, pigmentary, patches), (ii) they belong to a very
116 speciose family whose phylogeny is well established and readily available [40, 41], (iii) they live only in the Americas,
117 especially in the tropics where numerous species can coexist locally [37] (iv) there is an extensive documentation of hy-
118 bridisation between co-occurring species (see for example [42, 43] for our region of interest), which creates the perfect
119 opportunity to study reproductive interference and (v) almost all species are available in museum collections and their
120 colour can be objectively measured using spectrometric measurements [44].

121 Presence/absence data for hummingbird assemblages at 189 sites in Ecuador (see map in fig. S3) were compiled
122 from data in peer-reviewed papers and reports from environmental organisations [45]. These sites cover a large variety
123 of elevation ranges (fig. S3) and habitats [45, 46]. This dataset was previously thoroughly reviewed by comparing the
124 observations with the known elevational and geographical ranges of each species [46] and includes observations of 112 of
125 the 132 hummingbirds species found in Ecuador [47].

126 **Colour measurements and analyses**

127 For each one of the 112 species, we borrowed one adult male in good condition from either the Museum National d'Histoire
128 Naturelle (MNHN) in Paris or the Musée des Confluences, in Lyon (full list in Online Supplementary Information). studies
129 show that even low sampling per species (as low as one individual/species in some cases) can accurately capture colour

130 characteristics of the species [48]. Additionally, preliminary analyses on an independent dataset of 834 points across
131 18 hummingbird species, with up to 5 individuals measured by species, showed that intraspecific coefficient of variation
132 (standard deviation divided by the average) of hue is very low (1.69%) but could be higher for brightness (23.18%)
133 (detailed values for each species in table S3). When comparing intra- to interspecific variation, intraspecific however
134 always remains negligible compared to interspecific variation (intraclass coefficient reported in table S3). We ensured that
135 the specimen colouration was representative of the other specimens available in the collections to the human eye. When
136 multiple subspecies were living in the area where presence was recorded, we randomly picked one of them. Whenever
137 possible, we picked specimens collected in Ecuador (88 % of the cases), or when not available in neighbouring
138 countries, such as Colombia or North Peru (11 % of the cases), as to minimise the effect of regional variability in colour.

139 We consistently took spectral reflectance measurements on the 8 following patches (described in fig. S1): crown,
140 back, rump, tail, throat, breast, belly, wing. We also made additional measurements on patches that visually differed in
141 colouration from these 8 main ones, as in Gomez and Théry [7] and Doutrelant et al. [22].

142 We measured reflectance using a setup similar to Meadows et al. [49], relying on the use of two separate optical fibres.
143 Light was conducted from an Oceanoptics DH-2000 lamp emitting over the 300-700 nm range of wavelengths to which
144 birds are sensitive [50] to the sample through an illuminating FC-UV200-2-1.5 x 100 optical fibre (named illumination
145 fibre). Light reflected by the sample was then collected by a second identical optical fibre (named collection fibre) and
146 conducted toward an Oceanoptics USB4000 spectrophotometer (used with the SpectraSuite 2.0.162 software). This setup
147 allows for a precise independent rotation of the illumination and the collection fibres, necessary for the measurements of
148 iridescent colours [6]. For more details about the measurement conditions as recommended in White et al. [51], see the
149 supplementary materials (ESM).

150 For every patch, we recorded a first reflectance spectrum at the position of the fibres which maximised total reflectance.
151 To measure hue angle dependency (iridescence), we then moved both fibres 10° away from the previous position and
152 recorded a second spectrum, as in Meadows et al. [52]. More recent measurement methods revealed that it would be more
153 accurate to keep the angular span between the illumination and collection fibres constant [53]. We however confirmed
154 that this did not impact our results by running our analyses once with all data and once with only data at a given angular
155 span (which represented 94% of the total data). All measurements were performed in a dark room with temperature
156 control. Recorded spectra were normalised by an Avantes WS-1 white standard and a measurement with the lamp shut
157 down (dark reference) and integration times were determined for each sample as to maximise the intensity of the signal
158 without saturating the spectrometer.

159 Final values were averaged over 5 consecutive measurements and spectra were smoothed using a loess algorithm and
160 interpolated every 1 nm and negative values were set to zero using the R package `pavo` [54].

161 We analysed spectra using Endler and Mielke [55] model with relative quantum catches Q_i (without Fechner's law).
162 All birds are tetrachromats and can see light with wavelengths from 300 to 700 nm, which includes ultra-violet light (UV)
163 [56]. But different bird species vary in their sensitivity [57]: some are UV-sensitive (UVS) while others are violet-sensitive
164 (VS). Literature on colour vision in hummingbirds suggests that both types are found within the family (see Chen and

165 Goldsmith [50] and Herrera et al. [58] for UVS species and Ödeen and Håstad [59] for VS species). Because we did not
166 have enough information to compute ancestral states and vision type for all species in our study and because it was
167 found to have little influence in previous studies [7, 28], we ran our analyses as if all species were VS, using the spectral
168 sensitivities of a typical VS bird, *Puffinus pacificus* [60], whose photoreceptor absorbances match closely those reported for
169 hummingbirds [59]. We used different illuminants defined in Endler [8], depending on the habitat of the species described
170 in Stotz et al. [61] (detailed in SI): "large gaps" illumination was used for species living in the canopy while "forest shade"
171 was used for species living in the understory. Hue was a tridimensional variable defined by the position (x , y and z) of the
172 reflectance spectrum in the tetrahedron representing bird colour vision space [55] and brightness was defined as in Endler
173 and Mielke [55] (perceived intensity of colour, also sometimes referred to as luminance). We ensured that all indices were
174 repeatable (table S1) by measuring twice the same individual and patch on 20 patches and computing the intra-class
175 coefficient (ICC) with the rptR R package [62]. We add another variable to describe iridescence: hue shift, defined as
176 the difference between hue at maximum reflectance and hue at 10° away from maximum reflectance, in a similar fashion
177 to Dakin and Montgomerie [35]. Because it is the difference of two tridimensional variables (hue at the position where
178 reflectance was maximum and hue at 10° away), hue shift is tridimensional as well. Dakin and Montgomerie [35] found a
179 high correlation between hue and hue shift at the intraspecific level in the peacock *Pavo cristatus*, we also report a high
180 correlation at the interspecific level in hummingbirds by performing a linear regression in \mathbb{R}^3 between hue and hue shift
181 ($R^2 = 0.51$, $F(3; 1372) = 469.7$, $p < 0.0001$). New measurement methods have since been developed and propose a new
182 definition for hue shift which is not correlated to hue but they were not available at the time of this study [53].

183 We analysed the colour volume for each species by measuring the convex hull volume of all colour patches on the
184 bird, as suggested in Stoddard and Prum [63]. We compared the relationship between the colour volume of a community
185 and the number of species within this community relative to a null model (prediction 5) obtained by creating random
186 assemblages from a species pool containing all species from all communities. In other words, actual assemblages are
187 compared to fictional assemblages with exactly the same number of species but no abiotic or biotic constraints on the
188 species composition.

189 However, the colour volume does not take into account the patch location on the bird body, raising several concerns.
190 First, two species could use the same colour but at different places on their body. They would then look different to
191 an observer but not identified as such in this analysis. Additionally, we expect different evolutionary signals on different
192 patches, that could even each other out, and blur the outcome at the bird level. For these reasons, we also performed
193 our analyses separately for each one of the following eight patches: crown, back, rump, tail, throat, breast, belly, wing
194 (locations shown in fig. S1).

195 **Trochilidae phylogeny and comparative analyses**

196 A distribution of 100 phylogenetic trees of the Trochilidae family was downloaded from birdtree.org [40] to take into
197 account phylogenetic uncertainty in the comparative analyses [64]. The 112 species included in this study constitute a
198 fairly even sampling of the hummingbird phylogeny (fig. S2).

199 We used the method developed by Hardy and Senterre [65] and Baraloto et al. [66] to analyse respectively the phyloge-
200 netic (Π_{ST}) and phenotypic (τ_{ST}) structures of the hummingbird communities of Ecuador (clustering or overdispersion).
201 This method relies on computing indices inspired by the Simpson index and the fixation index F_{ST} , comparing the ob-
202 served diversity within and between the communities. For phylogeny, Π_{ST} can reveal phylogenetic clustering ($\Pi_{ST} > 0$)
203 or phylogenetic overdispersion ($\Pi_{ST} < 0$) within communities. Likewise, for phenotypic traits, τ_{ST} can reveal phenotypic
204 clustering ($\tau_{ST} > 0$) or phenotypic overdispersion ($\tau_{ST} < 0$) within communities. Statistical significance of overdispersion
205 or clustering is obtained from comparing the observed value to that obtained [for the same patch location](#) from 1000
206 random communities (created by drawing from the total species pool, using algorithm 1s from Hardy [67], which keeps
207 the local species richness per site constant). This approach compares the phenotypic structure to what would be expected
208 by chance.

209 To disentangle the relative effect of ecological (species sorting) and evolutionary mechanisms (selection), we also perform
210 our analyses by taking into account the phylogenetic relationships between species. If the species in the community are
211 more clustered or overdispersed than expected given their phylogenetic relationships, this is taken as evidence that the
212 trait has not evolved in a Brownian fashion (detailed in table 1). To this end, we used the `decouple` function [68], which
213 returns phylogenetically predicted and residual trait values by performing a linear regression of individual trait values
214 explained by the phylogeny. We computed the value of τ_{ST} on trait values decoupled from the phylogeny. This value is
215 hereafter denoted $dc\tau_{ST}$. Similarly to the classical τ_{ST} , the sign of $dc\tau_{ST}$ indicates phenotypic clustering ($dc\tau_{ST} > 0$) or
216 overdispersion ($dc\tau_{ST} < 0$) once the effect of the phylogenetic structure of the communities has been decoupled.

	$\tau_{ST} < 0$	$\tau_{ST} = 0$	$\tau_{ST} > 0$
$d\tau_{ST} < 0$	<p>Phenotypic overdispersion</p> <p>Co-occurring species are less similar than expected by chance because of character displacement.</p>	<p>No community structure</p> <p>Co-occurring species are nor more neither less similar than expected by chance despite character displacement because closely related species co-occur more often than expected at random (phylogenetic clustering; $\Pi_{ST} > 0$).</p>	<p>Phenotypic clustering</p> <p>Co-occurring species are more similar than expected by chance despite character displacement because closely related species co-occur more often than expected at random (phylogenetic clustering; $\Pi_{ST} > 0$).</p>
<p>Character displacement (divergence): co-occurring species are more dissimilar than expected given their phylogenetic relationships, which means they evolved towards dissimilarity in their colours.</p>			
$d\tau_{ST} = 0$	<p>Competitive exclusion: co-occurring species are more dissimilar than expected by chance because distantly-related (and therefore dissimilar) species co-occur more often than expected at random (phylogenetic overdispersion; $\Pi_{ST} < 0$).</p>	<p>Co-occurring species are not more similar nor more different than expected by change or than predicted given their phylogenetic relationships.</p>	<p>Environmental filtering: co-occurring species are more similar than expected by chance because closely-related (and therefore similar) species co-occur more often than expected at random (phylogenetic clustering; $\Pi_{ST} > 0$).</p>
$d\tau_{ST} > 0$	<p>Co-occurring species are less similar than expected by chance despite evolutionary convergence because distantly-related species co-occur more often than expected at random (phylogenetic overdispersion; $\Pi_{ST} < 0$).</p>	<p>Co-occurring species are neither more nor less similar than expected by chance despite evolutionary because distantly-related species co-occur more often than expected at random (phylogenetic overdispersion; $\Pi_{ST} < 0$).</p>	<p>Co-occurring species are more similar than expected by chance because of evolutionary convergence.</p>
<p>Evolutionary convergence : co-occurring species are more similar than expected given their phylogenetic relationships, which means they evolved towards similarity in their colours.</p>			

Table 1: Summary of the different evolutionary and ecological scenarios and their results in terms of values of τ_{ST} and decoupled $d\tau_{ST}$.

Analyses performed on a tree distribution (Π_{ST} and $d\tau_{ST}$) with n trees return a distribution of n statistics values and n p-values p_i . We summarised this information by computing the median of the statistics and the overall p-value p by using Jost's formula [69]:

$$p = k \sum_{i=0}^{n-1} \frac{(-\ln(k))^i}{i!} \quad \text{where } k = \prod_{i=1}^n p_i \quad (1)$$

Results

We find a strong phylogenetic clustering within communities ($\Pi_{ST} = 0.062 > 0$, $p < 0.0001$), indicating that co-occurring species are more closely related than expected by chance.

Phenotypic structure of the communities (predictions 1 - 4)

When looking at the bird entire body (when all patches are included simultaneously) by computing the overlap of the colour volumes, we did not find any phenotypic structure.

When the different major patches (crown, back, rump, tail, throat, breast, belly and wing) are examined separately (table 2 and table S2), we find clustering ($\tau_{ST} > 0$) in hue and hue shift on the back, rump, tail, belly and wing. Once we decouple the effect of the shared evolutionary history, we find clustering on the crown and the back ($d\tau_{ST} > 0$) but overdispersion on the belly for both hue and hue shift ($d\tau_{ST} < 0$). Hue shift is also overdispersed on the rump and the tail ($d\tau_{ST} < 0$). There is no phenotypic structure on the throat, breast or wing for hue and hue shift nor on the rump or the tail for hue.

We find no phenotypic structure (neither clustering nor overdispersion) for brightness on any patches before phylogenetic correction. After phylogenetic correction, brightness values for the throat, breast and belly are clustered among co-occurring species ($d\tau_{ST} > 0$) but show no phenotypic structure for the crown, the back, the wing and the tail.

Effect of community species richness on colour characteristics (prediction 5)

We found that the brightness range within a community increased in the same way as a null model built from random species assemblages (fig. 1b). For colour volume, we find some outliers with a higher colour volume than expected for community with the same number of species (fig. 1a).

Discussion

Our findings ~~suggest~~ are consistent with our hypothesis that colour structure within hummingbird communities likely results from the interplay between two selective pressures, acting in opposite directions: selection by the local environment (e.g. camouflage from predators, leading to phenotypic clustering on dorsal patches, and selection for species recognition, leading to phenotypic overdispersion on ventral and facial patches. We also discuss other possible effects that might have contributed to the observed pattern.

Variable	Phenotypic structure (τ_{ST})	Decoupled phenotypic structure ($dc\tau_{ST}$)
Hue		
Brightness		
Hue shift (=iridescence)		

Table 2: Phenotypic structure of hummingbird communities for different variables (hue, brightness and hue shift) on the patches studied (crown, back, rump, tail, throat, breast, belly, wing; names and locations illustrated in fig. S1). Hue is a tridimensional variable defined by the reflectance spectrum position x , y and z in the tetrahedron representing avian colour space. Blue plus signs $+$ indicate significant phenotypic clustering (τ_{ST} or $dc\tau_{ST} > 0$), orange minus signs $-$ indicate significant phenotypic overdispersion (τ_{ST} or $dc\tau_{ST} < 0$), and green zeros 0 represent the absence of phenotypic structure. The left column shows the raw phenotypic structure of the community (columns in table 1), which may be influenced by the phylogenetic structure while the right column shows the phenotypic structure of the community, decoupled from all effects caused by the phylogeny (rows in table 1). By comparing the values of τ_{ST} and $dc\tau_{ST}$ for each trait colour variable (hue, brightness and hue shift), we can assume a probable evolutionary scenario for each patches, based on the explanation in table 1. Exact values for the statistics are available in table S2.

245 Evidence for different evolutionary scenarios depending on patch location

246 At the entire bird level (i.e. when pooling together all patches), we did not find any phenotypic structure. But as mentioned
247 earlier, this was expected since different locations on the birds are ~~expected~~thought to be under different selection regimes
248 [7, 28].

249 In accordance with our prediction 5, community colour volume (as estimated by the convex hull of hue and brightness
250 range within a community) increases slightly faster with the number of species in the community than predicted by a null
251 model. This suggests that co-occurring species in these communities tend to use more similar colours than expected by
252 chance. However, this is not the ~~cause~~case for the majority of communities, where co-occurring species do not use more
253 nor less similar colours than expected by chance. This is further confirmed by the absence of phenotypic structure on the
254 colour volume and the brightness when the effect of the phylogeny is not decoupled.

255 This could be the consequence of similar selective pressures between the communities we studied, leading colours in
256 all assemblages to be randomly determined. This is however not very likely because the communities we studied differ a
257 lot in both their vegetation background and therefore in the pressure for crypsis [45] and in their species composition. A
258 more likely hypothesis is that co-occurring species tend to use the same colours but not necessarily on the same patches,
259 which would also explain the absence of phenotypic structure when we pool all patches without taking into account their
260 location. This is confirmed by our analysis patch by patch, where we find either clustering or overdispersion depending
261 on the location of the patch.

262 Selection for convergence and phenotypic clustering

263 In accordance with our first two predictions, co-occurring hummingbird species tend to have similar hues on patches more
264 likely dedicated to camouflage (back, rump, tail, wing; prediction 1) but not on patches more likely used in communication
265 (crown, throat, breast; prediction 2), as shown in table 2 and table S2. This new result for iridescence colours matches what
266 has been previously described for non-iridescent colours [7, 28]. The phenotypic clustering observed for hue on the rump,
267 the tail and the wing vanishes after decoupling the clustering effect due to phylogenetic structure. This ~~means~~suggests
268 that phenotypic clustering of hue on the rump, the tail and the wing is not caused by convergent evolution of co-occurring
269 species but by environmental filtering, leading related, similar-looking species to live in the same area (as explained in
270 table 1). This is confirmed by the high value of phylogenetic clustering. This sign of phylogenetic clustering completes
271 the results from Graham et al. [45] on the same dataset. We showed that intra-community species relatedness is high
272 compared to inter-community species relatedness (Π_{ST}), while they showed that intra-community species relatedness (Net
273 Relatedness Index) is higher than expected from random assemblages in 71 % of the cases [45]. This phylogenetic clustering
274 may be caused by a strong niche conservatism but our study cannot discriminate whether such niche conservatism involves
275 colour or other ecological traits. ~~However,~~Our data does not allow to assert with certainty the evolutionary history from
276 the pattern we observe but the predominance of green and brown hues on the back and the wing respectively, as shown
277 in fig. S4, hints to a role in camouflage. Alternatively, this phylogenetic clustering could be caused by hummingbirds'
278 costly hovering flight at high elevation due to weaker lift caused by the decreasing atmospheric pressure [70–72] ~~and~~, high

279 foraging specialisation [73] ~~likely contribute to this pattern. Alternatively, phylogenetic clustering could also be caused by~~
280 ~~a very or~~ low dispersal ability ~~of hummingbirds~~, but this last hypothesis remains quite unlikely as the rare studies on this
281 topic have shown that different hummingbird species display a wide variation in their dispersal ability [74, 75].

282 Contrary to our prediction 2, we also find clustering of hue on the belly before the use of the `decouple` function.
283 However, the fact that it turns into overdispersion after the use of the `decouple` function, and not simply into a random
284 phenotypic structure (as opposed to the rump, the tail and the wing mentioned just before), suggests this initial clustering
285 (right column in table 1) is mainly caused by environmental filtering on another trait but that hue on the belly is still
286 under selection for divergence (first row in table 1). This other trait may be the colour of another patch or other ecological
287 traits, as we explained previously.

288 We found a significant clustering of brightness on the throat, breast and belly after controlling for the phylogeny,
289 indicating that brightness on those patches is more similar than expected given the phylogeny among co-occurring species
290 (prediction 3bis). This suggests that the same patches have been selected to be involved either in communication or
291 in camouflage among species living in the same environment. This is seen after controlling for the phylogeny and it
292 is therefore not caused by the phylogenetic relatedness of co-occurring species. This is not surprising as many studies
293 showed the paramount importance of the throat in the courtship display of many hummingbird species [29–32, 76] Two
294 main hypotheses can explain why co-occurring species tend to communicate (or camouflage themselves) using the same
295 patches: (i) There may be selective pressures for the use of specific patches in camouflage in a given environment (e. g.,
296 patches that are more exposed to predators' sight). (ii) Convergence in patches used in communication may be selected
297 because it improves competitor identification in the case of a strong ecological niche overlap (convergence by agonistic
298 character displacement as shown in Grether et al. [13] and Tobias et al. [77]).

299 All those results suggest a strong effect of the environment in the evolution of colour in agreement with McNaught
300 and Owens [78] who found that bird plumage colour was due to the light environment and not to reproductive character
301 displacement in Australian birds. However, we do not find clustering on all patches, which ~~means that~~ suggests that, for
302 some patches, the effect of habitat pressure is somehow limited or counterbalanced by reproductive or agonistic character
303 displacement. On the contrary, for some patches, we found patterns that are likely the result of character displacement.

304 **Character displacement and phenotypic overdispersion**

305 In agreement with our prediction 2, after decoupling the effect of the phylogeny, there is overdispersion of hue on the
306 belly, likely caused by character displacement (table 1). At a completely different taxonomic scale, focusing on a single
307 hummingbird genus (*Coeligena*) with 11 species, Parra [33] also found that the belly was always involved in the difference
308 in hue between subspecies. It was sometimes even the only patch causing those differences, as for example between
309 *Coeligena torquata fulgidigula* and *Coeligena torquata torquata*. This suggests that the interspecific divergence we found
310 on the belly at the community level on the whole Trochilidae family can be observed at different geographic and taxonomic
311 scales, and even between subspecies of the same species.

312 As predicted, we also find more phenotypic overdispersion for hue shift than hue after decoupling the effect of the

313 phylogeny, for example, on the rump and on the tail (prediction 4). It is possible that hue shift is less sensitive to selection
314 for convergence because it may vary without disturbing camouflage efficacy. However, we did not find the expected
315 relaxing of clustering on hue shift on patches such as the back. This is likely caused by the fact that hue shift is highly
316 correlated with hue, as found in this study and in Dakin and Montgomerie [35], who used the same indices to quantify
317 iridescence. This correlation is due to the optics controlling iridescence, meaning that species that display similar hues
318 should also display the same hue shift if they use the same underlying multilayer structures. The fact that the correlation
319 is not perfect and that we nonetheless get different phenotypic patterns for hue and hue shift on some patches suggests
320 that co-occurring species use different multilayer structures (as recently confirmed by [79]), which can produce different
321 iridescent effects while displaying the same hue (functional convergence on hue).

322 Against our prediction 2, we did not find phenotypic overdispersion on any of the colour variables on patches such as
323 the throat or the crown, that are thought to be sexually selected and often used in courtship displays [29, 80]. Several
324 hypotheses can explain this fact: (i) The overdispersion on some patches (hue on the belly and hue shift on the rump and
325 tail) is sufficient to enable species recognition. (ii) The current phenotypic structure, which is neither overdispersed nor
326 clustered, on those patches is sufficient to enable species recognition. Indeed, the absence of phenotypic overdispersion
327 does not mean that species look the same. It simply means that colour differences between species living in the same
328 community and species in different communities occur in similar ranges. This difference may be sufficient to relax the
329 selective pressure towards reproductive character displacement. (iii) The pressure towards overdispersion is balanced by
330 habitat filtering (for both ventral and dorsal patches), resulting in no apparent phenotypic structure. The latter hypothesis
331 was also a candidate explanation of the pattern found by Martin et al. [21], where sympatric closely related species are
332 more divergent than allopatric ones, but only when the range overlap is limited. They suggested that local adaptation
333 could hinder divergence when species ranges was exactly the same. (iv) Species recognition is achieved by additional means
334 and divergence occurs on others traits, such as modified feathers [81], song [82, 83] or non-vocal noises [84–86] and size.
335 Notably, different species of hummingbirds can have very different courtship behaviour: leks for hermits [87, 88], dives
336 and shuttle displays for bees [31, 85, 89], for instance.

337 Taken together, our results suggest that hummingbird iridescent colours are determined by different evolutionary
338 mechanisms depending on their location. Within a community, co-occurring hummingbird species tend to display the
339 same hues on dorsal patches ~~probably because of~~ which is what we expect if colour on these patches is mainly driven
340 by selective pressures related to the local environment, such as selection for crypsis by predators, causing phenotypic
341 clustering at the community level. This phenotypic clustering does not seem to be caused by adaptive convergence on
342 colours but rather by environmental filtering perhaps linked to other ecological traits such as elevation tolerance or flight
343 ability. In spite ~~of such~~ this suspected environmental filtering, ~~character displacement leads to~~ there is overdispersion
344 for hue on the belly and hue shift on the rump and the tail. ~~Iridescence may therefore~~ This suggest a possible role of
345 character displacement, which could mean that iridescence could be used a way to enable species recognition without
346 affecting camouflage efficacy of birds, by opening up a new dimension in the sensory space: hue shift.

347 Acknowledgments

348 This project heavily relied on museum specimens which were made available by the work of collection curators: Patrick
349 Boussès, Anne Previato, and Jérôme Fuchs (Muséum National d’Histoire Naturelle), Cédric Audibert and Harold Labrique
350 (Musée des Confluences).

351 Conflict of interest disclosure

352 The authors of this preprint declare that they have no financial conflict of interest with the content of this article. Marianne
353 Elias is part of the managing board of PCIEvolBiol and is one of the PCIEvolBiol recommenders.

354 References

- 355 [1] JW Bradbury and SL Vehrencamp. *Principles of Animal Communication*. 2. ed. OCLC: 759797180. Sunderland,
356 Mass: Sinauer Associates, 2011. 697 pp. ISBN: 978-0-87893-045-6.
- 357 [2] AR Parker. “515 Million Years of Structural Colour”. In: *Journal of Optics A: Pure and Applied Optics* 2.6 (2000),
358 R15–R28. DOI: 10.1088/1464-4258/2/6/201.
- 359 [3] P Vukusic. “Natural Photonics”. In: *Physics World* 17.2 (2004), p. 35. DOI: 10.1088/2058-7058/17/2/34.
- 360 [4] SM Doucet and MG Meadows. “Iridescence: A Functional Perspective”. In: *Journal of The Royal Society Interface*
361 6 (Suppl 2 2009), S115–S132. DOI: 10.1098/rsif.2008.0395.focus. pmid: 19336344.
- 362 [5] RO Prum. “Anatomy, Physics, and Evolution of Structural Colors”. In: *Bird Coloration, Volume 1: Mechanisms and*
363 *Measurements*. Ed. by GE Hill and KJ McGraw. Vol. 1. 2 vols. Bird Coloration. Harvard University Press, 2006,
364 p. 640. ISBN: 978-0-674-01893-8.
- 365 [6] DC Osorio and AD Ham. “Spectral Reflectance and Directional Properties of Structural Coloration in Bird Plumage”.
366 In: *Journal of Experimental Biology* 205.14 (2002), pp. 2017–2027. pmid: 12089207.
- 367 [7] D Gomez and M Théry. “Simultaneous Crypsis and Conspicuousness in Color Patterns: Comparative Analysis of
368 a Neotropical Rainforest Bird Community”. In: *The American Naturalist* 169.s1 (2007), S42–S61. DOI: 10.1086/
369 510138.
- 370 [8] JA Endler. “The Color of Light in Forests and Its Implications”. In: *Ecological Monographs* 63.1 (1993), pp. 1–27.
371 DOI: 10.2307/2937121.
- 372 [9] D Gomez and M Théry. “Influence of Ambient Light on the Evolution of Colour Signals: Comparative Analysis of
373 a Neotropical Rainforest Bird Community”. In: *Ecology Letters* 7.4 (2004), pp. 279–284. DOI: 10.1111/j.1461-
374 0248.2004.00584.x.
- 375 [10] J Gröning and A Hochkirch. “Reproductive Interference Between Animal Species”. In: *The Quarterly Review of*
376 *Biology* 83.3 (2008), pp. 257–282. DOI: 10.1086/590510.

- 377 [11] WL Brown and EO Wilson. “Character Displacement”. In: *Systematic Biology* 5.2 (1956), pp. 49–64. DOI: 10.2307/
378 2411924.
- 379 [12] R Butlin. “Speciation by Reinforcement”. In: *Trends in Ecology & Evolution* 2.1 (1987), pp. 8–13. DOI: 10.1016/
380 0169-5347(87)90193-5.
- 381 [13] GF Grether, N Losin, CN Anderson, and K Okamoto. “The Role of Interspecific Interference Competition in Char-
382 acter Displacement and the Evolution of Competitor Recognition”. In: *Biological Reviews* 84.4 (2009), pp. 617–635.
383 DOI: 10.1111/j.1469-185X.2009.00089.x. pmid: 19681844.
- 384 [14] JA Tobias and N Seddon. “Signal Design and Perception in *Hypocnemis* Antbirds: Evidence for Convergent Evolution
385 via Social Selection”. In: *Evolution* 63.12 (2009), pp. 3168–3189. DOI: 10.1111/j.1558-5646.2009.00795.x. pmid:
386 19659594.
- 387 [15] HC Gerhardt. “Reproductive Character Displacement of Female Mate Choice in the Grey Treefrog, *Hyla Chrysoscelis*”.
388 In: *Animal Behaviour* 47.4 (1994), pp. 959–969. DOI: 10.1006/anbe.1994.1127.
- 389 [16] BR Grant and PR Grant. “Songs of Darwin’s Finches Diverge When a New Species Enters the Community”. In:
390 *Proceedings of the National Academy of Sciences* 107.47 (2010), pp. 20156–20163. DOI: 10.1073/pnas.1015115107.
391 pmid: 21048082.
- 392 [17] PMB Bacquet, O Brattström, HL Wang, CE Allen, C Löfstedt, et al. “Selection on Male Sex Pheromone Composition
393 Contributes to Butterfly Reproductive Isolation”. In: *Proceedings of the Royal Society B: Biological Sciences* 282.1804
394 (2015), p. 20142734. DOI: 10.1098/rspb.2014.2734. pmid: 25740889.
- 395 [18] GP Sætre, T Moum, S Bureš, M Král, M Adamjan, et al. “A Sexually Selected Character Displacement in Flycatchers
396 Reinforces Premating Isolation”. In: *Nature* 387.6633 (1997), pp. 589–592. DOI: 10.1038/42451. pmid: 847.
- 397 [19] RE Naisbit, CD Jiggins, and J Mallet. “Disruptive Sexual Selection against Hybrids Contributes to Speciation
398 between *Heliconius Cydno* and *Heliconius Melpomene*”. In: *Proceedings of the Royal Society of London B: Biological*
399 *Sciences* 268.1478 (2001), pp. 1849–1854. DOI: 10.1098/rspb.2001.1753. pmid: 11522205.
- 400 [20] VA Lukhtanov, NP Kandul, JB Plotkin, AV Dantchenko, D Haig, et al. “Reinforcement of Pre-Zygotic Isolation
401 and Karyotype Evolution in *Agrodiaetus* Butterflies”. In: *Nature* 436.7049 (2005), pp. 385–389. DOI: 10.1038/
402 nature03704.
- 403 [21] PR Martin, R Montgomerie, and SC Loughheed. “Color Patterns of Closely Related Bird Species Are More Divergent
404 at Intermediate Levels of Breeding-Range Sympatry”. In: *The American Naturalist* 185.4 (2015), pp. 443–451. DOI:
405 10.1086/680206.
- 406 [22] C Doutrelant, M Paquet, JP Renoult, A Grégoire, PA Crochet, et al. “Worldwide Patterns of Bird Colouration on
407 Islands”. In: *Ecology Letters* 19.5 (2016), pp. 537–545. DOI: 10.1111/ele.12588.
- 408 [23] CR Hemingson, PF Cowman, JR Hodge, and DR Bellwood. “Colour Pattern Divergence in Reef Fish Species Is
409 Rapid and Driven by Both Range Overlap and Symmetry”. In: *Ecology Letters* 22.1 (2019), pp. 190–199. DOI:
410 10.1111/ele.13180.


- 411 [24] ATD Bennett, IC Cuthill, and KJ Norris. “Sexual Selection and the Mismeasure of Color”. In: *The American*
412 *Naturalist* 144.5 (1994), pp. 848–860. DOI: 10.1086/285711.
- 413 [25] IC Cuthill, ATD Bennett, JC Partridge, and EJ Maier. “Plumage Reflectance and the Objective Assessment of Avian
414 Sexual Dichromatism”. In: *The American Naturalist* 153.2 (1999), pp. 183–200. DOI: 10.1086/303160.
- 415 [26] MD Eaton. “Human Vision Fails to Distinguish Widespread Sexual Dichromatism among Sexually “Monochromatic”
416 Birds”. In: *Proceedings of the National Academy of Sciences* 102.31 (2005), pp. 10942–10946. DOI: 10.1073/pnas.
417 0501891102. pmid: 16033870.
- 418 [27] R Montgomerie. “Analyzing Colors”. In: *Bird Coloration, Volume 1: Mechanisms and Measurements*. Ed. by GE Hill
419 and KJ McGraw. Vol. 1. 2 vols. Bird Coloration. Harvard University Press, 2006, p. 640. ISBN: 978-0-674-01893-8.
- 420 [28] K Delhey. “Revealing The Colourful Side of Birds: Spatial Distribution of Conspicuous Plumage Colours on The
421 Body of Australian Birds”. In: *bioRxiv* (2019), p. 647727. DOI: 10.1101/647727.
- 422 [29] FG Stiles. “Aggressive and Courtship Displays of the Male Anna’s Hummingbird”. In: *The Condor* 84.2 (1982),
423 pp. 208–225. DOI: 10.2307/1367674. JSTOR: 1367674.
- 424 [30] RK Simpson and KJ McGraw. “Experimental Trait Mis-Matches Uncover Specificity of Evolutionary Links between
425 Multiple Signaling Traits and Their Interactions in Hummingbirds”. In: *Evolution* 73.3 (2019), pp. 436–451. DOI:
426 10.1111/evo.13662.
- 427 [31] RK Simpson and KJ McGraw. “Two Ways to Display: Male Hummingbirds Show Different Color-Display Tactics
428 Based on Sun Orientation”. In: *Behavioral Ecology* 29.3 (2018), pp. 637–648. DOI: 10.1093/beheco/ary016.
- 429 [32] BG Hogan and MC Stoddard. “Synchronization of Speed, Sound and Iridescent Color in a Hummingbird Aerial
430 Courtship Dive”. In: *Nature Communications* 9.1 (2018), p. 5260. DOI: 10.1038/s41467-018-07562-7.
- 431 [33] JL Parra. “Color Evolution in the Hummingbird Genus *Coeligena*”. In: *Evolution* 64.2 (2010), pp. 324–335. DOI:
432 10.1111/j.1558-5646.2009.00827.x. pmid: 19703221.
- 433 [34] BD Wilts, K Michielsen, J Kuipers, H De Raedt, and DG Stavenga. “Brilliant Camouflage: Photonic Crystals in the
434 Diamond Weevil, *Entimus Imperialis*”. In: *Proceedings of the Royal Society B: Biological Sciences* 279.1738 (2012),
435 pp. 2524–2530. DOI: 10.1098/rspb.2011.2651.
- 436 [35] R Dakin and R Montgomerie. “Eye for an Eyespot: How Iridescent Plumage Ocelli Influence Peacock Mating Success”.
437 In: *Behavioral Ecology* 24.5 (2013), pp. 1048–1057. DOI: 10.1093/beheco/art045.
- 438 [36] S Villéger, NWH Mason, and D Mouillot. “New Multidimensional Functional Diversity Indices for a Multifaceted
439 Framework in Functional Ecology”. In: *Ecology* 89.8 (2008), pp. 2290–2301. DOI: 10.1890/07-1206.1.
- 440 [37] J Del Hoyo, A Elliott, J Sargatal, DA Christie, and E de Juana. *Handbook of the Birds of the World Alive*. 2017.
441 URL: hbw.com.
- 442 [38] J Dorst. “Recherches sur la structure des plumes des trochilidés”. OCLC: 14220401. Paris: Université de Paris, 1951.
443 260 pp.

- 444 [39] H Dürer. “Schillerfarben der Vogelfeder als Evolutionsproblem”. Medizinischen Fakultät der Universität Basel, 1977.
- 445 [40] W Jetz, GH Thomas, JB Joy, K Hartmann, and AO Mooers. “The Global Diversity of Birds in Space and Time”.
446 In: *Nature* 491.7424 (2012), pp. 444–448. DOI: 10.1038/nature11631. pmid: 23123857.
- 447 [41] JA McGuire, CC Witt, JVJ Remsen, A Corl, DL Rabosky, et al. “Molecular Phylogenetics and the Diversification of
448 Hummingbirds”. In: *Current Biology* 24.8 (2014), pp. 910–916. DOI: 10.1016/j.cub.2014.03.016. pmid: 24704078.
- 449 [42] GR Graves and RL Zusi. “An Intergeneric Hybrid Hummingbird (*Heliodoxa leadbeateri* *Heliangelus amethysticollis*)
450 from Northern Colombia”. In: *The Condor* 92.3 (1990), pp. 754–760. DOI: 10.2307/1368695.
- 451 [43] FG Stiles and JO Cortés-Herrera. “Diagnosis and Observations of a Hybrid Hummingbird (*Metallura tyrianthina* x
452 *Aglaiocercus kingi*) in the Eastern Andes of Colombia”. In: *Revista de la Academia Colombiana de Ciencias Exactas,*
453 *Físicas y Naturales* 39.153 (2015), pp. 481–490. DOI: 10.18257/raccefyn.260.
- 454 [44] SM Doucet and GE Hill. “Do Museum Specimens Accurately Represent Wild Birds? A Case Study of Carotenoid,
455 Melanin, and Structural Colours in Long-Tailed Manakins *Chiroxiphia linearis*”. In: *Journal of Avian Biology* 40.2
456 (2009), pp. 146–156. DOI: 10.1111/j.1600-048X.2009.03763.x.
- 457 [45] CH Graham, JL Parra, C Rahbek, and JA McGuire. “Phylogenetic Structure in Tropical Hummingbird Com-
458 munities”. In: *Proceedings of the National Academy of Sciences* 106 (Supplement 2 2009), pp. 19673–19678. DOI:
459 10.1073/pnas.0901649106. pmid: 19805042.
- 460 [46] JL Parra, JA McGuire, and CH Graham. “Incorporating Clade Identity in Analyses of Phylogenetic Community
461 Structure: An Example with Hummingbirds.” In: *The American Naturalist* 176.5 (2010), pp. 573–587. DOI: 10.
462 1086/656619. pmid: 20849270.
- 463 [47] RS Ridgely and PJ Greenfield. *The Birds of Ecuador: Status, Distribution and Taxonomy*. Ithaca, NY: Cornell
464 University Press, 2001. 880 pp. ISBN: 978-0-8014-8720-0.
- 465 [48] RL Dalrymple, FKC Hui, H Flores-Moreno, DJ Kemp, and AT Moles. “Roses Are Red, Violets Are Blue – so How
466 Much Replication Should You Do? An Assessment of Variation in the Colour of Flowers and Birds”. In: *Biological*
467 *Journal of the Linnean Society* 114.1 (2015), pp. 69–81. DOI: 10.1111/bij.12402.
- 468 [49] MG Meadows, NI Morehouse, RL Rutowski, JM Douglas, and KJ McGraw. “Quantifying Iridescent Coloration in
469 Animals: A Method for Improving Repeatability”. In: *Behavioral Ecology and Sociobiology* 65.6 (2011), pp. 1317–
470 1327. DOI: 10.1007/s00265-010-1135-5. pmid: 876.
- 471 [50] DM Chen and TH Goldsmith. “Four Spectral Classes of Cone in the Retinas of Birds”. In: *Journal of Comparative*
472 *Physiology A* 159.4 (1986), pp. 473–479. DOI: 10.1007/BF00604167.
- 473 [51] TE White, RL Dalrymple, DW Noble, JC O’Hanlon, DB Zurek, et al. “Reproducible Research in the Study of
474 Biological Coloration”. In: *Animal Behaviour* 106 (2015), pp. 51–57. DOI: 10.1016/j.anbehav.2015.05.007.
- 475 [52] MG Meadows, TE Roudybush, and KJ McGraw. “Dietary Protein Level Affects Iridescent Coloration in Anna’s
476 Hummingbirds, *Calypte anna*”. In: *Journal of Experimental Biology* 215.16 (2012), pp. 2742–2750. DOI: 10.1242/
477 jeb.069351. pmid: 22837446.

- 478 [53] H Gruson, C Andraud, W Daney de Marcillac, S Berthier, M Elias, et al. “Quantitative Characterization of Iridescent
479 Colours in Biological Studies: A Novel Method Using Optical Theory”. In: *Interface Focus* 9.1 (2019), p. 20180049.
480 DOI: 10.1098/rsfs.2018.0049.
- 481 [54] R Maia, H Gruson, JA Endler, and TE White. “Pavo 2: New Tools for the Spectral and Spatial Analysis of Colour
482 in R”. In: *Methods in Ecology and Evolution* 10.7 (2019), pp. 1097–1107. DOI: 10.1111/2041-210X.13174.
- 483 [55] JA Endler and PW Mielke. “Comparing Entire Colour Patterns as Birds See Them”. In: *Biological Journal of the
484 Linnean Society* 86.4 (2005), pp. 405–431. DOI: 10.1111/j.1095-8312.2005.00540.x.
- 485 [56] DC Osorio and M Vorobyev. “A Review of the Evolution of Animal Colour Vision and Visual Communication
486 Signals”. In: *Vision Research. Vision Research Reviews* 48.20 (2008), pp. 2042–2051. DOI: 10.1016/j.visres.2008.
487 06.018. pmid: 18627773.
- 488 [57] A Ödeen and O Håstad. “Complex Distribution of Avian Color Vision Systems Revealed by Sequencing the SWS1
489 Opsin from Total DNA”. In: *Molecular Biology and Evolution* 20.6 (2003), pp. 855–861. DOI: 10.1093/molbev/
490 msg108. pmid: 12716987.
- 491 [58] G Herrera, JC Zagal, M Diaz, MJ Fernández, A Vielma, et al. “Spectral Sensitivities of Photoreceptors and Their Role
492 in Colour Discrimination in the Green-Backed Firecrown Hummingbird (*Sephanoides Sephaniodes*)”. In: *Journal of
493 Comparative Physiology A* 194.9 (2008), p. 785. DOI: 10.1007/s00359-008-0349-8. pmid: 18584181.
- 494 [59] A Ödeen and O Håstad. “Pollinating Birds Differ in Spectral Sensitivity”. In: *Journal of Comparative Physiology A*
495 196.2 (2010), pp. 91–96. DOI: 10.1007/s00359-009-0474-z.
- 496 [60] NS Hart. “Microspectrophotometry of Visual Pigments and Oil Droplets in a Marine Bird, the Wedge-Tailed Shear-
497 water *Puffinus Pacificus*: Topographic Variations in Photoreceptor Spectral Characteristics”. In: *Journal of Exper-
498 imental Biology* 207.7 (2004), pp. 1229–1240. DOI: 10.1242/jeb.00857. pmid: 14978063.
- 499 [61] DF Stotz, JW Fitzpatrick, TA Parker III, and DK Moskovits. *Neotropical Birds: Ecology and Conservation*. Vol. 3.
500 OCLC: 32819832. University of Chicago Press, 1996. ISBN: 978-0-226-77629-3.
- 501 [62] MA Stoffel, S Nakagawa, and H Schielzeth. “rptR: Repeatability Estimation and Variance Decomposition by Gener-
502 alized Linear Mixed-Effects Models”. In: *Methods in Ecology and Evolution* 8.11 (2017). Ed. by S Goslee, pp. 1639–
503 1644. DOI: 10.1111/2041-210X.12797.
- 504 [63] MC Stoddard and RO Prum. “Evolution of Avian Plumage Color in a Tetrahedral Color Space: A Phylogenetic
505 Analysis of New World Buntings.” In: *The American Naturalist* 171.6 (2008), pp. 755–776. DOI: 10.1086/587526.
- 506 [64] M Pagel and F Lutzoni. “Accounting for Phylogenetic Uncertainty in Comparative Studies of Evolution and Adap-
507 tation”. In: *Biological Evolution and Statistical Physics*. Ed. by M Lässig and A Valleriani. Vol. 585. Lecture Notes
508 in Physics. Berlin, Heidelberg: Springer Berlin Heidelberg, 2002. ISBN: 978-3-540-43188-6. DOI: 10.1007/3-540-
509 45692-9.
- 510 [65] OJ Hardy and B Senterre. “Characterizing the Phylogenetic Structure of Communities by an Additive Partitioning of
511 Phylogenetic Diversity”. In: *Journal of Ecology* 95.3 (2007), pp. 493–506. DOI: 10.1111/j.1365-2745.2007.01222.x.

- 512 [66] C Baraloto, OJ Hardy, CET Paine, KG Dexter, C Cruaud, et al. “Using Functional Traits and Phylogenetic Trees
513 to Examine the Assembly of Tropical Tree Communities”. In: *Journal of Ecology* 100.3 (2012), pp. 690–701. DOI:
514 10.1111/j.1365-2745.2012.01966.x.
- 515 [67] OJ Hardy. “Testing the Spatial Phylogenetic Structure of Local Communities: Statistical Performances of Different
516 Null Models and Test Statistics on a Locally Neutral Community”. In: *Journal of Ecology* 96.5 (2008), pp. 914–926.
517 DOI: 10.1111/j.1365-2745.2008.01421.x.
- 518 [68] F de Bello, P Šmilauer, JAF Diniz-Filho, CP Carmona, Z Lososová, et al. “Decoupling Phylogenetic and Functional
519 Diversity to Reveal Hidden Signals in Community Assembly”. In: *Methods in Ecology and Evolution* 8.10 (2017),
520 pp. 1200–1211. DOI: 10.1111/2041-210X.12735.
- 521 [69] VN Balasubramanian, S Chakraborty, and S Panchanathan. “Conformal Predictions for Information Fusion”. In:
522 *Annals of Mathematics and Artificial Intelligence* 74.1-2 (2015), pp. 45–65. DOI: 10.1007/s10472-013-9392-4.
- 523 [70] RK Suarez. “Hummingbird Flight: Sustaining the Highest Mass-Specific Metabolic Rates among Vertebrates”. In:
524 *Experientia* 48.6 (1992), pp. 565–570. DOI: 10.1007/BF01920240.
- 525 [71] DL Altshuler, R Dudley, and JA McGuire. “Resolution of a Paradox: Hummingbird Flight at High Elevation Does
526 Not Come without a Cost”. In: *Proceedings of the National Academy of Sciences* 101.51 (2004), pp. 17731–17736.
527 DOI: 10.1073/pnas.0405260101. pmid: 15598748.
- 528 [72] DL Altshuler, FG Stiles, and R Dudley. “Of Hummingbirds and Helicopters: Hovering Costs, Competitive Ability,
529 and Foraging Strategies.” In: *The American Naturalist* 163.1 (2004), pp. 16–25. DOI: 10.1086/380511. pmid:
530 14767833.
- 531 [73] AB Lindberg and JM Olesen. “The Fragility of Extreme Specialization: *Passiflora Mixta* and Its Pollinating Hum-
532 mingbird *Ensifera Ensifera*”. In: *Journal of Tropical Ecology* 17.2 (2001), pp. 323–329. DOI: 10.1017/S0266467401001213.
- 533 [74] RP Moore, WD Robinson, IJ Lovette, and TR Robinson. “Experimental Evidence for Extreme Dispersal Limitation
534 in Tropical Forest Birds”. In: *Ecology Letters* 11.9 (2008), pp. 960–968. DOI: 10.1111/j.1461-0248.2008.01196.x.
- 535 [75] LN Céspedes, LI Pavan, JA Hazlehurst, and JE Jankowski. “The Behavior and Diet of the Shining Sunbeam (*Aglae-
536 actis Cupripennis*): A Territorial High-Elevation Hummingbird”. In: *The Wilson Journal of Ornithology* 131.1 (2019),
537 pp. 24–34. DOI: 10.1676/18-79.
- 538 [76] RK Simpson and KJ McGraw. “It’s Not Just What You Have, but How You Use It: Solar-Positional and Behavioural
539 Effects on Hummingbird Colour Appearance during Courtship”. In: *Ecology Letters* 21.9 (2018), pp. 1413–1422. DOI:
540 10.1111/ele.13125.
- 541 [77] JA Tobias, R Planqué, DL Cram, and N Seddon. “Species Interactions and the Structure of Complex Communication
542 Networks”. In: *Proceedings of the National Academy of Sciences* 111.3 (2014), pp. 1020–1025. DOI: 10.1073/pnas.
543 1314337111. pmid: 24395769.

- 544 [78] MK McNaught and IPF Owens. “Interspecific Variation in Plumage Colour among Birds: Species Recognition
545 or Light Environment?” In: *Journal of Evolutionary Biology* 15.4 (2002), pp. 505–514. DOI: 10.1046/j.1420-
546 9101.2002.00431.x.
- 547 [79] H Gruson, M Elias, C Andraud, C Djediat, S Berthier, et al. “Hummingbird Iridescence: An Unsuspected Structural
548 Diversity Influences Colouration at Multiple Scales”. In: *bioRxiv* (2019), p. 699744. DOI: 10.1101/699744.
- 549 [80] CJ Clark, TJ Feo, and I Escalante. “Courtship Displays and Natural History of Scintillant (*Selasphorus Scintilla*)
550 and Volcano (*S. Flammula*) Hummingbirds”. In: *The Wilson Journal of Ornithology* 123.2 (2011), pp. 218–228. DOI:
551 10.1676/10-076.1.
- 552 [81] CM Eliason, MD Shawkey, and JA Clarke. “Evolutionary Shifts in the Melanin-Based Color System of Birds”. In:
553 *Evolution* 70.2 (2016), pp. 445–454. DOI: 10.1111/evo.12855. pmid: 26044706.
- 554 [82] P Matyjasiak. “Birds Associate Species-Specific Acoustic and Visual Cues: Recognition of Heterospecific Rivals by
555 Male Blackcaps”. In: *Behavioral Ecology* 16.2 (2005), pp. 467–471. DOI: 10.1093/beheco/ari012.
- 556 [83] D Luther. “The Influence of the Acoustic Community on Songs of Birds in a Neotropical Rain Forest”. In: *Behavioral*
557 *Ecology* 20.4 (2009), pp. 864–871. DOI: 10.1093/beheco/arp074.
- 558 [84] CJ Clark and TJ Feo. “The Anna’s Hummingbird Chirps with Its Tail: A New Mechanism of Sonation in Birds”.
559 In: *Proceedings of the Royal Society of London B: Biological Sciences* 275.1637 (2008), pp. 955–962. DOI: 10.1098/
560 rspb.2007.1619. pmid: 18230592.
- 561 [85] CJ Clark, DO Elias, and RO Prum. “Aeroelastic Flutter Produces Hummingbird Feather Songs”. In: *Science* 333.6048
562 (2011), pp. 1430–1433. DOI: 10.1126/science.1205222. pmid: 21903810.
- 563 [86] CJ Clark. “Wing, Tail, and Vocal Contributions to the Complex Acoustic Signals of Courting Calliope Humming-
564 birds”. In: *Current Zoology* 57.2 (2011), pp. 187–196. DOI: 10.1093/czoolo/57.2.187.
- 565 [87] FG Stiles and LL Wolf. “Ecology and Evolution of Lek Mating Behavior in the Long-Tailed Hermit Hummingbird”.
566 In: *Ornithological Monographs* 27 (1979), pp. iii–78. DOI: 10.2307/40166760. JSTOR: 40166760.
- 567 [88] MA Pizo. “Lek Behavior of the Plovercrest (*Stephanoxis Lalandi*, Trochilidae)”. In: *The Wilson Journal of Ornithol-*
568 *ogy* 124.1 (2012), pp. 106–112. DOI: 10.1676/11-055.1.
- 569 [89] TA Hurly, RD Scott, and SD Healy. “The Function of Displays of Male Rufous Hummingbirds”. In: *The Condor*
570 103.3 (2001), pp. 647–651. DOI: 10.1650/0010-5422(2001)103[0647:TFODOM]2.0.CO;2.
- 571 [90] S Nakagawa and H Schielzeth. “Repeatability for Gaussian and Non-Gaussian Data: A Practical Guide for Biologists”.
572 In: *Biological Reviews* 85.4 (2010), pp. 935–956. DOI: 10.1111/j.1469-185X.2010.00141.x. pmid: 20569253.



Supplementary figure 1: Locations and names of the 8 patches measured on all species. Additional patches were measured for each species as soon as they differed from one of the 8 patches listed here for a human observer, as detailed in the methods section and as in Gomez and Théry [7].

Table 3: List of species with their provenance (Confluences = Musée des Confluences, Lyon, France, MNHN = Muséum National d’Histoire Naturelle, Paris, France), strata, and place of collection (when known). Strata data were extracted from Stotz et al. [61] and used in vision models.

Species	Clade	Provenance	Strata	Location
<i>Adelomyia melanogenys</i>	Coquette	Confluences	Understory	
<i>Aglaeactis cupripennis</i>	Brilliant	MNHN	Canopy	Ecuador
<i>Aglaiocercus coelestis</i>	Coquette	MNHN	Canopy	Ecuador
<i>Aglaiocercus kingi mocoa</i>	Coquette	MNHN	Canopy	Ecuador
<i>Amazilia amabilis</i>	Emerald	MNHN	Understory	Ecuador
<i>Amazilia amazilia</i>	Emerald	MNHN	Understory	Ecuador
<i>Amazilia fimbriata fluviatilis</i>	Emerald	MNHN	Canopy	Ecuador
<i>Amazilia franciae</i>	Emerald	MNHN	Canopy	Ecuador
<i>Amazilia grayi meridionalis</i>	Emerald	MNHN	Canopy	Ecuador
<i>Amazilia rosenbergi</i>	Emerald	MNHN	Understory	Ecuador
<i>Amazilia sapphirina</i>	Emerald	MNHN	Canopy	Brasil
<i>Amazilia tzacatl jucunda</i>	Emerald	MNHN	Canopy	Ecuador
<i>Androdon aequatorialis</i>	Mangoe	MNHN	Understory	Ecuador
<i>Anthracothorax nigricollis</i>	Mangoe	MNHN	Canopy	Colombia
<i>Avocettula recurvirostris</i>	Mangoe	Confluences	Understory	
<i>Boissonneaua flavescens</i>	Brilliant	MNHN	Canopy	Ecuador
<i>Boissonneaua matthewsii</i>	Brilliant	MNHN	Canopy	Ecuador
<i>Calliphlox amethystina</i>	Bee	MNHN	Canopy	Ecuador
<i>Calliphlox mitchellii</i>	Bee	Confluences	Canopy	
<i>Campylopterus falcatus</i>	Emerald	MNHN	Understory	Colombia
<i>Campylopterus largipennis</i>	Emerald	MNHN	Understory	Peru
<i>Campylopterus villaviscensio</i>	Emerald	MNHN	Understory	Ecuador
<i>Chaetocercus bombus</i>	Bee	MNHN	Canopy	Ecuador
<i>Chaetocercus mulsant</i>	Bee	MNHN	Understory	Ecuador
<i>Chalcostigma herrani</i>	Coquette	MNHN	Canopy	Ecuador
<i>Chalcostigma ruficeps</i>	Coquette	Confluences	Understory	
<i>Chalcostigma stanleyi stanleyi</i>	Coquette	MNHN	Canopy	Ecuador
<i>Chalybura buffonii intermedia</i>	Emerald	Confluences	Understory	
<i>Chalybura urochrysia urochrysia</i>	Emerald	Confluences	Understory	
<i>Chlorestes notata obsoletus-puruensis</i>	Emerald	Confluences	Canopy	
<i>Chlorostilbon melanorhynchus</i>	Emerald	MNHN	Understory	Ecuador

Species	Clade	Provenance	Strata	Location
<i>Chlorostilbon mellisugus phoeopygus</i>	Emerald	Confluences	Understory	
<i>Chrysuronia oenone</i>	Emerald	MNHN	Canopy	Ecuador
<i>Coeligena coeligena</i>	Brilliant	MNHN	Understory	Ecuador
<i>Coeligena iris hesperus</i>	Brilliant	MNHN	Understory	Ecuador
<i>Coeligena iris iris</i>	Brilliant	MNHN	Understory	Ecuador
<i>Coeligena lutetiae</i>	Brilliant	MNHN	Understory	Ecuador
<i>Coeligena torquata fulgidigula</i>	Brilliant	MNHN	Understory	Ecuador
<i>Coeligena torquata torquata</i>	Brilliant	MNHN	Understory	Ecuador
<i>Coeligena wilsoni</i>	Brilliant	MNHN	Understory	Ecuador
<i>Colibri coruscans</i>	Mangoe	MNHN	Canopy	Ecuador
<i>Colibri delphinae</i>	Mangoe	MNHN	Canopy	Ecuador
<i>Colibri thalassinus</i>	Mangoe	MNHN	Canopy	Colombia
<i>Damophila julie</i>	Emerald	MNHN	Understory	Ecuador
<i>Discosura conversii</i>	Coquette	MNHN	Canopy	Ecuador
<i>Discosura langsdorffi</i>	Coquette	Confluences	Canopy	
<i>Discosura popelairii</i>	Coquette	MNHN	Canopy	Ecuador
<i>Doryfera johannae</i>	Mangoe	MNHN	Understory	Ecuador
<i>Doryfera ludovicae</i>	Mangoe	MNHN	Understory	Ecuador
<i>Ensifera ensifera</i>	Brilliant	MNHN	Understory	Ecuador
<i>Eriocnemis alinae</i>	Brilliant	MNHN	Understory	Ecuador
<i>Eriocnemis luciani</i>	Brilliant	MNHN	Understory	Ecuador
<i>Eriocnemis mosquera</i>	Brilliant	Confluences	Understory	
<i>Eriocnemis nigrivestis</i>	Brilliant	MNHN	Understory	Ecuador
<i>Eriocnemis vestita smaragdinicollis</i>	Brilliant	MNHN	Understory	Ecuador
<i>Eutoxeres aquila</i>	Hermit	MNHN	Understory	Ecuador
<i>Eutoxeres condamini</i>	Hermit	Confluences	Understory	
<i>Florisuga mellivora</i>	Topazes	MNHN	Canopy	Ecuador
<i>Glaucis aeneus</i>	Hermit	MNHN	Understory	
<i>Glaucis hirsutus affinis</i>	Hermit	MNHN	Understory	Peru
<i>Haplophaedia aureliae russata</i>	Brilliant	Confluences	Understory	
<i>Haplophaedia lugens</i>	Brilliant	Confluences	Understory	
<i>Heliangelus amethysticollis laticlavus</i>	Coquette	Confluences	Understory	
<i>Heliangelus exortis</i>	Coquette	MNHN	Understory	Ecuador
<i>Heliangelus micraster</i>	Coquette	MNHN	Understory	Ecuador


Species	Clade	Provenance	Strata	Location
<i>Heliangelus strophianus</i>	Coquette	MNHN	Understory	Ecuador
<i>Heliangelus viola</i>	Coquette	MNHN	Understory	Ecuador
<i>Heliodoxa aurescens</i>	Brilliant	MNHN	Understory	Colombia
<i>Heliodoxa imperatrix</i>	Brilliant	MNHN	Understory	Ecuador
<i>Heliodoxa jacula jamesoni</i>	Brilliant	MNHN	Understory	Ecuador
<i>Heliodoxa leadbeateri</i>	Brilliant	MNHN	Understory	Ecuador
<i>Heliodoxa rubinoides aequatorialis</i>	Brilliant	MNHN	Understory	Ecuador
<i>Heliodoxa schreibersii</i>	Brilliant	MNHN	Understory	Ecuador
<i>Heliomaster longirostris</i>	MtGem	MNHN	Canopy	Colombia
<i>Heliothyx auritus</i>	Mangoe	MNHN	Canopy	Ecuador
<i>Heliothyx barroti</i>	Mangoe	MNHN	Canopy	Ecuador
<i>Klais guimeti</i>	Emerald	MNHN	Understory	Ecuador
<i>Lafresnaya lafresnayi gayi</i>	Brilliant	Confluences	Understory	
<i>Lesbia nuna gracilis</i>	Coquette	MNHN	Canopy	Ecuador
<i>Leucippus baeri</i>	Emerald	Confluences	Understory	
<i>Leucippus chlorocercus</i>	Emerald	Confluences	Canopy	
<i>Lophornis chalybeus verreauxi</i>	Coquette	MNHN	Canopy	Colombia
<i>Metallura baroni</i>	Coquette	MNHN	Canopy	Ecuador
<i>Metallura tyrianthina tyrianthina</i>	Coquette	MNHN	Understory	Ecuador
<i>Metallura williami primolina</i>	Coquette	MNHN	Canopy	Ecuador
<i>Myrmia micrura</i>	Bee	MNHN	Canopy	Peru
<i>Ocreatus underwoodii melanantherus</i>	Brilliant	MNHN	Understory	Ecuador
<i>Opisthoprora euryptera</i>	Coquette	Confluences	Understory	
<i>Oreotrochilus chimborazo chimborazo</i>	Coquette	MNHN	Understory	Ecuador
<i>Oreotrochilus chimborazo jamesonii</i>	Coquette	MNHN	Understory	Ecuador
<i>Patagona gigas</i>	Patagona	MNHN	Canopy	Ecuador
<i>Phaethornis atrimentalis atrimentalis</i>	Hermit	Confluences	Understory	
<i>Phaethornis bourcierii</i>	Hermit	MNHN	Understory	
<i>Phaethornis griseogularis</i>	Hermit	MNHN	Understory	Ecuador
<i>Phaethornis guy</i>	Hermit	MNHN	Understory	Ecuador
<i>Phaethornis hispidus</i>	Hermit	Confluences	Understory	
<i>Phaethornis longirostris</i>	Hermit	Confluences	Understory	
<i>Phaethornis malaris</i>	Hermit	Confluences	Understory	
<i>Phaethornis ruber</i>	Hermit	Confluences	Understory	

Species	Clade	Provenance	Strata	Location
<i>Phaethornis syrmatophorus columbianus</i>	Hermit	MNHN	Understory	Ecuador
<i>Phaethornis yaruqui yaruqui</i>	Hermit	MNHN	Understory	Ecuador
<i>Phlogophilus hemileucurus</i>	Coquette	MNHN	Understory	Ecuador
<i>Polytmus theresiae leucorrhous</i>	Mangoe	MNHN	Understory	Ecuador
<i>Pterophanes cyanopterus</i>	Brilliant	MNHN	Understory	Ecuador
<i>Ramphomicron microrhynchum</i>	Coquette	MNHN	Canopy	Ecuador
<i>Schistes geoffroyi</i>	Mangoe	MNHN	Understory	Ecuador
<i>Taphrospilus hypostictus</i>	Emerald	MNHN	Understory	Ecuador
<i>Thalurania fannyi verticeps</i>	Emerald	MNHN	Understory	Ecuador
<i>Thalurania furcata viridipectus</i>	Emerald	MNHN	Understory	
<i>Thaumastura cora</i>	Bee	Confluences	Canopy	
<i>Threnetes leucurus cervinicauda</i>	Hermit	Confluences	Understory	
<i>Threnetes ruckeri</i>	Hermit	MNHN	Understory	Ecuador
<i>Urochroa bougueri</i>	Brilliant	Confluences	Understory	
<i>Urochroa bougueri leucura</i>	Brilliant	Confluences	Understory	
<i>Urosticte benjamini</i>	Brilliant	MNHN	Understory	Ecuador
<i>Urosticte ruficrissa</i>	Brilliant	Confluences	Understory	


Table 4: Measurement of intraspecific variability for hue (H1) and brightness (B2) by computing the coefficient of variation (standard deviation divided the average) on an independent dataset of hummingbirds living in French Guiana (Gomez *et al*, unpublished data), in which between 2 and 5 males (last column) were measured for each species. The measurement protocol differs slightly from the one used in this study, because we used a bifurcated probe at 45°, which may increase the intraspecific variability in brightness. In spite of the apparently high values of the coefficient of variation for brightness, it remains highly repeatable as estimated by the intra-class coefficient [90]: $R = 0.809$, $p < 0.0001$ for brightness and $R = 0.661$, $p < 0.0001$ for hue.

Species	CV brightness (%)	CV_ hue (%)	n
<i>Anthracothorax nigricollis</i>	20.57	2	3
<i>Calliphlox amethystina</i>	24.37	1.13	5
<i>Campylopterus largipennis</i>	17.43	0.1	2
<i>Chlorestes notatus</i>	19.79	1.96	5
<i>Discosura longicauda</i>	26.27	2.51	5
<i>Florisuga mellivora</i>	22.41	2.1	5
<i>Glaucis hirsuta</i>	33.75	0	4
<i>Heliomaster longirostris</i>	26.88	2.26	4
<i>Heliothyx aurita</i>	22.82	1.26	5
<i>Hylocharis cyanus</i>	29.75	2.55	3

Species	CV brightness (%)	CV_ hue (%)	n
<i>Hylocharis sapphirina</i>	23.32	3.36	4
<i>Lophornis ornatus</i>	23.38	1.55	5
<i>Phaethornis longuemareus</i>	18.59	0.15	4
<i>Phaethornis malaris</i>	21.44	0.1	2
<i>Phaethornis superciliosus</i>	27.88	0.1	5
<i>Thalurania furcata</i>	84.13	12.4	2
<i>Threnetes niger</i>	16.42	0.1	2
<i>Topaza pella</i>	23.04	1.83	5



Supplementary figure 2: Phylogenetic coverage of the *Trochilidae* family in our dataset (species and lineages in red).



Supplementary figure 3: Study sites locations (red dots) plotted on an altitudinal map of Ecuador. Communities outside the borders of the map are on islands or close enough to Ecuador borders to be taken into account in our study.





Figure 1: (a) community total colour volume and (b) brightness range increase with the number of species within the community. Each point is a community. The black solid line represents the mean value of (a) colour volume or (b) brightness range from 10 000 random communities with a given species count (null model) and the gray ribbon represents two standard deviations from the mean of the null model.

Variable	Diffuse		Directional		Both	
	R	p-value	R	p-value	R	p-value
x	0.734	0.002	0.877	<0.0001	0.925	<0.0001
Hue y	0.923	<0.0001	0.785	0.0006	0.951	<0.0001
z	0.780	0.0006	0.880	<0.0001	0.940	<0.0001
Brightness	0.411	0.090	0.055	0.48	0.373	0.04

Supplementary table 1: We quantified the repeatability R (intra-class coefficient ICC) and the related p-value by bootstrapping using the `rptR` R package [90] of indices used in this study by performing the same measurements twice on two patches for 12 species (*Coeligena torquata*, *Colibri coruscans*, *Doryfera ludovicae*, *Heliangelus strophianus*, *Heliodoxa jamesonii*, *Heliosthryx barroti*, *Juliamyia julie*, *Lesbia nuna*, *Metallura tyrianthina*, *Ramphomicron microrhynchum*, *Schistes albogularis*, *Urosticte benjamini*). Patches were selected to be of similar hue from a human point of view.



Supplementary figure 4: Colour of the 8 main patches for each species in our dataset. The colour corresponds to the colour in the human visual system (CIE10). The x-axis on the phylogeny is in millions years.

variable	value	Crown	Back	Rump	Tail	Throat	Breast	Belly	Wing
Hue	τ_{st}	-0.0073	0.055	0.055	0.044	0.027	0.03	0.05	0.058
	$p_{\tau_{st}<0}$	0.4	1	1	1	0.9	0.9	1	1
	$p_{\tau_{st}>0}$	0.6	0.01	0.01	0.03	0.09	0.06	0.005	0.006
	$d\mathcal{T}_{st}$	0.0099	0.026	-0.0021	0.0034	-0.0021	-0.0032	-0.01	0.00073
	$p_{\tau_{st}<0}$	1	1	0.8	1	0.9	0.3	<0.0001	1
	$p_{\tau_{st}>0}$	<0.0001	<0.0001	1	0.2	1	1	1	1
Brightness	τ_{st}	-0.021	0.0078	0.0032	-0.0064	0.00015	0.0041	-0.0031	0.0091
	$p_{\tau_{st}<0}$	0.1	0.7	0.6	0.5	0.5	0.6	0.5	0.6
	$p_{\tau_{st}>0}$	0.9	0.3	0.4	0.5	0.5	0.4	0.5	0.4
	$d\mathcal{T}_{st}$	-0.0014	0.0028	0.00037	0.00068	0.013	0.023	0.007	-0.0058
	$p_{\tau_{st}<0}$	0.3	1	0.9	1	1	1	1	0.2
	$p_{\tau_{st}>0}$	0.8	0.7	0.7	0.8	<0.0001	<0.0001	0.002	1
Hue shift	τ_{st}	-0.007	0.051	0.052	0.043	0.027	0.029	0.049	0.058
	$p_{\tau_{st}<0}$	0.4	1	1	1	0.9	0.9	1	1
	$p_{\tau_{st}>0}$	0.6	0.01	0.01	0.03	0.08	0.06	0.006	0.006
	$d\mathcal{T}_{st}$	0.0087	0.0059	-0.0068	-0.006	-0.0033	0.0023	-0.0098	-0.0018
	$p_{\tau_{st}<0}$	1	1	0.005	0.01	0.6	1	<0.0001	1
	$p_{\tau_{st}>0}$	<0.0001	0.03	1	1	1	0.9	1	1

Supplementary table 2: Numerical values for τ_{st} and decoupled τ_{st} (denoted $d\mathcal{T}_{st}$). P-values were computed by comparison of the actual value with the null distribution (obtained by randomisation of the communities using method 1s of Hardy [67]). Significant p-values are in bold and green. Positive values of $d\mathcal{T}_{st}$ indicate phenotypic clustering whereas negative values indicate overdispersion.