

1 Distribution of iridescent colours in hummingbird communities results
2 from the interplay between selection for camouflage and communication

3 Hugo Gruson^{*a}, Marianne Elias^b, Juan L. Parra^c, Christine Andraud^d, Serge Berthier^e, Claire
4 Doutrelant^a, and Doris Gomez^{a,e}

5 ^aCEFE, Univ Montpellier, CNRS, Univ Paul Valéry Montpellier 3, EPHE, IRD, Montpellier, France

6 ^bISYEB, CNRS, MNHN, Sorbonne Université, EPHE, 45 rue Buffon CP50, Paris, France

7 ^cGrupo de Ecología y Evolución de Vertebrados, Instituto de Biología, Universidad de Antioquia,
8 Medellín, Colombia

9 ^dCRC, MNHN, Ministère de la Culture et de la Communication, CNRS, Paris, France

10 ^eINSP, Sorbonne Université, CNRS, Paris, France

11 **Abstract**

12 Identification errors between closely related, co-occurring, species may lead to misdirected social interactions such
13 as costly interbreeding or misdirected aggression. This selects for divergence in traits involved in species identification
14 among co-occurring species, resulting from character displacement. On the other hand, predation may select for
15 crypsis, potentially leading co-occurring species that share the same environment ~~to have and predators to have a~~
16 similar appearance. ~~Few~~ However, few studies have explored how these antagonistic processes influence colour at the
17 community level. Here, we assess colour clustering and overdispersion in multiple hummingbird communities across
18 Ecuador and identify the processes at stake by controlling for species phylogenetic relatedness. In hummingbirds, most
19 colours are iridescent structural colours, defined as colours that change with the illumination or observation angle.
20 Because small variations in the underlying structures can have dramatic effects on the resulting colours and because
21 iridescent structures can produce virtually any hue and brightness, we expect iridescent colours to respond finely to
22 selective pressures. Moreover, we predict that hue angular dependence – a specific aspect of iridescent colours – may
23 be used as an additional channel for species recognition. In our hummingbird assemblages in Ecuador, we find support
24 for colour overdispersion in specific body patches at the community level even after controlling for the phylogeny,
25 especially on iridescence-related traits, suggesting character displacement among co-occurring species. We also find
26 colour clustering at the community level on ~~patches dorsal patches, suspected to be~~ involved in camouflage, ~~which~~
27 ~~may counter-balance the effect of character displacement~~ suggesting that the same cryptic colours are selected among
28 co-occurring species.

*hugo.gruson@normalesup.org

29 **Keywords:** Reproductive Character Displacement, Agonistic Character Displacement, Camouflage, Structural Colours,
30 Angle-Dependent Colouration, Community structure, Ecuador

31

32 Colour is a complex communication channel widespread among various taxa and involved in many ecological and
33 evolutionary processes [1]. It can be described by multiple variables, including hue (colour in its common sense, such as
34 red, green, blue, etc.) and brightness (average level of grey of a colour, i.e. whether the object is light or dark). Colours
35 can be produced by two non-mutually exclusive means: pigmentary colours are produced by the selective absorption of
36 incoming light by pigments, while structural colours are produced by the interaction of incoming light with nanostructures,
37 causing diffraction, interferences or scattering [2]. Among structural colours, iridescent colours are characterised by a shift
38 in hue with changes in illumination or observation angle [3]. Iridescent colours are found in many bird families such as
39 Anatidae (ducks) Phasianidae (fowls), Sturnidae (starlings), or Trochilidae (hummingbirds), and thought to be involved
40 in numerous adaptations [4]. But evolution of iridescent colours at the community level remains poorly understood. Yet,
41 they may display evolutionary patterns that differ from non-iridescent colours. Indeed, as opposed to other types of
42 colours, iridescent colours can produce virtually any hue and are expected to respond more readily and finely to selection,
43 because large changes of hue can be achieved by small changes in the underlying structures [5]. They can also result in
44 directional colours only seen at specific angles, as well as highly reflective colours [6].

45 Because colours are involved in many different ecological processes, they are subject to multiple selection pressures,
46 often with opposite effects [7]. ~~For example~~ Colour may indeed increase or decrease detectability of an animal depending
47 on the colour contrast with its surroundings. In particular, colour can reduce predation risk via crypsis or aposematism
48 or serve as a means of species identification. In this case, two opposite evolutionary forces act on colours: (i) On the
49 one hand, species living in the same environment are likely experiencing similar selective pressures, such as predation.
50 The environment is characterised by ambient light and vegetation, which both influence greatly which colours are poorly
51 detectable and which colours are highly detectable [8, 9]. We thus expect co-occurring species to ~~converge in coloration and~~
52 ~~harbour poorly detectable~~ harbour the same, poorly detectable, colours as this would decrease the risk of being detected
53 by predators. ~~Colour,~~ thereby causing a clustering pattern in colouration at the community level, all else being equal.
54 This colour clustering can result from convergence between sympatric species (evolutionary process), from environmental
55 filtering (ecological process), i.e. species ~~assortment~~ sorting locally according to the traits they harbour, or a mixture of
56 the two (detailed in table 1). (ii) On the other hand, sympatric closely-related species are more likely to face problems
57 of species recognition, eventually resulting in reproductive interference - a phenomenon where an individual courts or
58 mates with individuals of another species, producing no offspring or low fertility hybrids, leading to costly interbreeding
59 [10]. Species misidentification can also lead to misdirected aggression and costly fighting when individuals compete over
60 resources or territories. Hence, any feature that would enhance species recognition is expected to be selected for. In this
61 context, closely related species living in sympatry should be under strong selective pressure to diverge in traits involved
62 in communication, if divergence enhances species recognition. Divergence can result from a process called character
63 displacement (RCD for reproductive character displacement, ACD for agonistic character displacement; evolutionary

64 process) [11–13] or from species ~~assortment-sorting~~ (ecological process). For ACD, it is worth ~~noticing-noting~~ that traits
65 are expected to diverge only in case of moderate ecological competition, whereas they should converge in case of high
66 competition [13, 14]. Multiple empirical studies have shown character displacement for songs (e.g. Gerhardt [15] in frogs
67 and Grant and Grant [16] in birds), or olfactory signals [17]. However, fewer studies have looked at divergence in colour
68 patterns (but see Sætre et al. [18], Naisbit et al. [19], Lukhtanov et al. [20], Martin et al. [21], Doutrelant et al. [22], and
69 Hemingson et al. [23]). Almost all these studies were at the species level, and at best involved comparison between closely
70 related species. Many of them also did not use objective spectrometry measurements and instead relied on human vision,
71 which ~~is likely to have biased their results~~ did not allow them to analyse colours as perceived by the intended
72 receiver, in the case of this study: birds [24–27].

73 In birds, it has been ~~argued~~ showed that colouration is under different selective pressures depending on the body patch
74 location: dorsal patches ~~are~~, which are exposed to aerial predators, are mainly involved in camouflage while ventral and
75 facial patches are mainly involved in communication ~~[7, 28]~~ [7, 29]. In this study, we test this hypothesis for iridescent
76 colours at the community level by looking at phenotypic structure in hummingbird local assemblages across different body
77 parts. Hummingbirds are an interesting study system to test this hypothesis as various published accounts of sexual
78 displays and aggressive encounters among hummingbirds have made clear that certain feather patches such as the crown
79 and throat are consistently used during these displays [30–33]. On the other hand, colours displayed on the dorsal side
80 of hummingbirds tend to resemble background colours and thus have been suggested to be cryptic [28]. Accordingly,
81 we predict that co-occurring hummingbird species should display similar hues on dorsal patches, leading to phenotypic
82 clustering of hues (i.e. co-occurring species are more similar than expected by chance, prediction 1) and different hues
83 on ventral patches, resulting in a phenotypic overdispersion pattern (i.e. co-occurring species are more dissimilar than
84 expected by chance, prediction 2). For brightness, we can formulate two alternative predictions: on the one hand, it
85 might evolve in the same way as hue, also because of reproductive character displacement and selection for camouflage,
86 leading to the same outcome as for hue (prediction 3, equivalent to predictions 1 and 2 but for brightness). On the other
87 hand, because brightness level positively correlates with signal conspicuousness, poorly detectable signals have similar
88 brightness, and highly detectable signals have similar brightness. Hence, we may instead expect that species co-occurring
89 should converge for brightness on all patches (prediction 3bis) if the same patches are involved in the same ecological
90 process (communication or camouflage).

91 Compared to other types of colouration, iridescent colours might enable species recognition on another dimension in
92 the sensory space. Two species can have the same hue or brightness at a given angle but can differ at another angle, via
93 an additional variable we call "hue shift". Because hue shift cannot be seen at ~~large-long~~ distances, it may allow species
94 to diverge without interfering with camouflage against predators ~~[4]~~ [4, 34]. Accordingly, we predict overdispersion for hue
95 shift not only on ventral patches, but also on dorsal patches (prediction 4). However, hue shift is often highly correlated
96 with hue due to the optics underlying iridescence (Dakin and Montgomerie [35] for example reported $R^2 \geq 0.95$ for the
97 correlation between hue and hue shift). We test this correlation with the data from this article and discuss how it may
98 impact our results.

99 At the community level, we predict that community colour volume (also known as functional richness **FRic** in functional
100 ecology [36]) and brightness range increase with species richness more than expected in a random species assemblage (null
101 model) because co-occurring species would use different colours (hue or brightness) (prediction 5).

102 Here we test our five predictions by quantifying both iridescent and non-iridescent colours of 189 hummingbird assem-
103 blages in Ecuador that include 112 species and span a large variety of habitats, and by assessing the phenotypic structure
104 (clustering, random distribution, overdispersion of colours) and ~~correct that for the expectation given~~ investigate the
105 underlying processes by taking into account species phylogenetic relatedness within these assemblages. Comparing the
106 uncorrected and the phylogenetically-corrected phenotypic structure of hummingbird communities will allow us to identify
107 which mechanisms (character displacement, species ~~assortment~~ sorting with mutual exclusion of similar species, environ-
108 mental filtering; as detailed in table 1) underlie the community structure of iridescent colours in hummingbirds.

109 **Materials and methods**

110 All scripts and data used to produce the results and figures from this article are available at [https://doi.org/10.5281/](https://doi.org/10.5281/zenodo.3355444)
111 [zenodo.3355444](https://doi.org/10.5281/zenodo.3355444)

112 **Community data**

113 Hummingbirds are particularly suited as a study system to explore the possible effect of reproductive character displace-
114 ment on iridescent colours because (i) they display a large variety of hues [37] and all species harbour some iridescent
115 patches, many of which have a very strong angular dependence, rapidly shifting from e.g. pink to green or black [38,
116 39] (but note that many hummingbirds species also have non-iridescent, pigmentary, patches), (ii) they belong to a very
117 speciose family whose phylogeny is well established and readily available [40, 41], (iii) they live only in the Americas,
118 especially in the tropics where numerous species can coexist locally [37] ~~and~~ (iv) there is an extensive documentation of
119 hybridisation between co-occurring species (see for example [42, 43] for our region of interest), which creates the perfect
120 opportunity to study reproductive interference and (v) almost all species are available in museum collections and their
121 colour can be objectively measured using spectrometric measurements [44].

122 Presence/absence data for hummingbird assemblages at 189 sites in Ecuador (see map in fig. S3) were compiled
123 from data in peer-reviewed papers and reports from environmental organisations [45]. These sites cover a large variety
124 of elevation ranges (fig. S3) and habitats [45, 46]. This dataset was previously thoroughly reviewed by comparing the
125 observations with the known elevational and geographical ranges of each species [46] and includes observations of 112 of
126 the 132 hummingbirds species found in Ecuador [47].

127 **Colour measurements and analyses**

128 For each one of the 112 species, we borrowed one ~~male~~ adult male in good condition from either the Museum National
129 d’Histoire Naturelle (MNHN) in Paris or the Musée des Confluences, in Lyon (full list in Online Supplementary Informa-

tion). [We ensured that the specimen colouration was representative of the other specimens available in the collections to the human eye.](#) When multiple subspecies were living in the area where presence was recorded, we randomly picked one of them. We consistently took spectral reflectance measurements on the 8 following patches (described in fig. S1): crown, back, rump, tail, throat, breast, belly, wing. We also made additional measurements on patches that visually differed in colouration from these 8 main ones, as in Gomez and Théry [7] and Doutrelant et al. [22].

We measured reflectance using a setup similar to Meadows et al. [48], relying on the use of two separate optical fibres. Light was conducted from an Oceanoptics DH-2000 lamp emitting over the 300-700 nm range of wavelengths to which birds are sensitive [49] to the sample through an illuminating FC-UV200-2-1.5 x 100 optical fibre (named illumination fibre). Light reflected by the sample was then collected by a second identical optical fibre (named collection fibre) and conducted toward an Oceanoptics USB4000 spectrophotometer (used with the SpectraSuite 2.0.162 software). This setup allows for a precise independent rotation of the illumination and the collection fibres, necessary for the measurements of iridescent colours [6]. For more details about the measurement conditions as recommended in White et al. [50], see [the supplementary materials \(ESM\).](#)

For every patch, we recorded a first reflectance spectrum at the position of the fibres which maximised total reflectance. To measure hue angle dependency (iridescence), we then moved both fibres 10° away from the previous position and recorded a second spectrum, as in Meadows et al. [51]. More recent measurement methods revealed that it would be more accurate to keep the angular span between the illumination and collection fibres constant [52]. We however confirmed that this did not impact our results by running our analyses once with all data and once with only data at a given angular span (which represented 94% of the total data). All measurements were performed in a dark room with temperature control. Recorded spectra were normalised by an Avantes WS-1 white standard and a measurement with the lamp shut down (dark reference) and integration times were determined for each sample as to maximise the intensity of the signal without saturating the spectrometer.

Final values were averaged over 5 consecutive measurements and spectra were smoothed using a loess algorithm and interpolated every ~~1 nm~~ [1 nm](#) and negative values were set to zero using the R package `pavo` ~~[53]~~[\[54\]](#).

We analysed spectra using Endler and Mielke [55] model with relative quantum catches Q_i (without Fechner's law). All birds are tetrachromats and can see light with wavelengths from 300 to 700 nm, which includes ultra-violet light (UV) [56]. But different bird species vary in their sensitivity [57]: some are ~~called~~ UV-sensitive (UVS) while others are violet-sensitive (VS). Literature on colour vision in hummingbirds suggests that both types are found within the family (see Chen and Goldsmith [49] and Herrera et al. [58] for UVS species and Ödeen and Håstad [59] for VS species). Because we did not have enough information to compute ancestral states and vision type for all species in our study and because it was found to have little influence in previous studies ~~[7]~~[\[7, 29\]](#), we ran our analyses as if all species were VS, using the spectral sensitivities of a typical VS bird, *Puffinus pacificus* [60], [whose photoreceptor absorbances match closely those reported for hummingbirds \[59\]](#). We used different illuminants defined in Endler [8], depending on the habitat of the species described in Stotz et al. [61] (detailed in SI): "large gaps" illumination was used for species living in the canopy while "forest shade" was used for species living in the understory. Hue was a tridimensional variable defined by the position (x , y and z) of

165 the reflectance spectrum in the tetrahedron representing bird colour vision space [55] and brightness was defined as in
166 Endler and Mielke [55] (perceived intensity of colour, also sometimes referred to as luminance). We ensured that all indices
167 were repeatable (table S1) ~~using the by measuring twice the same individual and patch on 20 patches and computing the~~
168 intra-class coefficient (ICC) with the rptR R package [62]. We add another variable to describe iridescence: hue shift,
169 defined as the difference between hue at maximum reflectance and hue at 10° away from maximum reflectance, in a similar
170 fashion to Dakin and Montgomerie [35]. Because it is the difference of two tridimensional variables (hue at the position
171 where reflectance was maximum and hue at 10° away), hue shift is tridimensional as well. Dakin and Montgomerie [35]
172 found a high correlation between hue and hue shift at the intraspecific level in the peacock *Pavo cristatus*, we also report
173 a high correlation at the interspecific level in hummingbirds by performing a linear regression in \mathbb{R}^3 between hue and hue
174 shift ($R^2 = 0.51$, $F(3; 1372) = 469.7$, $p < 0.0001$). New measurement methods have since been developed and propose a
175 new definition for hue shift which is not correlated to hue but they were not available at the time of this study [52].

176 We analysed the colour volume for each species by measuring the convex hull volume of all colour patches on the
177 bird, as suggested in Stoddard and Prum [63]. We compared the relationship between the colour volume of a community
178 and the number of species within this community relative to a null model (prediction 5) obtained by creating random
179 assemblages from a species pool containing all species from all communities. In other words, actual assemblages are
180 compared to fictional assemblages with exactly the same number of species but no abiotic or biotic constraints on the
181 species composition.

182 However, the colour volume does not ~~differentiate the different patches~~ take into account the patch location on the
183 bird body, raising several concerns. First, two species could use the same colour but at different places on their body.
184 They would then look different to an observer but not identified as such in this analysis. Additionally, we expect different
185 evolutionary signals on different patches, that could even each other out, and blur the outcome at the bird level. For these
186 reasons, we also performed our analyses separately for each one of the following eight patches: crown, back, rump, tail,
187 throat, breast, belly, wing (locations shown in fig. S1).

188 Trochilidae phylogeny and comparative analyses

189 A distribution of 100 phylogenetic trees of the Trochilidae family was downloaded from birdtree.org [40] to take into
190 account phylogenetic uncertainty in the comparative analyses [64]. The 112 species included in this study constitute a
191 fairly even sampling of the hummingbird phylogeny (fig. S2).

192 We used the method developed by Hardy and Senterre [65] and Baraloto et al. [66] to analyse respectively the phyloge-
193 netic (Π_{ST}) and phenotypic (τ_{ST}) structures of the hummingbird communities of Ecuador (clustering or overdispersion).
194 This method relies on computing indices inspired by the Simpson index and the fixation index F_{ST} , comparing the ob-
195 served diversity within and between the communities. For phylogeny, Π_{ST} can reveal phylogenetic clustering ($\Pi_{ST} > 0$)
196 or phylogenetic overdispersion ($\Pi_{ST} < 0$) within communities. Likewise, for phenotypic traits, τ_{ST} can reveal phenotypic
197 clustering ($\tau_{ST} > 0$) or phenotypic overdispersion ($\tau_{ST} < 0$) within communities. Statistical significance of overdispersion
198 or clustering is obtained from comparing the observed value to that obtained from 1000 random communities (created by

199 drawing from the total species pool, using algorithm 1s from Hardy [67], which keeps the local species richness per site
200 constant). This approach compares the phenotypic structure to what would be expected by chance.

201 To disentangle the relative effect of ecological (species ~~assortment~~sorting) and evolutionary mechanisms (selection),
202 we also perform our analyses by taking into account the phylogenetic relationships between species. If the species in
203 the community are more clustered or overdispersed than expected given their phylogenetic relationships, this is taken as
204 evidence that the trait has not evolved in a Brownian fashion (detailed in table 1). To this end, we used the `decouple`
205 function [68], which returns phylogenetically predicted and residual trait values by performing a linear regression of
206 individual trait values explained by the phylogeny. We computed the value of τ_{ST} on trait values decoupled from the
207 phylogeny. This value is hereafter denoted $dc\tau_{ST}$. Similarly to the classical τ_{ST} , the sign of $dc\tau_{ST}$ indicates phenotypic
208 clustering ($dc\tau_{ST} > 0$) or overdispersion ($dc\tau_{ST} < 0$) once the effect of the phylogenetic structure of the communities has
209 been ~~removed~~decoupled.

	$\tau_{ST} < 0$ Phenotypic overdispersion	$\tau_{ST} = 0$ No community structure	$\tau_{ST} > 0$ Phenotypic clustering
$d\tau_{ST} < 0$	Co-occurring species are less similar than expected by chance because of character displacement.	Co-occurring species are not more by chance despite character displacement because closely related species co-occur more often than expected at random (phylogenetic clustering: $\Pi_{ST} > 0$).	Co-occurring species are more similar than expected by chance despite character displacement because closely related species co-occur more often than expected at random (phylogenetic clustering: $\Pi_{ST} > 0$).
Character displacement (divergence): co-occurring species are more dissimilar than expected given their phylogenetic relationships, which means they evolved towards dissimilarity in their colours.			
$d\tau_{ST} = 0$ Brownian trait evolution	Competitive exclusion: co-occurring species are more dissimilar than expected by chance because distantly-related (and therefore dissimilar) species co-occur more often than expected at random (phylogenetic overdispersion; $\Pi_{ST} < 0$).	Co-occurring species are not more similar nor more different than expected by chance or than predicted given their phylogenetic relationships.	Environmental filtering: co-occurring species are more similar than expected by chance because closely-related (and therefore similar) species co-occur more often than expected at random (phylogenetic clustering: $\Pi_{ST} > 0$).
$d\tau_{ST} > 0$	Co-occurring species are less similar than expected by chance despite evolutionary convergence because distantly-related species co-occur more often than expected at random (phylogenetic overdispersion; $\Pi_{ST} < 0$).	Co-occurring species are neither more nor less similar than expected by chance despite evolutionary convergence because distantly-related species co-occur more often than expected at random (phylogenetic overdispersion; $\Pi_{ST} < 0$).	Co-occurring species are more similar than expected by chance because of evolutionary convergence.
Evolutionary convergence : co-occurring species are more similar than expected given their phylogenetic relationships, which means they evolved towards similarity in their colours.			

Table 1: Summary of the different evolutionary and ecological scenarios and their results in terms of values of τ_{ST} and decoupled $d\tau_{ST}$.

Analyses performed on a tree distribution (Π_{ST} and $dc\tau_{ST}$) with n trees return a distribution of n statistics values and n p-values p_i . We summarised this information by computing the median of the statistics and the overall p-value p by using Jost's formula [69]:

$$p = k \sum_{i=0}^{n-1} \frac{(-\ln(k))^i}{i!} \quad \text{where } k = \prod_{i=1}^n p_i \quad (1)$$

Results

We find a strong phylogenetic clustering within communities ($\Pi_{ST} = 0.062 > 0$, $p < 0.0001$), indicating that co-occurring species are more closely related than expected by chance.

Phenotypic structure of the communities (predictions 1 - 4)

When looking at the bird entire body (when all patches are included simultaneously) by computing the overlap of the colour volumes, we did not find any phenotypic structure.

When the different major patches (crown, back, rump, tail, throat, breast, belly and wing) are examined separately (table 2 and table S2), we find clustering ($\tau_{ST} > 0$) in hue and hue shift on the back, rump, tail, belly and wing. Once we ~~remove-decouple~~ the effect of the shared evolutionary history ~~with the decouple function~~, we find clustering on the crown and the back ($dc\tau_{ST} > 0$) but overdispersion on the belly for both hue and hue shift ($dc\tau_{ST} < 0$). Hue shift is also overdispersed on the rump and the tail ($dc\tau_{ST} < 0$). There is no phenotypic structure on the throat, breast or wing for hue and hue shift nor on the rump or the tail for hue.

We find no phenotypic structure (neither clustering nor overdispersion) for brightness on any patches before phylogenetic correction. After phylogenetic correction, brightness values for the throat, breast and belly are clustered among co-occurring species ($dc\tau_{ST} > 0$) but show no phenotypic structure for the crown, the back, the wing and the tail.

Effect of community species richness on colour characteristics (prediction 5)

We found that the brightness range within a community increased in the same way as a null model built from random species assemblages (fig. 1b). For colour volume, we find some outliers with a higher colour volume than expected for community with the same number of species (fig. 1a).

Discussion

Our findings suggest that colour structure within hummingbird communities ~~results from a trade-off between selection for camouflage (likely results from the interplay between two selective pressures, acting in opposite directions: selection by the local environment (e.g. camouflage from predators, leading to phenotypic clustering) and species recognition (on dorsal patches, and selection for species recognition, leading to phenotypic overdispersion)). This balance between selective~~







Variable	Phenotypic structure (τ_{ST})	Decoupled phenotypic structure ($dc\tau_{ST}$)
Hue		
Brightness		
Hue shift (=iridescence)		

Table 2: Phenotypic structure of hummingbird communities for different variables (hue, brightness and hue shift) on the patches studied (crown, back, rump, tail, throat, breast, belly, wing; names and locations illustrated in fig. S1). Hue is a tridimensional variable defined by the reflectance spectrum position x , y and z in the tetrahedron representing avian colour space. Blue plus ~~sign signs~~ + ~~patterns~~ indicate significant phenotypic clustering (τ_{ST} or $dc\tau_{ST} > 0$), orange minus ~~sign signs~~ - indicate significant phenotypic overdispersion (τ_{ST} or $dc\tau_{ST} < 0$), and green ~~zero-zeros~~ 0 ~~patterns~~ represent the absence of phenotypic structure. The left column shows the raw phenotypic structure of the community ([columns in table 1](#)), which may be influenced by the phylogenetic structure while the right column shows the phenotypic structure of the community, decoupled from all effects caused by the phylogeny ([rows in table 1](#)). [By comparing the values of \$\tau_{ST}\$ and \$dc\tau_{ST}\$ for each trait colour variable \(hue, brightness and hue shift\), we can assume a probable evolutionary scenario for each patches, based on the explanation in table 1.](#) Exact values for the statistics are available in table S2.

~~pressure acting in opposite directions produces a complex phenotypic structure when looking at different patches on the body on ventral and facial patches.~~

Evidence for different evolutionary scenarios depending on patch location

At the entire bird level (i.e. when pooling together all patches), we did not find any phenotypic structure. ~~But as mentioned earlier, this was expected since different locations on the birds are expected to be under different selection regimes [7, 29].~~

~~As predicted in-~~ In accordance with our prediction 5, community colour volume (as estimated by the convex hull of hue and brightness range within a community) increases slightly faster with the number of species in the community than predicted by a null model. This suggests that co-occurring species in these communities tend to use ~~less different~~ more similar colours than expected by chance. However, this is not the cause for the majority of communities, where co-occurring species do not use more nor less similar colours than expected by chance. This is further confirmed by the absence of phenotypic structure on the colour volume and the brightness when the effect of the phylogeny is not ~~removed~~ decoupled.

This could be the consequence of similar selective pressures between the communities we studied, leading colours in all assemblages to be randomly determined. This is however not very likely because the communities we studied differ a lot in both their vegetation background and therefore in the pressure for crypsis [45] and in their species composition. A more likely hypothesis is that co-occurring species tend to use the same colours but not necessarily on the same patches, which would also explain the absence of phenotypic structure when we pool all patches without taking into account their location. This is confirmed by our analysis patch by patch, where we find either clustering or overdispersion depending on the location of the patch.

Selection for convergence and phenotypic clustering

In accordance with our predictions, co-occurring hummingbird species tend to have similar hues on patches more likely dedicated to camouflage (back, rump, tail, wing; prediction 1) but not on patches more likely used in communication (crown, throat, breast; prediction 2), as shown in table 2 and table S2. This new result for iridescence colours matches what has been previously described for non-iridescent colours ~~[7]~~ [7, 29]. The phenotypic clustering observed for hue on the rump, the tail and the wing vanishes after ~~removing~~ decoupling the clustering effect due to phylogenetic structure. This means that phenotypic clustering of hue on the rump, the tail and the wing is not caused by convergent evolution of co-occurring species but by environmental filtering, leading related, similar-looking species to live in the same area (as explained in table 1). This is confirmed by the high value of phylogenetic clustering. ~~Using different methods-~~ This sign of phylogenetic clustering completes the results from Graham et al. [45] on the same dataset, ~~Graham et al. [45] also found significant phylogenetic clustering in 37 communities and overdispersion in only one.~~ We showed that intra-community species relatedness is high compared to inter-community species relatedness (Π_{ST}), while they showed that intra-community species relatedness (Net Relatedness Index) is higher than expected from random assemblages in 71 % of the cases [45].

270 This phylogenetic clustering may be caused by a strong niche conservatism but our study cannot discriminate whether
271 such niche conservatism involves colour or other ecological traits. However, hummingbirds' costly hovering flight at high
272 elevation due to weaker lift caused by the decreasing atmospheric pressure [70–72] and high foraging specialisation [73]
273 likely contribute to this pattern. Alternatively, phylogenetic clustering could also be caused by a very low dispersal ability
274 of hummingbirds, but this remains quite unlikely as the rare studies on this topic have shown that different hummingbird
275 species display a wide variation in their dispersal ability [74, 75].

276 Contrary to our prediction 2, we also find clustering of hue on the belly before the use of the `decouple` function.
277 However, the fact that it turns into overdispersion after the use of the `decouple` function, and not simply into a random
278 phenotypic structure (as opposed to the rump, the tail and the wing mentioned just before), suggests this initial clustering
279 (right column in table 1) is mainly caused by environmental filtering on another trait but that hue on the belly is still
280 under selection for divergence (first row in table 1). This other trait may be the colour of another patch or other ecological
281 traits, as we explained previously.

282 We found a significant clustering of brightness on the throat, breast and belly after controlling for the phylogeny,
283 indicating that brightness on those patches is more similar than expected given the phylogeny among co-occurring species
284 (prediction 3bis). This suggests that the same patches have been selected to be involved either in communication or
285 in camouflage among species living in the same environment. This is seen after controlling for the phylogeny and it
286 is therefore not caused by the phylogenetic relatedness of co-occurring species. This is not surprising as many studies
287 showed the paramount importance of the throat in the courtship display of many hummingbird species [30–33, 76]. Two
288 main hypotheses can explain why co-occurring species tend to communicate (or camouflage themselves) using the same
289 patches: (i) There may be selective pressures for the use of specific patches in camouflage in a given environment (e. g.,
290 patches that are more exposed to predators' sight). (ii) Convergence in patches used in communication may be selected
291 because it improves competitor identification in the case of a strong ecological niche overlap (convergence by agonistic
292 character displacement as shown in Grether et al. [13] and Tobias et al. [77]).

293 All those results suggest a strong effect of the environment in the evolution of colour in agreement with McNaught
294 and Owens [78] who found that bird plumage colour was due to the light environment and not to reproductive character
295 displacement in Australian birds. However, we do not find clustering on all patches, which means that the effect of habitat
296 pressure is somehow limited or counterbalanced by reproductive or agonistic character displacement. On the contrary, for
297 some patches, we found patterns that are likely the result of character displacement.

298 **Character displacement and phenotypic overdispersion**

299 In agreement with our prediction 2, after removing-decoupling the effect of the phylogeny, there is overdispersion of hue
300 on the belly, likely caused by character displacement (table 1). At a completely different taxonomic scale, focusing on a
301 single hummingbird genus (*Coeligena*) with 11 species, Parra [28] also found that the belly was always involved in the
302 difference in hue between subspecies. It was sometimes even the only patch causing those differences, as for example
303 between *Coeligena torquata fulgidigula* and *Coeligena torquata torquata*. This suggests that the interspecific divergence

304 we found on the belly at the community level on the whole Trochilidae family can be observed at different geographic and
305 taxonomic scales, and even between subspecies of the same species.

306 As predicted, we also find more phenotypic overdispersion for hue shift than hue after ~~removing~~ decoupling the effect
307 of the phylogeny, for example, on the rump and on the tail (prediction 4). It is possible that hue shift is less sensitive
308 to selection for convergence because it may vary without disturbing camouflage efficacy. However, we did not find the
309 expected relaxing of clustering on hue shift on patches such as the back. This is likely caused by the fact that hue shift
310 is highly correlated with hue, as found in this study and in Dakin and Montgomerie [35], who used the same indices to
311 quantify iridescence. This correlation is due to the optics controlling iridescence, meaning that species that display similar
312 hues should also display the same hue shift if they use the same underlying multilayer structures. The fact that the
313 correlation is not perfect and that we nonetheless get different phenotypic patterns for hue and hue shift on some patches
314 suggests that co-occurring species use different multilayer structures (as recently confirmed by [79]), which can produce
315 different iridescent effects while displaying the same hue (functional convergence on hue).

316 Against our prediction 2, we did not find phenotypic overdispersion on any of the colour variables on patches such as
317 the throat or the crown, that are thought to be sexually selected and often used in courtship displays [30, 80]. Several
318 hypotheses can explain this fact: (i) The overdispersion on some patches (hue on the belly and hue shift on the rump and
319 tail) is sufficient to enable species recognition. (ii) The current phenotypic structure, which is neither overdispersed nor
320 clustered, on those patches is sufficient to enable species recognition. Indeed, the absence of phenotypic overdispersion
321 does not mean that species look the same. It simply means that colour differences between species living in the same
322 community and species in different communities occur in similar ranges. This difference may be sufficient to relax the
323 selective pressure towards reproductive character displacement. (iii) The pressure towards overdispersion is balanced by
324 habitat filtering (for both ventral and dorsal patches), resulting in no apparent phenotypic structure. The latter hypothesis
325 was also a candidate explanation of the pattern found by Martin et al. [21], where sympatric closely related species are
326 more divergent than allopatric ones, but only when the range overlap is limited. They suggested that local adaptation
327 could hinder divergence when species ranges was exactly the same. (iv) Species recognition is achieved by additional means
328 and divergence occurs on others traits, such as modified feathers [81], song [82, 83] or non-vocal noises [84–86] and size.
329 Notably, different species of hummingbirds can have very different courtship behaviour: leks for hermits [87, 88], dives
330 and shuttle displays for bees [32, 85, 89], for instance.

331 Taken together, our results suggest that hummingbird iridescent colours are determined by different evolutionary mech-
332 anisms depending on their location. Within a community, co-occurring hummingbird species tend to ~~use the same hue on~~
333 ~~dorsal, large, display the same hues on dorsal~~ patches probably because of ~~the evolutionary pressure for camouflages~~ selective
334 pressures related to the local environment, such as selection for crypsis by predators, causing phenotypic clustering at the
335 community level. This phenotypic clustering does not seem to be caused by adaptive convergence on colours but rather
336 by environmental filtering perhaps linked to other ecological traits such as elevation tolerance or flight ability. In spite
337 of such environmental filtering, character displacement leads to overdispersion for hue on the belly and hue shift on the
338 rump and the tail. Iridescence may therefore enable species recognition without affecting camouflage efficacy of birds, by

339 opening up a new dimension in the sensory space: hue shift.

340 Acknowledgments

341 This project heavily relied on museum specimens which were made available by the work of collection curators: Patrick
342 Boussès, Anne Previato, and Jérôme Fuchs (Muséum National d’Histoire Naturelle), Cédric Audibert and Harold Labrique
343 (Musée des Confluences).

344 Conflict of interest disclosure

345 The authors of this preprint declare that they have no financial conflict of interest with the content of this article. Marianne
346 Elias is part of the managing board of PCIEvolBiol and is one of the PCIEvolBiol recommenders.

347 References

- 348 [1] JW Bradbury and SL Vehrencamp. *Principles of Animal Communication*. 2. ed. OCLC: 759797180. Sunderland,
349 Mass: Sinauer Associates, 2011. 697 pp. ISBN: 978-0-87893-045-6.
- 350 [2] AR Parker. “515 Million Years of Structural Colour”. In: *Journal of Optics A: Pure and Applied Optics* 2.6 (2000),
351 R15–R28. DOI: 10.1088/1464-4258/2/6/201.
- 352 [3] P Vukusic. “Natural Photonics”. In: *Physics World* 17.2 (2004), p. 35. DOI: 10.1088/2058-7058/17/2/34.
- 353 [4] SM Doucet and MG Meadows. “Iridescence: A Functional Perspective”. In: *Journal of The Royal Society Interface*
354 6 (Suppl 2 2009), S115–S132. DOI: 10.1098/rsif.2008.0395.focus. pmid: 19336344.
- 355 [5] RO Prum. “Anatomy, Physics, and Evolution of Structural Colors”. In: *Bird Coloration, Volume 1: Mechanisms and*
356 *Measurements*. Ed. by GE Hill and KJ McGraw. Vol. 1. 2 vols. Bird Coloration. Harvard University Press, 2006,
357 p. 640. ISBN: 978-0-674-01893-8.
- 358 [6] DC Osorio and AD Ham. “Spectral Reflectance and Directional Properties of Structural Coloration in Bird Plumage”.
359 In: *Journal of Experimental Biology* 205.14 (2002), pp. 2017–2027. pmid: 12089207.
- 360 [7] D Gomez and M Théry. “Simultaneous Crypsis and Conspicuousness in Color Patterns: Comparative Analysis of
361 a Neotropical Rainforest Bird Community”. In: *The American Naturalist* 169.s1 (2007), S42–S61. DOI: 10.1086/
362 510138.
- 363 [8] JA Endler. “The Color of Light in Forests and Its Implications”. In: *Ecological Monographs* 63.1 (1993), pp. 1–27.
364 DOI: 10.2307/2937121.
- 365 [9] D Gomez and M Théry. “Influence of Ambient Light on the Evolution of Colour Signals: Comparative Analysis of
366 a Neotropical Rainforest Bird Community”. In: *Ecology Letters* 7.4 (2004), pp. 279–284. DOI: 10.1111/j.1461-
367 0248.2004.00584.x.

- 368 [10] J Gröning and A Hochkirch. “Reproductive Interference Between Animal Species”. In: *The Quarterly Review of*
369 *Biology* 83.3 (2008), pp. 257–282. DOI: 10.1086/590510.
- 370 [11] WL Brown and EO Wilson. “Character Displacement”. In: *Systematic Biology* 5.2 (1956), pp. 49–64. DOI: 10.2307/
371 2411924.
- 372 [12] R Butlin. “Speciation by Reinforcement”. In: *Trends in Ecology & Evolution* 2.1 (1987), pp. 8–13. DOI: 10.1016/
373 0169-5347(87)90193-5.
- 374 [13] GF Grether, N Losin, CN Anderson, and K Okamoto. “The Role of Interspecific Interference Competition in Char-
375 acter Displacement and the Evolution of Competitor Recognition”. In: *Biological Reviews* 84.4 (2009), pp. 617–635.
376 DOI: 10.1111/j.1469-185X.2009.00089.x. pmid: 19681844.
- 377 [14] JA Tobias and N Seddon. “Signal Design and Perception in *Hypocnemis* Antbirds: Evidence for Convergent Evolution
378 via Social Selection”. In: *Evolution* 63.12 (2009), pp. 3168–3189. DOI: 10.1111/j.1558-5646.2009.00795.x. pmid:
379 19659594.
- 380 [15] HC Gerhardt. “Reproductive Character Displacement of Female Mate Choice in the Grey Treefrog, *Hyla Chrysoscelis*”.
381 In: *Animal Behaviour* 47.4 (1994), pp. 959–969. DOI: 10.1006/anbe.1994.1127.
- 382 [16] BR Grant and PR Grant. “Songs of Darwin’s Finches Diverge When a New Species Enters the Community”. In:
383 *Proceedings of the National Academy of Sciences* 107.47 (2010), pp. 20156–20163. DOI: 10.1073/pnas.1015115107.
384 pmid: 21048082.
- 385 [17] PMB Bacquet, O Brattström, HL Wang, CE Allen, C Löfstedt, et al. “Selection on Male Sex Pheromone Composition
386 Contributes to Butterfly Reproductive Isolation”. In: *Proceedings of the Royal Society B: Biological Sciences* 282.1804
387 (2015), p. 20142734. DOI: 10.1098/rspb.2014.2734. pmid: 25740889.
- 388 [18] GP Sætre, T Moum, S Bureš, M Král, M Adamjan, et al. “A Sexually Selected Character Displacement in Flycatchers
389 Reinforces Premating Isolation”. In: *Nature* 387.6633 (1997), pp. 589–592. DOI: 10.1038/42451. pmid: 847.
- 390 [19] RE Naisbit, CD Jiggins, and J Mallet. “Disruptive Sexual Selection against Hybrids Contributes to Speciation
391 between *Heliconius Cydno* and *Heliconius Melpomene*”. In: *Proceedings of the Royal Society of London B: Biological*
392 *Sciences* 268.1478 (2001), pp. 1849–1854. DOI: 10.1098/rspb.2001.1753. pmid: 11522205.
- 393 [20] VA Lukhtanov, NP Kandul, JB Plotkin, AV Dantchenko, D Haig, et al. “Reinforcement of Pre-Zygotic Isolation
394 and Karyotype Evolution in *Agrodiaetus* Butterflies”. In: *Nature* 436.7049 (2005), pp. 385–389. DOI: 10.1038/
395 nature03704.
- 396 [21] PR Martin, R Montgomerie, and SC Loughheed. “Color Patterns of Closely Related Bird Species Are More Divergent
397 at Intermediate Levels of Breeding-Range Sympatry”. In: *The American Naturalist* 185.4 (2015), pp. 443–451. DOI:
398 10.1086/680206.
- 399 [22] C Doutrelant, M Paquet, JP Renoult, A Grégoire, PA Crochet, et al. “Worldwide Patterns of Bird Colouration on
400 Islands”. In: *Ecology Letters* 19.5 (2016), pp. 537–545. DOI: 10.1111/ele.12588.

- 401 [23] CR Hemingson, PF Cowman, JR Hodge, and DR Bellwood. “Colour Pattern Divergence in Reef Fish Species Is
402 Rapid and Driven by Both Range Overlap and Symmetry”. In: *Ecology Letters* 22.1 (2019), pp. 190–199. DOI:
403 10.1111/ele.13180.
- 404 [24] ATD Bennett, IC Cuthill, and KJ Norris. “Sexual Selection and the Mismeasure of Color”. In: *The American*
405 *Naturalist* 144.5 (1994), pp. 848–860. DOI: 10.1086/285711.
- 406 [25] IC Cuthill, ATD Bennett, JC Partridge, and EJ Maier. “Plumage Reflectance and the Objective Assessment of Avian
407 Sexual Dichromatism”. In: *The American Naturalist* 153.2 (1999), pp. 183–200. DOI: 10.1086/303160.
- 408 [26] MD Eaton. “Human Vision Fails to Distinguish Widespread Sexual Dichromatism among Sexually “Monochromatic”
409 Birds”. In: *Proceedings of the National Academy of Sciences* 102.31 (2005), pp. 10942–10946. DOI: 10.1073/pnas.
410 0501891102. pmid: 16033870.
- 411 [27] R Montgomerie. “Analyzing Colors”. In: *Bird Coloration, Volume 1: Mechanisms and Measurements*. Ed. by GE Hill
412 and KJ McGraw. Vol. 1. 2 vols. Bird Coloration. Harvard University Press, 2006, p. 640. ISBN: 978-0-674-01893-8.
- 413 [28] JL Parra. “Color Evolution in the Hummingbird Genus *Coeligena*”. In: *Evolution* 64.2 (2010), pp. 324–335. DOI:
414 10.1111/j.1558-5646.2009.00827.x. pmid: 19703221.
- 415 [29] K Delhey. “Revealing The Colourful Side of Birds: Spatial Distribution of Conspicuous Plumage Colours on The
416 Body of Australian Birds”. In: *bioRxiv* (2019), p. 647727. DOI: 10.1101/647727.
- 417 [30] FG Stiles. “Aggressive and Courtship Displays of the Male Anna’s Hummingbird”. In: *The Condor* 84.2 (1982),
418 pp. 208–225. DOI: 10.2307/1367674. JSTOR: 1367674.
- 419 [31] RK Simpson and KJ McGraw. “Experimental Trait Mis-Matches Uncover Specificity of Evolutionary Links between
420 Multiple Signaling Traits and Their Interactions in Hummingbirds”. In: *Evolution* (2018). DOI: 10.1111/evo.13662.
- 421 [32] RK Simpson and KJ McGraw. “Two Ways to Display: Male Hummingbirds Show Different Color-Display Tactics
422 Based on Sun Orientation”. In: *Behavioral Ecology* 29.3 (2018), pp. 637–648. DOI: 10.1093/beheco/ary016.
- 423 [33] BG Hogan and MC Stoddard. “Synchronization of Speed, Sound and Iridescent Color in a Hummingbird Aerial
424 Courtship Dive”. In: *Nature Communications* 9.1 (2018), p. 5260. DOI: 10.1038/s41467-018-07562-7.
- 425 [34] BD Wilts, K Michielsen, J Kuipers, H De Raedt, and DG Stavenga. “Brilliant Camouflage: Photonic Crystals in the
426 Diamond Weevil, *Entimus Imperialis*”. In: *Proceedings of the Royal Society B: Biological Sciences* 279.1738 (2012),
427 pp. 2524–2530. DOI: 10.1098/rspb.2011.2651.
- 428 [35] R Dakin and R Montgomerie. “Eye for an Eyespot: How Iridescent Plumage Ocelli Influence Peacock Mating Success”.
429 In: *Behavioral Ecology* 24.5 (2013), pp. 1048–1057. DOI: 10.1093/beheco/art045.
- 430 [36] S Villéger, NWH Mason, and D Mouillot. “New Multidimensional Functional Diversity Indices for a Multifaceted
431 Framework in Functional Ecology”. In: *Ecology* 89.8 (2008), pp. 2290–2301. DOI: 10.1890/07-1206.1.
- 432 [37] J Del Hoyo, A Elliott, J Sargatal, DA Christie, and E de Juana. *Handbook of the Birds of the World Alive*. 2017.
433 URL: hbw.com.

- 434 [38] J Dorst. “Recherches sur la structure des plumes des trochilidés”. OCLC: 14220401. Paris: Université de Paris, 1951.
435 260 pp.
- 436 [39] H Dürer. “Schillerfarben der Vogelfeder als Evolutionsproblem”. Medizinischen Fakultät der Universität Basel, 1975.
- 437 [40] W Jetz, GH Thomas, JB Joy, K Hartmann, and AO Mooers. “The Global Diversity of Birds in Space and Time”.
438 In: *Nature* 491.7424 (2012), pp. 444–448. DOI: 10.1038/nature11631. pmid: 23123857.
- 439 [41] JA McGuire, CC Witt, JVJ Remsen, A Corl, DL Rabosky, et al. “Molecular Phylogenetics and the Diversification of
440 Hummingbirds”. In: *Current Biology* 24.8 (2014), pp. 910–916. DOI: 10.1016/j.cub.2014.03.016. pmid: 24704078.
- 441 [42] GR Graves and RL Zusi. “An Intergeneric Hybrid Hummingbird (*Heliodoxa Leadbeateri* *Heliangelus Amethysti-*
442 *collis*) from Northern Colombia”. In: *The Condor* 92.3 (1990), pp. 754–760. DOI: 10.2307/1368695.
- 443 [43] FG Stiles and JO Cortés-Herrera. “Diagnosis and Observations of a Hybrid Hummingbird (*Metallura Tyrianthina* x
444 *Aglaiocercus Kingi*) in the Eastern Andes of Colombia”. In: *Revista de la Academia Colombiana de Ciencias Exactas,*
445 *Físicas y Naturales* 39.153 (2015), pp. 481–490. DOI: 10.18257/raccefyn.260.
- 446 [44] SM Doucet and GE Hill. “Do Museum Specimens Accurately Represent Wild Birds? A Case Study of Carotenoid,
447 Melanin, and Structural Colours in Long-Tailed Manakins *Chiroxiphia Linearis*”. In: *Journal of Avian Biology* 40.2
448 (2009), pp. 146–156. DOI: 10.1111/j.1600-048X.2009.03763.x.
- 449 [45] CH Graham, JL Parra, C Rahbek, and JA McGuire. “Phylogenetic Structure in Tropical Hummingbird Com-
450 munities”. In: *Proceedings of the National Academy of Sciences* 106 (Supplement 2 2009), pp. 19673–19678. DOI:
451 10.1073/pnas.0901649106. pmid: 19805042.
- 452 [46] JL Parra, JA McGuire, and CH Graham. “Incorporating Clade Identity in Analyses of Phylogenetic Community
453 Structure: An Example with Hummingbirds.” In: *The American Naturalist* 176.5 (2010), pp. 573–587. DOI: 10.
454 1086/656619. pmid: 20849270.
- 455 [47] RS Ridgely and PJ Greenfield. *The Birds of Ecuador: Status, Distribution and Taxonomy*. Ithaca, NY: Cornell
456 University Press, 2001. 880 pp. ISBN: 978-0-8014-8720-0.
- 457 [48] MG Meadows, NI Morehouse, RL Rutowski, JM Douglas, and KJ McGraw. “Quantifying Iridescent Coloration in
458 Animals: A Method for Improving Repeatability”. In: *Behavioral Ecology and Sociobiology* 65.6 (2011), pp. 1317–
459 1327. DOI: 10.1007/s00265-010-1135-5. pmid: 876.
- 460 [49] DM Chen and TH Goldsmith. “Four Spectral Classes of Cone in the Retinas of Birds”. In: *Journal of Comparative*
461 *Physiology A* 159.4 (1986), pp. 473–479. DOI: 10.1007/BF00604167.
- 462 [50] TE White, RL Dalrymple, DW Noble, JC O’Hanlon, DB Zurek, et al. “Reproducible Research in the Study of
463 Biological Coloration”. In: *Animal Behaviour* 106 (2015), pp. 51–57. DOI: 10.1016/j.anbehav.2015.05.007.
- 464 [51] MG Meadows, TE Roudybush, and KJ McGraw. “Dietary Protein Level Affects Iridescent Coloration in Anna’s
465 Hummingbirds, *Calypte Anna*”. In: *Journal of Experimental Biology* 215.16 (2012), pp. 2742–2750. DOI: 10.1242/
466 jeb.069351. pmid: 22837446.

- 467 [52] H Gruson, C Andraud, W Daney de Marcillac, S Berthier, M Elias, et al. “Quantitative Characterization of Iridescent
468 Colours in Biological Studies: A Novel Method Using Optical Theory”. In: *Interface Focus* 9.1 (2019), p. 20180049.
469 DOI: 10.1098/rsfs.2018.0049.
- 470 [53] R Maia, CM Eliason, PP Bitton, SM Doucet, and MD Shawkey. “Pavo: An R Package for the Analysis, Visualization
471 and Organization of Spectral Data”. In: *Methods in Ecology and Evolution* 4.10 (2013), pp. 906–913. DOI: 10.1111/
472 2041-210X.12069.
- 473 [54] R Maia, H Gruson, JA Endler, and TE White. “Pavo 2: New Tools for the Spectral and Spatial Analysis of Colour
474 in R”. In: *Methods in Ecology and Evolution* 10.7 (2019), pp. 1097–1107. DOI: 10.1111/2041-210X.13174.
- 475 [55] JA Endler and PW Mielke. “Comparing Entire Colour Patterns as Birds See Them”. In: *Biological Journal of the
476 Linnean Society* 86.4 (2005), pp. 405–431. DOI: 10.1111/j.1095-8312.2005.00540.x.
- 477 [56] DC Osorio and M Vorobyev. “A Review of the Evolution of Animal Colour Vision and Visual Communication
478 Signals”. In: *Vision Research*. Vision Research Reviews 48.20 (2008), pp. 2042–2051. DOI: 10.1016/j.visres.2008.
479 06.018. pmid: 18627773.
- 480 [57] A Ödeen and O Håstad. “Complex Distribution of Avian Color Vision Systems Revealed by Sequencing the SWS1
481 Opsin from Total DNA”. In: *Molecular Biology and Evolution* 20.6 (2003), pp. 855–861. DOI: 10.1093/molbev/
482 msg108. pmid: 12716987.
- 483 [58] G Herrera, JC Zagal, M Diaz, MJ Fernández, A Vielma, et al. “Spectral Sensitivities of Photoreceptors and Their Role
484 in Colour Discrimination in the Green-Backed Firecrown Hummingbird (*Sephanoides Sephaniodes*)”. In: *Journal of
485 Comparative Physiology A* 194.9 (2008), p. 785. DOI: 10.1007/s00359-008-0349-8. pmid: 18584181.
- 486 [59] A Ödeen and O Håstad. “Pollinating Birds Differ in Spectral Sensitivity”. In: *Journal of Comparative Physiology A*
487 196.2 (2010), pp. 91–96. DOI: 10.1007/s00359-009-0474-z.
- 488 [60] NS Hart. “Microspectrophotometry of Visual Pigments and Oil Droplets in a Marine Bird, the Wedge-Tailed Shear-
489 water *Puffinus Pacificus*: Topographic Variations in Photoreceptor Spectral Characteristics”. In: *Journal of Exper-
490 imental Biology* 207.7 (2004), pp. 1229–1240. DOI: 10.1242/jeb.00857. pmid: 14978063.
- 491 [61] DF Stotz, JW Fitzpatrick, TA Parker III, and DK Moskovits. *Neotropical Birds: Ecology and Conservation*. Vol. 3.
492 OCLC: 32819832. University of Chicago Press, 1996. ISBN: 978-0-226-77629-3.
- 493 [62] MA Stoffel, S Nakagawa, and H Schielzeth. “rptR: Repeatability Estimation and Variance Decomposition by Gener-
494 alized Linear Mixed-Effects Models”. In: *Methods in Ecology and Evolution* 8.11 (2017). Ed. by S Goslee, pp. 1639–
495 1644. DOI: 10.1111/2041-210X.12797.
- 496 [63] MC Stoddard and RO Prum. “Evolution of Avian Plumage Color in a Tetrahedral Color Space: A Phylogenetic
497 Analysis of New World Buntings.” In: *The American Naturalist* 171.6 (2008), pp. 755–776. DOI: 10.1086/587526.

- 498 [64] M Pagel and F Lutzoni. “Accounting for Phylogenetic Uncertainty in Comparative Studies of Evolution and Adap-
499 tation”. In: *Biological Evolution and Statistical Physics*. Ed. by M Lässig and A Valleriani. Vol. 585. Lecture Notes
500 in Physics. Berlin, Heidelberg: Springer Berlin Heidelberg, 2002. ISBN: 978-3-540-43188-6. DOI: 10.1007/3-540-
501 45692-9.
- 502 [65] OJ Hardy and B Senterre. “Characterizing the Phylogenetic Structure of Communities by an Additive Partitioning of
503 Phylogenetic Diversity”. In: *Journal of Ecology* 95.3 (2007), pp. 493–506. DOI: 10.1111/j.1365-2745.2007.01222.x.
- 504 [66] C Baraloto, OJ Hardy, CET Paine, KG Dexter, C Cruaud, et al. “Using Functional Traits and Phylogenetic Trees
505 to Examine the Assembly of Tropical Tree Communities”. In: *Journal of Ecology* 100.3 (2012), pp. 690–701. DOI:
506 10.1111/j.1365-2745.2012.01966.x.
- 507 [67] OJ Hardy. “Testing the Spatial Phylogenetic Structure of Local Communities: Statistical Performances of Different
508 Null Models and Test Statistics on a Locally Neutral Community”. In: *Journal of Ecology* 96.5 (2008), pp. 914–926.
509 DOI: 10.1111/j.1365-2745.2008.01421.x.
- 510 [68] F de Bello, P Šmilauer, JAF Diniz-Filho, CP Carmona, Z Lososová, et al. “Decoupling Phylogenetic and Functional
511 Diversity to Reveal Hidden Signals in Community Assembly”. In: *Methods in Ecology and Evolution* 8.10 (2017),
512 pp. 1200–1211. DOI: 10.1111/2041-210X.12735.
- 513 [69] VN Balasubramanian, S Chakraborty, and S Panchanathan. “Conformal Predictions for Information Fusion”. In:
514 *Annals of Mathematics and Artificial Intelligence* 74.1-2 (2015), pp. 45–65. DOI: 10.1007/s10472-013-9392-4.
- 515 [70] RK Suarez. “Hummingbird Flight: Sustaining the Highest Mass-Specific Metabolic Rates among Vertebrates”. In:
516 *Experientia* 48.6 (1992), pp. 565–570. DOI: 10.1007/BF01920240.
- 517 [71] DL Altshuler, R Dudley, and JA McGuire. “Resolution of a Paradox: Hummingbird Flight at High Elevation Does
518 Not Come without a Cost”. In: *Proceedings of the National Academy of Sciences* 101.51 (2004), pp. 17731–17736.
519 DOI: 10.1073/pnas.0405260101. pmid: 15598748.
- 520 [72] DL Altshuler, FG Stiles, and R Dudley. “Of Hummingbirds and Helicopters: Hovering Costs, Competitive Ability,
521 and Foraging Strategies.” In: *The American Naturalist* 163.1 (2004), pp. 16–25. DOI: 10.1086/380511. pmid:
522 14767833.
- 523 [73] AB Lindberg and JM Olesen. “The Fragility of Extreme Specialization: *Passiflora Mixta* and Its Pollinating Hum-
524 mingbird *Ensifera Ensifera*”. In: *Journal of Tropical Ecology* 17.2 (2001), pp. 323–329. DOI: 10.1017/S0266467401001213.
- 525 [74] RP Moore, WD Robinson, IJ Lovette, and TR Robinson. “Experimental Evidence for Extreme Dispersal Limitation
526 in Tropical Forest Birds”. In: *Ecology Letters* 11.9 (2008), pp. 960–968. DOI: 10.1111/j.1461-0248.2008.01196.x.
- 527 [75] LN Céspedes, LI Pavan, JA Hazlehurst, and JE Jankowski. “The Behavior and Diet of the Shining Sunbeam (*Aglae-
528 actis Cupripennis*): A Territorial High-Elevation Hummingbird”. In: *The Wilson Journal of Ornithology* 131.1 (2019),
529 pp. 24–34. DOI: 10.1676/18-79.

- 530 [76] RK Simpson and KJ McGraw. “It’s Not Just What You Have, but How You Use It: Solar-Positional and Behavioural
531 Effects on Hummingbird Colour Appearance during Courtship”. In: *Ecology Letters* 0.0 (2018). DOI: 10.1111/ele.
532 13125.
- 533 [77] JA Tobias, R Planqué, DL Cram, and N Seddon. “Species Interactions and the Structure of Complex Communication
534 Networks”. In: *Proceedings of the National Academy of Sciences* 111.3 (2014), pp. 1020–1025. DOI: 10.1073/pnas.
535 1314337111. pmid: 24395769.
- 536 [78] MK McNaught and IPF Owens. “Interspecific Variation in Plumage Colour among Birds: Species Recognition
537 or Light Environment?” In: *Journal of Evolutionary Biology* 15.4 (2002), pp. 505–514. DOI: 10.1046/j.1420-
538 9101.2002.00431.x.
- 539 [79] H Gruson, M Elias, C Andraud, C Djediat, S Berthier, et al. “Hummingbird Iridescence: An Unsuspected Structural
540 Diversity Influences Colouration at Multiple Scales”. In: *bioRxiv* (2019), p. 699744. DOI: 10.1101/699744.
- 541 [80] CJ Clark, TJ Feo, and I Escalante. “Courtship Displays and Natural History of Scintillant (*Selasphorus Scintilla*)
542 and Volcano (*S. Flammula*) Hummingbirds”. In: *The Wilson Journal of Ornithology* 123.2 (2011), pp. 218–228. DOI:
543 10.1676/10-076.1.
- 544 [81] CM Eliason, MD Shawkey, and JA Clarke. “Evolutionary Shifts in the Melanin-Based Color System of Birds”. In:
545 *Evolution* 70.2 (2016), pp. 445–454. DOI: 10.1111/evo.12855. pmid: 26044706.
- 546 [82] P Matyjasiak. “Birds Associate Species-Specific Acoustic and Visual Cues: Recognition of Heterospecific Rivals by
547 Male Blackcaps”. In: *Behavioral Ecology* 16.2 (2005), pp. 467–471. DOI: 10.1093/beheco/ari012.
- 548 [83] D Luther. “The Influence of the Acoustic Community on Songs of Birds in a Neotropical Rain Forest”. In: *Behavioral*
549 *Ecology* 20.4 (2009), pp. 864–871. DOI: 10.1093/beheco/arp074.
- 550 [84] CJ Clark and TJ Feo. “The Anna’s Hummingbird Chirps with Its Tail: A New Mechanism of Sonation in Birds”.
551 In: *Proceedings of the Royal Society of London B: Biological Sciences* 275.1637 (2008), pp. 955–962. DOI: 10.1098/
552 rspb.2007.1619. pmid: 18230592.
- 553 [85] CJ Clark, DO Elias, and RO Prum. “Aeroelastic Flutter Produces Hummingbird Feather Songs”. In: *Science* 333.6048
554 (2011), pp. 1430–1433. DOI: 10.1126/science.1205222. pmid: 21903810.
- 555 [86] CJ Clark. “Wing, Tail, and Vocal Contributions to the Complex Acoustic Signals of Courting Calliope Humming-
556 birds”. In: *Current Zoology* 57.2 (2011), pp. 187–196. DOI: 10.1093/czoolo/57.2.187.
- 557 [87] FG Stiles and LL Wolf. “Ecology and Evolution of Lek Mating Behavior in the Long-Tailed Hermit Hummingbird”.
558 In: *Ornithological Monographs* 27 (1979), pp. iii–78. DOI: 10.2307/40166760. JSTOR: 40166760.
- 559 [88] MA Pizo. “Lek Behavior of the Plovercrest (*Stephanoxis Lalandi*, Trochilidae)”. In: *The Wilson Journal of Ornithol-*
560 *ogy* 124.1 (2012), pp. 106–112. DOI: 10.1676/11-055.1.
- 561 [89] TA Hurly, RD Scott, and SD Healy. “The Function of Displays of Male Rufous Hummingbirds”. In: *The Condor*
562 103.3 (2001), pp. 647–651. DOI: 10.1650/0010-5422(2001)103[0647:TFODOM]2.0.CO;2.

563 [90] S Nakagawa and H Schielzeth. “Repeatability for Gaussian and Non-Gaussian Data: A Practical Guide for Biologists”.
 564 In: *Biological Reviews* 85.4 (2010), pp. 935–956. DOI: 10.1111/j.1469-185X.2010.00141.x. pmid: 20569253.

565 ~~$\tau_{ST} < 0$~~

566 ~~**Phenotypic overdispersion** $\tau_{ST} = 0$~~

567 ~~**No community structure** $\tau_{ST} > 0$~~

568 ~~**Phenotypic clustering** $dc\tau_{ST} < 0$~~

569 ~~**Character displacement (divergence):** co-occurring species are more dissimilar than expected given their phylogenetic
 570 relationships, which means they evolved towards dissimilarity in their colours.—Co-occurring species are less similar
 571 than expected by chance because of character displacement. Co-occurring species are not more neither less similar than
 572 expected by chance despite character displacement because closely related species co-occur more often than expected
 573 at random (phylogenetic clustering; $\Pi_{ST} > 0$). Co-occurring species are more similar than expected by chance despite
 574 character displacement because closely related species co-occur more often than expected at random (phylogenetic clustering; $\Pi_{ST} > 0$).~~

575 ~~**Brownian trait evolution** **Competitive exclusion:** co-occurring species are more dissimilar than expected by chance
 576 because distantly-related (and therefore dissimilar) species co-occur more often than expected at random (phylogenetic overdispersion
 577 species are not more similar nor more different than expected by chance or than predicted given their phylogenetic
 578 relationships. **Environmental filtering:** co-occurring species are more similar than expected by chance because closely-related
 579 (and therefore similar) species co-occur more often than expected at random (phylogenetic clustering; $Pi_{ST} > 0$). $dc\tau_{ST} > 0$~~

580 ~~**Evolutionary convergence** $\tau_{ST} < 0$: co-occurring species are more similar than expected given their phylogenetic relationships,
 581 which means they evolved towards similarity in their colours.—Co-occurring species are less similar than expected by
 582 chance despite evolutionary convergence because distantly-related species co-occur more often than expected at random
 583 (phylogenetic overdispersion; $\Pi_{ST} < 0$).—Co-occurring species are neither more nor less similar than expected by chance
 584 despite evolutionary because distantly-related species co-occur more often than expected at random (phylogenetic overdispersion; $\Pi_{ST} < 0$).
 585 species are more similar than expected by chance because of evolutionary convergence.—Summary of the different
 586 evolutionary and ecological scenarios and their results in terms of values of τ_{ST} and decoupled $dc\tau_{ST}$.~~

Table 3: List of species with their provenance (Confluences = Musée des Confluences, Lyon, France, MNHN = Muséum National d’Histoire Naturelle, Paris, France) and strata. Strata data were extracted from Stotz et al. [61] and used in vision models.


Species	Clade	Provenance	Strata
Adelomyia melanogenys	Coquette	Confluences	Understory
Aglaeactis cupripennis	Brilliant	MNHN	Canopy
Aglaiocercus coelestis	Coquette	MNHN	Canopy

Species	Clade	Provenance	Strata
<i>Aglaiocercus kingi</i> mocoa	Coquette	MNHN	Canopy
<i>Amazilia amabilis</i>	Emerald	MNHN	Understory
<i>Amazilia amazilia</i>	Emerald	MNHN	Understory
<i>Amazilia fimbriata</i> fluviatilis	Emerald	MNHN	Canopy
<i>Amazilia franciae</i>	Emerald	MNHN	Canopy
<i>Amazilia grayi</i> meridionalis	Emerald	MNHN	Canopy
<i>Amazilia rosenbergi</i>	Emerald	MNHN	Understory
<i>Amazilia sapphirina</i>	Emerald	MNHN	Canopy
<i>Amazilia tzacatl</i> jucunda	Emerald	MNHN	Canopy
<i>Androdon aequatorialis</i>	Mangoe	MNHN	Understory
<i>Anthracothorax nigricollis</i>	Mangoe	MNHN	Canopy
<i>Avocettula recurvirostris</i>	Mangoe	Confluences	Understory
<i>Boissonneaua flavescens</i>	Brilliant	MNHN	Canopy
<i>Boissonneaua matthewsii</i>	Brilliant	MNHN	Canopy
<i>Calliphlox amethystina</i>	Bee	MNHN	Canopy
<i>Calliphlox mitchellii</i>	Bee	Confluences	Canopy
<i>Campylopterus falcatus</i>	Emerald	MNHN	Understory
<i>Campylopterus largipennis</i>	Emerald	MNHN	Understory
<i>Campylopterus villaviscensio</i>	Emerald	MNHN	Understory
<i>Chaetocercus bombus</i>	Bee	MNHN	Canopy
<i>Chaetocercus mulsant</i>	Bee	MNHN	Understory
<i>Chalcostigma herrani</i>	Coquette	MNHN	Canopy
<i>Chalcostigma ruficeps</i>	Coquette	Confluences	Understory
<i>Chalcostigma stanleyi</i> stanleyi	Coquette	MNHN	Canopy
<i>Chalybura buffonii</i> intermedia	Emerald	Confluences	Understory
<i>Chalybura urochrysia</i> urochrysia	Emerald	Confluences	Understory
<i>Chlorestes notata</i> obsoletus-puruensis	Emerald	Confluences	Canopy
<i>Chlorostilbon melanorhynchus</i>	Emerald	MNHN	Understory
<i>Chlorostilbon mellisugus</i> phoeopygus	Emerald	Confluences	Understory
<i>Chrysuronia oenone</i>	Emerald	MNHN	Canopy
<i>Coeligena coeligena</i>	Brilliant	MNHN	Understory
<i>Coeligena iris</i> hesperus	Brilliant	MNHN	Understory
<i>Coeligena iris</i> iris	Brilliant	MNHN	Understory
<i>Coeligena lutetiae</i>	Brilliant	MNHN	Understory


Species	Clade	Provenance	Strata
<i>Coeligena torquata fulgidigula</i>	Brilliant	MNHN	Understory
<i>Coeligena torquata torquata</i>	Brilliant	MNHN	Understory
<i>Coeligena wilsoni</i>	Brilliant	MNHN	Understory
<i>Colibri coruscans</i>	Mangoe	MNHN	Canopy
<i>Colibri delphinae</i>	Mangoe	MNHN	Canopy
<i>Colibri thalassinus</i>	Mangoe	MNHN	Canopy
<i>Damophila julie</i>	Emerald	MNHN	Understory
<i>Discosura conversii</i>	Coquette	MNHN	Canopy
<i>Discosura langsdorffi</i>	Coquette	Confluences	Canopy
<i>Discosura popelairii</i>	Coquette	MNHN	Canopy
<i>Doryfera johannae</i>	Mangoe	MNHN	Understory
<i>Doryfera ludovicae</i>	Mangoe	MNHN	Understory
<i>Ensifera ensifera</i>	Brilliant	MNHN	Understory
<i>Eriocnemis alinae</i>	Brilliant	MNHN	Understory
<i>Eriocnemis luciani</i>	Brilliant	MNHN	Understory
<i>Eriocnemis mosquera</i>	Brilliant	Confluences	Understory
<i>Eriocnemis nigrivestis</i>	Brilliant	MNHN	Understory
<i>Eriocnemis vestita smaragdinicollis</i>	Brilliant	MNHN	Understory
<i>Eutoxeres aquila</i>	Hermit	MNHN	Understory
<i>Eutoxeres condamini</i>	Hermit	Confluences	Understory
<i>Florisuga mellivora</i>	Topazes	MNHN	Canopy
<i>Glaucis aeneus</i>	Hermit	MNHN	Understory
<i>Glaucis hirsutus affinis</i>	Hermit	MNHN	Understory
<i>Haplophaedia aureliae russata</i>	Brilliant	Confluences	Understory
<i>Haplophaedia lugens</i>	Brilliant	Confluences	Understory
<i>Heliangelus amethysticollis laticlavus</i>	Coquette	Confluences	Understory
<i>Heliangelus exortis</i>	Coquette	MNHN	Understory
<i>Heliangelus exortis</i>	Coquette	MNHN	Understory
<i>Heliangelus micraster</i>	Coquette	MNHN	Understory
<i>Heliangelus strophianus</i>	Coquette	MNHN	Understory
<i>Heliangelus viola</i>	Coquette	MNHN	Understory
<i>Heliodoxa aurescens</i>	Brilliant	MNHN	Understory
<i>Heliodoxa imperatrix</i>	Brilliant	MNHN	Understory
<i>Heliodoxa jacula jamesoni</i>	Brilliant	MNHN	Understory

Species	Clade	Provenance	Strata
<i>Heliodoxa leadbeateri</i>	Brilliant	MNHN	Understory
<i>Heliodoxa rubinoides aequatorialis</i>	Brilliant	MNHN	Understory
<i>Heliodoxa schreibersii</i>	Brilliant	MNHN	Understory
<i>Heliomaster longirostris</i>	MtGem	MNHN	Canopy
<i>Heliothyx auritus</i>	Mangoe	MNHN	Canopy
<i>Heliothyx barroti</i>	Mangoe	MNHN	Canopy
<i>Klais guimeti</i>	Emerald	MNHN	Understory
<i>Lafresnaya lafresnayi gayi</i>	Brilliant	Confluences	Understory
<i>Lesbia nuna gracilis</i>	Coquette	MNHN	Canopy
<i>Leucippus baeri</i>	Emerald	Confluences	Understory
<i>Leucippus chlorocercus</i>	Emerald	Confluences	Canopy
<i>Lophornis chalybeus verreauxi</i>	Coquette	MNHN	Canopy
<i>Metallura baroni</i>	Coquette	MNHN	Canopy
<i>Metallura tyrianthina tyrianthina</i>	Coquette	MNHN	Understory
<i>Metallura williamsi primolina</i>	Coquette	MNHN	Canopy
<i>Myrmia micrura</i>	Bee	MNHN	Canopy
<i>Ocreatus underwoodii melanantherus</i>	Brilliant	MNHN	Understory
<i>Opisthoprora euryptera</i>	Coquette	Confluences	Understory
<i>Oreotrochilus chimborazo chimborazo</i>	Coquette	MNHN	Understory
<i>Oreotrochilus chimborazo jamesonii</i>	Coquette	MNHN	Understory
<i>Patagona gigas</i>	Patagona	MNHN	Canopy
<i>Phaethornis atrimentalis atrimentalis</i>	Hermit	Confluences	Understory
<i>Phaethornis bourcierii</i>	Hermit	MNHN	Understory
<i>Phaethornis griseogularis</i>	Hermit	MNHN	Understory
<i>Phaethornis griseogularis</i>	Hermit	MNHN	Understory
<i>Phaethornis guy</i>	Hermit	MNHN	Understory
<i>Phaethornis hispidus</i>	Hermit	Confluences	Understory
<i>Phaethornis longirostris</i>	Hermit	Confluences	Understory
<i>Phaethornis malaris</i>	Hermit	Confluences	Understory
<i>Phaethornis ruber</i>	Hermit	Confluences	Understory
<i>Phaethornis syrmatophorus columbianus</i>	Hermit	MNHN	Understory
<i>Phaethornis yaruqui yaruqui</i>	Hermit	MNHN	Understory
<i>Phlogophilus hemileucurus</i>	Coquette	MNHN	Understory
<i>Polytmus theresiae leucorrhous</i>	Mangoe	MNHN	Understory


Species	Clade	Provenance	Strata
<i>Pterophanes cyanopterus</i>	Brilliant	MNHN	Understory
<i>Ramphomicron microrhynchum</i>	Coquette	MNHN	Canopy
<i>Schistes geoffroyi</i>	Mangoe	MNHN	Understory
<i>Taphrospilus hypostictus</i>	Emerald	MNHN	Understory
<i>Thalurania fannyi verticeps</i>	Emerald	MNHN	Understory
<i>Thalurania furcata viridipectus</i>	Emerald	MNHN	Understory
<i>Thaumastura cora</i>	Bee	Confluences	Canopy
<i>Threnetes leucurus cervinicauda</i>	Hermit	Confluences	Understory
<i>Threnetes ruckeri</i>	Hermit	MNHN	Understory
<i>Urochroa bougueri</i>	Brilliant	Confluences	Understory
<i>Urochroa bougueri leucura</i>	Brilliant	Confluences	Understory
<i>Urosticte benjamini</i>	Brilliant	MNHN	Understory
<i>Urosticte ruficrissa</i>	Brilliant	Confluences	Understory



Supplementary figure 1: Locations and names of the 8 patches measured on all species. Additional patches were measured for each species as soon as they differed from one of the 8 patches listed here for a human observer, as detailed in the methods section and as in Gomez and Théry [7].



Supplementary figure 2: Phylogenetic coverage of the *Trochilidae* family in our dataset (species and lineages in red).



Supplementary figure 3: Study sites locations (red dots) plotted on an altitudinal map of Ecuador. Communities outside the borders of the map are on islands or close enough to Ecuador borders to be taken into account in our study.





Figure 1: (a) community total colour volume and (b) brightness range increase with the number of species within the community. Each point is a community. The black solid line represents the mean value of (a) colour volume or (b) brightness range from 10 000 random communities with a given species count (null model) and the gray ribbon represents two standard deviations from the mean of the null model.

Variable	Diffuse		Directional		Both		
	R	p-value	<u>R</u>	<u>p-value</u>	<u>R</u>	<u>p-value</u>	
Hue	x	<u>0.734</u>	<u>0.002</u>	<u>0.877</u>	<u><0.0001</u>	0.925	<0.0001
	y	<u>0.923</u>	<u><0.0001</u>	<u>0.785</u>	<u>0.0006</u>	0.951	<0.0001
	z	<u>0.780</u>	<u>0.0006</u>	<u>0.880</u>	<u><0.0001</u>	0.940	<0.0001
Brightness	<u>0.411</u>	<u>0.090</u>	<u>0.055</u>	<u>0.48</u>	0.373	0.04	

Supplementary table 1: We quantified the repeatability R (intra-class coefficient ICC) and the related p-value by bootstrapping using the `rptR` R package [90] of indices used in this study by performing the same measurements twice on two patches for 12 species (*Coeligena torquata*, *Colibri coruscans*, *Doryfera ludovicae*, *Heliangelus strophianus*, *Heliodoxa jamesonii*, *Heliostyris barroti*, *Juliamyia julie*, *Lesbia nuna*, *Metallura tyrianthina*, *Ramphomicron microrhynchum*, *Schistes albogularis*, *Urosticte benjamini*). Patches were selected to be of similar hue from a human point of view.



Supplementary figure 4: [Colour of the 8 main patches for each species in our dataset. The colour corresponds to the colour in the human visual system \(CIE10\). The x-axis on the phylogeny is in millions years.](#)

variable	value	Crown	Back	Rump	Tail	Throat	Breast	Belly	Wing
Hue	τ_{st}	-0.0073	0.055	0.055	0.044	0.027	0.03	0.05	0.058
	$p_{\tau_{st}<0}$	0.4	1	1	1	0.9	0.9	1	1
	$p_{\tau_{st}>0}$	0.6	0.01	0.01	0.03	0.09	0.06	0.005	0.006
	$d\mathcal{T}_{st}$	0.0099	0.026	-0.0021	0.0034	-0.0021	-0.0032	-0.01	0.00073
	$p_{\tau_{st}<0}$	1	1	0.8	1	0.9	0.3	<0.0001	1
	$p_{\tau_{st}>0}$	<0.0001	<0.0001	1	0.2	1	1	1	1
Brightness	τ_{st}	-0.021	0.0078	0.0032	-0.0064	0.00015	0.0041	-0.0031	0.0091
	$p_{\tau_{st}<0}$	0.1	0.7	0.6	0.5	0.5	0.6	0.5	0.6
	$p_{\tau_{st}>0}$	0.9	0.3	0.4	0.5	0.5	0.4	0.5	0.4
	$d\mathcal{T}_{st}$	-0.0014	0.0028	0.00037	0.00068	0.013	0.023	0.007	-0.0058
	$p_{\tau_{st}<0}$	0.3	1	0.9	1	1	1	1	0.2
	$p_{\tau_{st}>0}$	0.8	0.7	0.7	0.8	<0.0001	<0.0001	0.002	1
Hue shift	τ_{st}	-0.007	0.051	0.052	0.043	0.027	0.029	0.049	0.058
	$p_{\tau_{st}<0}$	0.4	1	1	1	0.9	0.9	1	1
	$p_{\tau_{st}>0}$	0.6	0.01	0.01	0.03	0.08	0.06	0.006	0.006
	$d\mathcal{T}_{st}$	0.0087	0.0059	-0.0068	-0.006	-0.0033	0.0023	-0.0098	-0.0018
	$p_{\tau_{st}<0}$	1	1	0.005	0.01	0.6	1	<0.0001	1
	$p_{\tau_{st}>0}$	<0.0001	0.03	1	1	1	0.9	1	1

Supplementary table 2: Numerical values for τ_{st} and decoupled τ_{st} (denoted $d\mathcal{T}_{st}$). P-values were computed by comparison of the actual value with the null distribution (obtained by randomisation of the communities using method 1s of Hardy [67]). Significant p-values are in bold and green. Positive values of $d\mathcal{T}_{st}$ indicate phenotypic clustering whereas negative values indicate overdispersion.